

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All causes

All races 3/, both sexes

2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	873.1	728.7	32.9	16.4	20.9	68.2	96.0	99.0	116.3	162.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8

All races 3/, male

2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	874.7	799.9	36.5	18.3	25.0	94.9	142.0	141.9	157.3	209.8
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6

All races 3/, female

2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	871.6	654.3	29.1	14.5	16.6	40.0	48.2	56.5	76.0	115.1
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All causes

White, both sexes

2003.....	890.1	591.3	28.5	13.7	17.8	64.7	89.4	86.1	101.9	145.5
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	915.5	598.4	29.1	15.1	19.8	65.2	86.6	87.9	103.6	146.3
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5

White, male

2003.....	877.6	658.7	31.5	15.3	21.4	88.3	129.1	120.0	137.0	187.6
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	905.8	656.2	32.5	16.9	23.8	89.2	127.5	125.8	142.1	191.6
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8

White, female

2003.....	902.3	520.7	25.5	12.1	14.0	39.6	46.7	50.0	65.2	102.2
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	924.9	537.9	25.4	13.3	15.6	39.7	43.6	49.6	64.9	100.5
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All causes

Black, both sexes

2003.....	763.6	1,273.8	46.8	19.7	25.7	80.0	147.3	168.6	194.1	275.2
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	809.6	1,505.6	56.1	24.5	28.0	89.0	155.7	175.6	214.9	288.5
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5

Black, male

2003.....	813.7	1,410.0	53.7	21.6	31.4	120.9	224.5	249.6	262.5	344.8
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	865.4	1,653.2	61.2	27.5	33.7	131.6	237.6	255.6	287.9	368.1
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8

Black, female

2003.....	717.9	1,132.2	39.7	17.8	19.8	37.8	70.6	93.6	132.5	213.1
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	759.1	1,352.7	50.8	21.4	22.1	44.8	75.4	102.2	150.2	217.8
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.8	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

2003.....	1.3	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

All races 3/, male

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

All races 3/, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*

Black, male

2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*

Black, female

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	1.0	0.2	*	*	0.1	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Septicemia (A40-A41)

All races 3/, both sexes

2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.3	7.1	0.7	0.2	0.1	0.2	0.3	0.6	0.9	1.5
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4

All races 3/, male

2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	10.1	7.1	0.7	0.2	*	0.2	0.4	0.7	1.0	1.5
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5

All races 3/, female

2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.6	7.2	0.6	*	*	*	0.3	0.5	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Septicemia (A40-A41)

White, both sexes

2003.....	11.5	4.6	0.4	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	11.0	5.6	0.5	0.2	0.1	0.2	0.2	0.4	0.7	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0

White, male

2003.....	10.2	4.6	0.6	*	*	0.3	0.4	0.4	0.8	1.3
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.6	5.7	0.6	*	*	*	0.3	0.5	0.8	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2

White, female

2003.....	12.8	4.5	*	*	*	*	0.4	0.6	0.7	1.1
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.2	5.5	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Septicemia (A40-A41)

Black, both sexes

2003.....	16.3	18.6	0.9	*	*	*	0.7	1.6	2.2	3.6
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.8	17.3	1.2	*	*	*	0.8	1.5	2.6	3.8
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7

Black, male

2003.....	15.3	21.1	*	*	*	*	*	1.6	2.7	3.6
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	15.4	16.9	*	*	*	*	*	2.2	2.6	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0

Black, female

2003.....	17.2	16.1	*	*	*	*	*	1.6	1.6	3.5
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	18.0	17.8	*	*	*	*	*	*	2.6	4.0
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Viral hepatitis (B15-B19)

All races 3/, both sexes

2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

All races 3/, male

2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.5	*	*	*	*	*	*	*	0.3	1.4
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2

All races 3/, female

2003.....	1.3	*	*	*	*	*	*	*	*	0.4
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Viral hepatitis (B15-B19)

White, both sexes

2003.....	1.8	*	*	*	*	*	*	*	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.9	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

White, male

2003.....	2.4	*	*	*	*	*	*	*	0.2	1.2
2002.....	2.6	*	*	*	*	*	*	*	0.4	1.5
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.3

White, female

2003.....	1.2	*	*	*	*	*	*	*	*	0.5
2002.....	1.3	*	*	*	*	*	*	*	*	0.5
2001.....	1.3	*	*	*	*	*	*	*	*	0.6
2000.....	1.3	*	*	*	*	*	*	*	*	0.6
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.3	*	*	*	0.1	0.2	0.8	3.6	9.1	13.0
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0

All races 3/, male

2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	8.1	*	*	*	*	*	0.8	4.2	12.7	19.6
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5

All races 3/, female

2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.6	*	*	*	*	*	0.7	3.1	5.7	6.5
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2003.....	2.5	*	*	*	*	*	0.3	0.9	2.9	6.1
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.9	*	*	*	*	*	0.3	1.7	4.8	7.7
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9

White, male

2003.....	4.2	*	*	*	*	*	0.3	1.4	4.5	9.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.8	*	*	*	*	*	0.4	2.2	7.5	12.7
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0

White, female

2003.....	0.9	*	*	*	*	*	*	0.5	1.3	2.4
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.1	2.0	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2003.....	19.6	*	*	*	*	0.6	3.2	11.2	23.2	46.0
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	22.2	*	*	*	*	0.7	3.7	15.8	38.6	50.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9

Black, male

2003.....	27.7	*	*	*	*	*	3.2	12.7	28.6	61.9
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	32.2	*	*	*	*	*	3.6	17.0	50.2	72.5
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9

Black, female

2003.....	12.3	*	*	*	*	*	3.1	9.8	18.3	31.9
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	13.2	*	*	*	*	*	3.8	14.8	28.3	31.5
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2003.....	2.6	4.3	0.5	0.1	0.1	0.2	0.2	0.3	0.4	0.7
2002.....	2.3	5.6	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7
2000.....	2.0	4.3	0.5	0.2	0.2	0.2	0.3	0.3	0.4	0.7
1999.....	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0

All races 3/, male

2003.....	2.8	4.2	0.5	*	*	0.3	0.2	0.3	0.5	0.9
2002.....	2.6	6.1	0.5	*	0.2	0.2	0.3	0.4	0.5	0.9
2001.....	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8
2000.....	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.5	0.9
1999.....	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3

All races 3/, female

2003.....	2.3	4.5	0.5	*	*	*	*	0.3	0.3	0.6
2002.....	2.1	5.0	0.3	0.3	*	0.2	0.2	0.3	0.4	0.6
2001.....	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6
2000.....	1.9	3.9	0.4	*	*	*	0.3	0.3	0.4	0.5
1999.....	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2003.....	2.6	3.5	0.4	0.1	*	0.2	0.2	0.2	0.4	0.6
2002.....	2.4	5.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.6
2001.....	2.0	3.8	0.4	0.2	0.1	0.2	0.2	0.3	0.3	0.6
2000.....	2.0	3.3	0.4	0.2	0.2	0.2	0.3	0.2	0.3	0.6
1999.....	2.5	4.5	0.5	0.2	0.2	0.2	0.2	0.4	0.5	0.8

White, male

2003.....	2.8	3.2	0.4	*	*	0.3	*	0.3	0.4	0.8
2002.....	2.6	5.3	0.5	*	0.2	*	*	0.3	0.5	0.7
2001.....	2.1	3.9	0.4	*	*	*	*	0.3	0.4	0.7
2000.....	2.1	3.8	0.4	*	*	*	0.3	*	0.4	0.7
1999.....	2.6	5.6	0.6	*	*	*	*	0.3	0.6	1.2

White, female

2003.....	2.4	3.9	0.5	*	*	*	*	*	0.3	0.5
2002.....	2.2	4.6	*	*	*	*	*	0.3	0.4	0.4
2001.....	1.9	3.8	*	*	*	*	*	*	0.2	0.5
2000.....	1.9	2.8	0.4	*	*	*	*	*	0.3	0.4
1999.....	2.5	3.4	*	*	*	*	*	0.4	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2003.....	2.5	8.0	1.0	*	*	*	*	*	0.7	1.4
2002.....	2.5	8.7	*	*	*	*	*	*	1.0	1.8
2001.....	2.2	9.5	*	*	*	*	*	*	1.0	1.4
2000.....	2.3	10.5	*	*	*	*	*	*	1.3	1.7
1999.....	3.5	9.9	0.9	*	*	*	*	1.1	2.1	2.5

Black, male

2003.....	2.9	8.9	*	*	*	*	*	*	*	1.6
2002.....	2.9	9.6	*	*	*	*	*	*	*	2.1
2001.....	2.4	10.8	*	*	*	*	*	*	*	1.7
2000.....	2.6	10.5	*	*	*	*	*	*	*	2.2
1999.....	3.6	10.1	*	*	*	*	*	*	2.6	2.7

Black, female

2003.....	2.2	7.1	*	*	*	*	*	*	*	*
2002.....	2.1	7.9	*	*	*	*	*	*	*	1.6
2001.....	2.0	8.2	*	*	*	*	*	*	*	*
2000.....	2.1	10.5	*	*	*	*	*	*	*	*
1999.....	3.4	9.7	*	*	*	*	*	*	1.6	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	200.9	2.4	2.8	2.5	2.6	3.7	5.2	7.6	13.1	25.0
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2

All races 3/, male

2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	212.5	2.5	3.0	2.7	2.9	4.4	6.1	7.8	12.0	21.7
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6

All races 3/, female

2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	189.9	2.2	2.5	2.3	2.3	3.1	4.3	7.4	14.1	28.3
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms (C00-C97)

White, both sexes

2003.....	203.7	2.1	2.6	2.8	2.6	3.5	4.7	6.7	11.3	22.2
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	212.2	2.5	2.7	2.5	2.7	3.7	5.1	7.6	12.7	23.7
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0

White, male

2003.....	213.1	1.8	2.9	3.1	2.7	4.3	5.1	7.0	10.6	19.3
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	222.5	2.7	3.1	2.7	3.0	4.2	6.0	7.9	11.9	20.8
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1

White, female

2003.....	194.6	2.3	2.3	2.4	2.5	2.6	4.2	6.5	12.0	25.2
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	202.2	2.2	2.4	2.4	2.4	3.0	4.2	7.4	13.4	26.6
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms (C00-C97)

Black, both sexes

2003.....	164.3	*	2.2	2.3	2.8	2.9	4.9	8.5	15.7	31.0
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	175.5	*	3.1	2.3	2.5	4.5	5.9	8.3	16.4	35.1
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8

Black, male

2003.....	178.3	*	2.4	2.4	3.1	3.5	5.6	7.0	13.5	21.7
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	195.6	*	3.0	2.7	2.8	5.3	7.0	7.6	13.5	28.1
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8

Black, female

2003.....	151.4	*	2.1	2.1	2.5	2.3	4.3	10.0	17.7	39.2
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	157.2	*	3.3	2.0	2.1	3.6	4.8	9.0	18.9	41.4
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	2.7	*	*	*	*	*	*	0.2	0.3
2002.....	2.7	*	*	*	*	*	0.1	0.2	0.4
2001.....	2.7	*	*	*	*	*	*	0.2	0.4
2000.....	2.7	*	*	*	*	*	*	0.2	0.4
1999.....	2.7	*	*	*	*	*	*	0.1	0.4

[illegible]

2003.....	1.7	*	*	*	*	*	*	*	*
2002.....	1.7	*	*	*	*	*	*	*	*
2001.....	1.8	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.8	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

[illegible]

2003.....	3.7	*	*	*	*	*	*	*	*
2002.....	3.7	*	*	*	*	*	*	*	*
2001.....	3.6	*	*	*	*	*	*	*	*
2000.....	3.5	*	*	*	*	*	*	*	*
1999.....	3.5	*	*	*	*	*	*	*	*

2003.....	1.8	*	*	*	*	*	*	*	*
2002.....	1.8	*	*	*	*	*	*	*	*
2001.....	1.9	*	*	*	*	*	*	*	*
2000.....	1.9	*	*	*	*	*	*	*	0.2
1999.....	1.9	*	*	*	*	*	*	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	2.9	*	*	*	*	*	*	*	*
2002.....	2.8	*	*	*	*	*	*	*	0.8
2001.....	2.9	*	*	*	*	*	*	*	0.8
2000.....	3.1	*	*	*	*	*	*	*	*
1999.....	3.2	*	*	*	*	*	*	*	0.8

2003.....	4.6	*	*	*	*	*	*	*	*
2002.....	4.3	*	*	*	*	*	*	*	*
2001.....	4.4	*	*	*	*	*	*	*	*
2000.....	5.0	*	*	*	*	*	*	*	*
1999.....	4.8	*	*	*	*	*	*	*	*

2003.....	1.4	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*
1999.....	1.7	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.6	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8

All races 3/, male

2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.4	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9

All races 3/, female

2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.8	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	5.4	*	*	*	*	*	*	*
2002.....	5.6	*	*	*	*	*	*	0.7
2001.....	5.6	*	*	*	*	*	*	0.8
2000.....	6.2	*	*	*	*	*	*	0.9
1999.....	6.1	*	*	*	*	*	*	0.8

2003.....	6.5	*	*	*	*	*	*	*	*
2002.....	6.6	*	*	*	*	*	*	*	*
2001.....	6.7	*	*	*	*	*	*	*	1.8
2000.....	7.3	*	*	*	*	*	*	*	1.9
1999.....	7.3	*	*	*	*	*	*	*	1.4

2003.....	4.3	*	*	*	*	*	*	*	*
2002.....	4.7	*	*	*	*	*	*	*	*
2001.....	4.5	*	*	*	*	*	*	*	*
2000.....	5.2	*	*	*	*	*	*	*	*
1999.....	5.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.9	*	*	*	*	*	0.1	0.4	1.0	2.2
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9

All races 3/, male

2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	21.1	*	*	*	*	*	*	0.5	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0

All races 3/, female

2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.6	*	*	*	*	*	*	0.4	0.8	1.9
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2003.....	20.2	*	*	*	*	*	0.1	0.5	0.9	1.9
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.9	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8

White, male

2003.....	20.5	*	*	*	*	*	*	0.6	0.9	2.0
2002.....	21.1	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	21.2	*	*	*	*	*	*	0.4	1.1	2.5
2000.....	22.0	*	*	*	*	*	*	0.5	1.1	2.2
1999.....	21.8	*	*	*	*	*	*	0.5	0.9	1.9

White, female

2003.....	19.8	*	*	*	*	*	*	0.4	0.9	1.8
2002.....	20.3	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	20.7	*	*	*	*	*	*	0.4	0.9	1.8
2000.....	21.7	*	*	*	*	*	*	0.4	0.7	1.6
1999.....	21.4	*	*	*	*	*	*	0.3	0.9	1.7

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	5.1	*	0.1	*	*	*	0.1	0.1	0.3	0.5
2002.....	4.9	*	0.2	*	*	*	*	0.2	0.3	0.6
2001.....	4.7	*	*	*	*	*	*	0.2	0.2	0.6
2000.....	4.7	*	0.2	*	*	*	*	0.2	0.3	0.5
1999.....	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6

2003.....	6.6	*	*	*	*	*	*	*	0.4	0.8
2002.....	6.4	*	*	*	*	*	*	0.2	0.4	0.8
2001.....	6.1	*	*	*	*	*	*	*	0.3	0.9
2000.....	6.1	*	*	*	*	*	*	*	0.4	0.7
1999.....	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	4.9	*	*	*	*	*	*	*	0.2	0.4
2002.....	4.8	*	0.2	*	*	*	*	*	0.2	0.4
2001.....	4.6	*	*	*	*	*	*	*	0.2	0.5
2000.....	4.6	*	*	*	*	*	*	*	0.2	0.4
1999.....	4.4	*	0.2	*	*	*	*	*	*	0.4

[illegible]

2003.....	3.5	*	*	*	*	*	*	*
2002.....	3.5	*	*	*	*	*	*	0.2
2001.....	3.3	*	*	*	*	*	*	0.3
2000.....	3.4	*	*	*	*	*	*	0.3
1999.....	3.3	*	*	*	*	*	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes

2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.7	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8

All races 3/, male

2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.6	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9

All races 3/, female

2003.....	10.6	*	*	*	*	*	*	*	*	0.5
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.7	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	56.5	*	*	*	*	*	*	0.2	0.8	3.0
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9

All races 3/, male

2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	67.2	*	*	*	*	*	*	0.3	0.8	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8

All races 3/, female

2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	46.3	*	*	*	*	*	*	*	0.9	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, both sexes

2003.....	58.7	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	60.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7

White, male

2003.....	66.9	*	*	*	*	*	*	*	0.6	2.3
2002.....	67.7	*	*	*	*	*	*	*	0.7	2.5
2001.....	68.6	*	*	*	*	*	*	*	0.6	2.8
2000.....	70.8	*	*	*	*	*	*	0.3	0.8	3.0
1999.....	69.1	*	*	*	*	*	*	*	0.8	2.6

White, female

2003.....	50.7	*	*	*	*	*	*	*	0.6	2.4
2002.....	50.7	*	*	*	*	*	*	*	0.5	2.8
2001.....	49.8	*	*	*	*	*	*	*	0.6	3.0
2000.....	50.6	*	*	*	*	*	*	*	0.8	2.5
1999.....	48.2	*	*	*	*	*	*	*	0.6	2.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant melanoma of skin (C43)

All races 3/, both sexes

2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.7	*	*	*	*	*	0.2	0.3	0.7	1.1
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0

All races 3/, male

2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.4	*	*	*	*	*	*	0.4	0.8	1.1
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3

All races 3/, female

2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant melanoma of skin (C43)

White, both sexes

2003.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.3
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3

White, male

2003.....	4.2	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	4.0	*	*	*	*	*	*	0.5	1.0	1.3
2001.....	4.1	*	*	*	*	*	*	0.5	0.7	1.4
2000.....	4.1	*	*	*	*	*	*	0.5	1.0	1.3
1999.....	4.0	*	*	*	*	*	*	0.5	0.8	1.6

White, female

2003.....	2.3	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	2.3	*	*	*	*	*	*	0.4	0.5	1.1
2001.....	2.3	*	*	*	*	*	*	0.4	0.6	0.9
2000.....	2.4	*	*	*	*	*	*	0.3	0.6	1.2
1999.....	2.3	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.4	*	*	*	*	*	*	0.5	2.0	4.4
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3

All races 3/, male

2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

All races 3/, female

2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.8	*	*	*	*	*	*	0.9	3.9	8.7
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of breast (C50)

White, both sexes

2003.....	15.0	*	*	*	*	*	*	0.2	1.5	3.8
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	16.0	*	*	*	*	*	*	0.4	1.7	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7

White, male

2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

White, female

2003.....	29.3	*	*	*	*	*	*	0.4	3.0	7.7
2002.....	29.5	*	*	*	*	*	*	0.5	3.0	7.2
2001.....	29.8	*	*	*	*	*	*	0.8	3.0	7.8
2000.....	31.1	*	*	*	*	*	*	0.9	3.4	7.8
1999.....	30.2	*	*	*	*	*	*	0.8	3.0	7.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of breast (C50)

Black, both sexes

2003.....	15.1	*	*	*	*	*	*	0.9	3.4	8.0
2002.....	14.9	*	*	*	*	*	*	0.8	3.7	8.1
2001.....	15.0	*	*	*	*	*	*	1.1	4.0	7.5
2000.....	15.4	*	*	*	*	*	*	0.8	3.9	7.9
1999.....	15.0	*	*	*	*	*	*	0.9	3.7	8.8

Black, male

2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*

Black, female

2003.....	28.5	*	*	*	*	*	*	1.6	6.5	15.1
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	28.9	*	*	*	*	*	*	*	7.3	14.6
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.8	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	3.0	*	*	*	*	*	*	0.9	1.6	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2003.....	1.2	*	*	*	*	*	*	0.3	0.6	1.1
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	2.5	*	*	*	*	*	*	0.5	1.2	2.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.5	2.4
2001.....	2.7	*	*	*	*	*	*	0.6	1.5	2.5
2000.....	2.7	*	*	*	*	*	*	0.9	1.4	2.5
1999.....	2.8	*	*	*	*	*	*	0.7	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2003.....	2.1	*	*	*	*	*	*	*	1.2	2.0
2002.....	2.2	*	*	*	*	*	*	*	1.1	2.1
2001.....	2.1	*	*	*	*	*	*	*	1.1	1.8
2000.....	2.5	*	*	*	*	*	*	*	1.5	2.1
1999.....	2.3	*	*	*	*	*	*	*	1.0	2.1

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	4.0	*	*	*	*	*	*	*	2.2	3.8
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.7	*	*	*	*	*	*	*	2.8	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

All races 3/, male

All races 3/, female

2003.....	4.7	*	*	*	*	*	*	*	0.3
2002.....	4.7	*	*	*	*	*	*	0.2	0.4
2001.....	4.7	*	*	*	*	*	*	0.3	0.4
2000.....	4.7	*	*	*	*	*	*	0.3	0.4
1999.....	4.5	*	*	*	*	*	*	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	2.4	*	*	*	*	*	*	*	*
2002.....	2.4	*	*	*	*	*	*	*	*
2001.....	2.4	*	*	*	*	*	*	*	0.1
2000.....	2.4	*	*	*	*	*	*	*	*
1999.....	2.4	*	*	*	*	*	*	*	*

[illegible]

2003.....	4.7	*	*	*	*	*	*	*	*
2002.....	4.7	*	*	*	*	*	*	*	*
2001.....	4.8	*	*	*	*	*	*	0.3	*
2000.....	4.7	*	*	*	*	*	*	*	*
1999.....	4.6	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of ovary (C56)

All races 3/, both sexes

2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	10.0	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of ovary (C56)

White, both sexes

2003.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	11.0	*	*	*	*	*	*	0.3	0.6	1.2
2002.....	11.1	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	11.0	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	11.1	*	*	*	*	*	*	0.5	0.7	1.3
1999.....	10.6	*	*	*	*	*	*	0.3	0.5	1.5

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	10.2	*	*	*	*	*	*	*	*	*
2002.....	10.6	*	*	*	*	*	*	*	*	*
2001.....	10.8	*	*	*	*	*	*	*	*	*
2000.....	11.3	*	*	*	*	*	*	*	*	*
1999.....	11.4	*	*	*	*	*	*	*	*	*

All races 3/, male

2003.....	20.7	*	*	*	*	*	*	*	*	*
2002.....	21.5	*	*	*	*	*	*	*	*	*
2001.....	22.0	*	*	*	*	*	*	*	*	*
2000.....	23.1	*	*	*	*	*	*	*	*	*
1999.....	23.2	*	*	*	*	*	*	*	*	*

All races 3/, female

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	10.3	*	*	*	*	*	*	*	*
2002.....	10.6	*	*	*	*	*	*	*	*
2001.....	10.8	*	*	*	*	*	*	*	*
2000.....	11.2	*	*	*	*	*	*	*	*
1999.....	11.4	*	*	*	*	*	*	*	*

White, male

2003.....	20.7	*	*	*	*	*	*	*	*
2002.....	21.5	*	*	*	*	*	*	*	*
2001.....	21.9	*	*	*	*	*	*	*	*
2000.....	22.8	*	*	*	*	*	*	*	*
1999.....	23.1	*	*	*	*	*	*	*	*

White, female

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of prostate (C61)

Black, both sexes

2003.....	12.8	*	*	*	*	*	*	*	*	*
2002.....	13.6	*	*	*	*	*	*	*	*	*
2001.....	14.2	*	*	*	*	*	*	*	*	*
2000.....	15.1	*	*	*	*	*	*	*	*	*
1999.....	14.8	*	*	*	*	*	*	*	*	*

Black, male

2003.....	26.9	*	*	*	*	*	*	*	*
2002.....	28.6	*	*	*	*	*	*	*	*
2001.....	29.7	*	*	*	*	*	*	*	*
2000.....	31.9	*	*	*	*	*	*	*	*
1999.....	31.2	*	*	*	*	*	*	*	*

Black, female

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	4.2	*	*	0.1	*	*	*	*	0.2	0.4
2002.....	4.2	*	*	0.1	*	*	*	*	0.2	0.4
2001.....	4.2	*	0.1	0.1	*	*	*	*	0.2	0.3
2000.....	4.3	*	*	0.1	*	*	*	*	0.1	0.4
1999.....	4.0	*	*	0.1	*	*	*	0.1	0.1	0.5

[illegible]

2003.....	3.1	*	*	*	*	*	*	*	0.2
2002.....	3.1	*	*	*	*	*	*	*	0.3
2001.....	3.1	*	*	0.2	*	*	*	*	*
2000.....	3.2	*	*	*	*	*	*	*	0.3
1999.....	3.0	*	*	*	*	*	*	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

[illegible]

2003.....	5.9	*	*	*	*	*	*	*	*
2002.....	5.9	*	*	*	*	*	*	*	*
2001.....	5.8	*	*	*	*	*	*	*	*
2000.....	5.8	*	*	*	*	*	*	*	*
1999.....	5.4	*	*	*	*	*	*	*	*

2003.....	3.4	*	*	*	*	*	*	*	*
2002.....	3.4	*	*	*	*	*	*	*	0.2
2001.....	3.4	*	*	*	*	*	*	*	*
2000.....	3.5	*	*	*	*	*	*	*	*
1999.....	3.2	*	*	*	*	*	*	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	2.4	*	*	*	*	*	*	*	*
2002.....	2.5	*	*	*	*	*	*	*	*
2001.....	2.4	*	*	*	*	*	*	*	*
2000.....	2.4	*	*	*	*	*	*	*	*
1999.....	2.5	*	*	*	*	*	*	*	*

Black, male

2003.....	2.8	*	*	*	*	*	*	*	*	*
2002.....	2.7	*	*	*	*	*	*	*	*	*
2001.....	2.5	*	*	*	*	*	*	*	*	*
2000.....	2.8	*	*	*	*	*	*	*	*	*
1999.....	2.9	*	*	*	*	*	*	*	*	*

Black, female

2003.....	2.1	*	*	*	*	*	*	*	*
2002.....	2.3	*	*	*	*	*	*	*	*
2001.....	2.2	*	*	*	*	*	*	*	*
2000.....	2.1	*	*	*	*	*	*	*	*
1999.....	2.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.6	0.5	0.7	0.8	0.8	0.5	0.6	0.9	1.3	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9

All races 3/, male

2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.2	*	0.7	0.9	0.9	0.6	0.7	1.1	1.8	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3

All races 3/, female

2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	4.0	*	0.6	0.8	0.6	0.4	0.6	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	1.0	1.3	2.1
2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.2	*	0.7	0.9	0.8	0.5	0.7	1.0	1.4	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2

White, male

2003.....	5.6	*	0.8	0.9	0.8	0.7	0.7	1.0	1.6	2.5
2002.....	5.7	*	0.7	1.1	0.7	0.7	0.8	0.7	1.6	2.3
2001.....	5.7	*	0.8	0.8	0.6	0.7	0.7	1.1	1.9	2.6
2000.....	5.8	*	0.7	0.9	0.9	0.6	0.8	1.2	1.8	2.9
1999.....	5.8	*	0.8	0.9	0.6	0.6	0.7	1.2	1.8	2.6

White, female

2003.....	4.4	*	0.6	0.9	0.8	0.6	0.5	0.9	1.0	1.8
2002.....	4.4	*	0.6	1.0	0.6	0.4	0.3	0.7	1.2	1.6
2001.....	4.4	*	0.7	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000.....	4.5	*	0.6	0.8	0.7	0.4	0.6	0.8	1.0	1.6
1999.....	4.6	*	0.8	0.8	0.7	0.5	0.5	0.9	1.0	1.7

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	2.1	*	*	0.8	0.6	*	*	*	*	1.2
2002.....	2.1	*	1.1	1.0	0.6	*	*	*	0.8	1.0
2001.....	2.0	*	*	*	0.7	*	*	*	0.7	*
2000.....	2.2	*	*	0.8	0.8	*	*	*	0.8	1.2
1999.....	2.1	*	*	1.1	*	0.7	*	*	0.9	0.9

Black, male

2003.....	2.5	*	*	*	*	*	*	*	*
2002.....	2.3	*	*	1.4	*	*	*	*	*
2001.....	2.2	*	*	*	*	*	*	*	*
2000.....	2.5	*	*	*	*	*	*	*	*
1999.....	2.3	*	*	*	*	*	*	*	*

Black, female

2003.....	1.8	*	*	*	*	*	*	*	*
2002.....	1.9	*	*	*	*	*	*	*	*
2001.....	1.8	*	*	*	*	*	*	*	*
2000.....	1.9	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.5	1.0	1.0	0.8	1.0	1.6	2.1	2.3	2.6	3.5
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7

All races 3/, male

2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	22.2	1.1	1.0	1.0	1.2	2.0	2.7	2.5	3.2	4.0
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4

All races 3/, female

2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.9	*	0.9	0.7	0.8	1.3	1.6	2.1	2.0	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, both sexes

2003.....	20.9	0.9	0.9	0.9	1.0	1.3	1.7	1.9	2.3	3.0
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	22.3	1.2	1.0	0.9	1.1	1.6	2.0	2.2	2.5	3.4
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6

White, male

2003.....	23.1	*	1.1	1.2	1.1	1.6	2.0	2.2	2.9	3.4
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	24.1	*	1.0	1.0	1.3	1.9	2.6	2.5	3.0	3.9
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3

White, female

2003.....	18.8	1.4	0.8	0.7	0.9	0.8	1.5	1.6	1.7	2.5
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.5	*	1.0	0.8	0.9	1.2	1.5	1.9	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2003.....	13.4	*	*	0.6	1.0	1.1	1.7	2.4	3.3	4.1
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	14.4	*	0.9	0.7	0.7	1.9	2.7	2.9	3.2	4.5
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6

Black, male

2003.....	13.9	*	*	*	1.2	*	2.1	2.5	4.2	4.7
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	15.2	*	*	*	*	2.3	3.2	2.7	4.7	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3

Black, female

2003.....	12.9	*	*	*	*	*	1.3	2.4	2.4	3.6
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.6	*	*	*	*	1.5	2.1	3.1	1.9	4.2
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Hodgkin's disease (C81)

All races 3/, both sexes

2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4

All races 3/, male

2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5

All races 3/, female

2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Hodgkin's disease (C81)

White, both sexes

2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4

White, male

2003.....	0.5	*	*	*	*	*	*	0.4	0.5	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5

White, female

2003.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3
2002.....	0.5	*	*	*	*	*	0.4	0.3	0.5	0.3
2001.....	0.4	*	*	*	*	*	*	0.4	0.3	0.3
2000.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	0.8	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.4	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.3	*	*	0.1	0.1	0.3	0.6	0.7	0.9	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4

All races 3/, male

2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.8	*	*	*	*	0.5	0.8	0.7	1.1	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8

All races 3/, female

2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.8	*	*	*	*	*	0.4	0.7	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2003.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.2	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

White, male

2003.....	8.9	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.7	*	*	*	*	0.5	0.8	0.7	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

White, female

2003.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.7	*	*	*	*	*	0.4	0.6	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Leukemia (C91-C95)

All races 3/, both sexes

2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.8	1.0	0.9	0.7	0.9	1.2	1.2	1.3	1.3	1.7
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6

All races 3/, male

2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.7	1.1	0.9	0.8	1.1	1.5	1.5	1.5	1.6	1.7
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8

All races 3/, female

2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.8	*	0.9	0.6	0.7	1.0	0.9	1.1	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Leukemia (C91-C95)

White, both sexes

2003.....	8.2	0.9	0.9	0.9	0.9	0.9	1.1	1.1	1.1	1.4
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.5	1.2	1.0	0.7	0.9	1.2	1.2	1.2	1.2	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6

White, male

2003.....	9.4	*	1.0	1.0	1.0	1.2	1.2	1.3	1.4	1.4
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.5	*	0.9	0.8	1.1	1.4	1.5	1.4	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8

White, female

2003.....	7.0	1.4	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.5	*	1.0	0.7	0.7	1.0	0.9	1.0	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Leukemia (C91-C95)

Black, both sexes

2003.....	4.6	*	*	*	0.8	0.7	1.0	1.0	1.4	2.0
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.9	*	*	*	0.6	1.5	1.3	1.5	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8

Black, male

2003.....	5.0	*	*	*	*	*	1.3	*	1.8	2.5
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.5	*	*	*	*	2.0	1.7	*	2.6	1.5
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8

Black, female

2003.....	4.2	*	*	*	*	*	*	*	*	1.7
2002.....	4.3	*	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	*	1.4
2000.....	4.4	*	*	*	*	*	*	1.6	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	*	1.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2003.....	21.5	*	0.7	0.6	0.7	1.2	1.3	1.6	1.9	3.1
2002.....	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3
2001.....	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2
2000.....	22.9	0.6	0.9	0.6	0.7	1.2	1.4	1.6	2.1	3.5
1999.....	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6

All races 3/, male

2003.....	22.3	*	0.8	0.8	0.7	1.5	1.6	1.9	2.4	3.5
2002.....	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7
2001.....	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5
2000.....	23.6	*	1.0	0.6	0.7	1.4	1.9	2.1	2.5	3.8
1999.....	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1

All races 3/, female

2003.....	20.8	*	0.5	0.5	0.7	0.8	1.1	1.2	1.4	2.7
2002.....	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8
2001.....	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0
2000.....	22.3	*	0.7	0.6	0.6	1.0	1.0	1.1	1.7	3.1
1999.....	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2003.....	23.1	*	0.6	0.7	0.7	1.3	1.4	1.6	1.8	3.1
2002.....	23.3	0.7	0.7	0.7	0.8	1.2	1.5	1.5	2.1	3.2
2001.....	23.5	*	0.8	0.6	0.8	1.2	1.5	1.7	1.9	3.1
2000.....	24.3	*	0.8	0.7	0.7	1.2	1.5	1.8	2.1	3.3
1999.....	25.3	*	0.8	0.7	0.7	1.2	1.4	1.6	2.2	3.5

White, male

2003.....	23.8	*	0.7	0.9	0.6	1.6	1.6	2.0	2.3	3.6
2002.....	24.2	*	0.8	0.7	0.7	1.4	1.9	2.0	2.5	3.6
2001.....	24.0	*	1.0	0.6	0.8	1.5	2.0	2.0	2.3	3.4
2000.....	24.9	*	1.0	0.7	0.7	1.4	1.9	2.3	2.6	3.7
1999.....	25.9	*	0.9	0.8	0.8	1.5	1.7	2.2	2.7	4.0

White, female

2003.....	22.5	*	0.5	0.5	0.7	0.9	1.1	1.1	1.3	2.6
2002.....	22.5	*	0.7	0.6	0.8	1.0	1.0	1.0	1.7	2.8
2001.....	22.9	*	0.7	0.5	0.7	1.0	0.8	1.4	1.5	2.8
2000.....	23.8	*	0.6	0.7	0.6	1.0	1.1	1.2	1.6	2.9
1999.....	24.6	*	0.7	0.7	0.6	1.0	1.1	1.0	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2003.....	17.3	*	0.9	0.6	0.9	0.9	1.3	1.9	2.7	3.7
2002.....	18.1	*	1.1	0.8	0.7	1.1	1.6	1.9	2.6	4.5
2001.....	18.3	*	*	*	0.8	1.2	1.3	2.0	2.6	4.3
2000.....	19.0	*	1.1	*	0.8	1.6	1.3	1.3	2.0	5.3
1999.....	19.5	*	*	0.6	0.9	1.3	1.6	1.5	2.4	4.9

Black, male

2003.....	18.3	*	*	*	*	1.2	1.7	1.7	3.0	3.6
2002.....	18.8	*	*	*	*	1.4	2.2	2.5	2.7	5.1
2001.....	19.4	*	*	*	*	1.5	1.9	2.5	3.0	4.4
2000.....	20.1	*	*	*	*	1.7	2.1	1.6	2.1	5.5
1999.....	20.6	*	*	*	*	1.9	2.0	1.5	3.1	5.1

Black, female

2003.....	16.5	*	*	*	*	*	*	2.0	2.3	3.9
2002.....	17.4	*	*	*	*	*	*	1.4	2.6	4.0
2001.....	17.3	*	*	*	*	*	*	1.5	2.3	4.2
2000.....	17.9	*	*	*	*	1.4	*	*	2.0	5.2
1999.....	18.5	*	*	*	*	*	*	1.5	1.8	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2003.....	4.7	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2002.....	4.6	1.7	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7
2000.....	4.9	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6

All races 3/, male

2003.....	4.7	1.6	0.4	0.2	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7
2000.....	4.9	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8
1999.....	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7

All races 3/, female

2003.....	4.7	1.5	0.3	*	*	*	0.2	0.3	0.5	0.6
2002.....	4.6	2.0	0.4	0.2	*	0.2	*	0.3	0.5	0.5
2001.....	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7
2000.....	5.0	1.5	0.3	0.3	0.2	*	0.2	0.4	0.5	0.6
1999.....	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2003.....	5.2	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.6
2002.....	5.1	1.9	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	5.3	1.2	0.4	0.2	0.2	0.2	0.3	0.4	0.4	0.7
2000.....	5.4	1.6	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7
1999.....	5.2	1.4	0.4	0.3	0.2	0.2	0.2	0.3	0.5	0.6

White, male

2003.....	5.2	1.6	0.3	*	*	*	0.3	0.4	0.5	0.7
2002.....	5.1	1.7	0.3	*	0.2	*	0.2	0.3	0.5	0.7
2001.....	5.3	1.2	0.4	*	*	*	0.3	0.4	0.4	0.7
2000.....	5.3	1.6	0.4	0.3	*	*	0.4	0.3	0.6	0.7
1999.....	5.2	1.4	0.4	0.3	*	*	0.3	0.3	0.5	0.7

White, female

2003.....	5.1	1.5	*	*	*	*	0.3	0.3	0.5	0.5
2002.....	5.1	2.1	0.4	0.3	*	*	*	*	0.4	0.5
2001.....	5.2	*	0.4	*	0.3	0.3	*	0.3	0.5	0.7
2000.....	5.5	1.6	*	0.3	*	*	*	0.3	0.5	0.6
1999.....	5.1	1.4	0.4	0.4	*	0.3	*	*	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Anemias (D50-D64)

All races 3/, both sexes

2003.....	1.6	0.5	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2002.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5
2001.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4
2000.....	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.4
1999.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4

All races 3/, male

2003.....	1.3	*	*	*	*	*	0.3	0.4	0.4	0.3
2002.....	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5
2001.....	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4
2000.....	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
1999.....	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4

All races 3/, female

2003.....	1.9	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4
2001.....	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2000.....	2.0	*	*	*	*	*	0.3	0.4	0.5	0.5
1999.....	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Anemias (D50-D64)

Black, both sexes

2003.....	2.7	*	*	*	*	*	1.9	1.9	2.2	2.2
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.7	*	*	*	*	*	1.7	1.8	2.9	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2

Black, male

2003.....	2.7	*	*	*	*	*	1.8	2.2	2.3	2.2
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.4	*	*	*	*	*	1.7	1.6	2.8	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5

Black, female

2003.....	2.8	*	*	*	*	*	2.1	1.7	2.2	2.2
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.1	*	*	*	*	*	1.6	2.0	3.0	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	25.2	*	*	*	*	0.2	0.6	1.2	2.1	3.1
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2

All races 3/, male

2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	23.5	*	*	*	*	0.3	0.7	1.3	2.7	3.6
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8

All races 3/, female

2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.8	*	*	*	*	*	0.6	1.0	1.6	2.6
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diabetes mellitus (E10-E14)

White, both sexes

2003.....	25.0	*	*	*	*	0.2	0.5	0.9	1.8	2.8
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.6	*	*	*	*	0.2	0.5	0.9	1.9	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8

White, male

2003.....	24.8	*	*	*	*	*	0.6	0.8	2.2	3.4
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	23.4	*	*	*	*	0.3	0.6	1.1	2.4	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4

White, female

2003.....	25.2	*	*	*	*	*	0.4	1.0	1.4	2.3
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.7	*	*	*	*	*	0.4	0.7	1.3	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diabetes mellitus (E10-E14)

Black, both sexes

2003.....	33.8	*	*	*	*	*	1.0	3.2	4.4	7.3
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	34.1	*	*	*	*	*	1.5	2.9	4.3	6.1
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6

Black, male

2003.....	29.8	*	*	*	*	*	*	4.1	5.2	8.4
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	28.4	*	*	*	*	*	1.5	3.2	4.7	7.3
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7

Black, female

2003.....	37.4	*	*	*	*	*	*	2.3	3.7	6.3
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	39.1	*	*	*	*	*	1.5	2.6	3.9	5.1
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	1.2	*	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*	*
2000.....	1.6	*	*	*	*	*	*	*	*	0.1
1999.....	1.6	*	*	*	*	*	*	*	*	*

2003.....	0.8	*	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*	*
2001.....	0.9	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*	*

2003.....	1.6	*	*	*	*	*	*	*	*
2002.....	1.8	*	*	*	*	*	*	*	*
2001.....	1.8	*	*	*	*	*	*	*	*
2000.....	2.1	*	*	*	*	*	*	*	*
1999.....	2.1	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2003.....	1.2	*	*	*	*	*	*	*	*
2002.....	1.3	*	*	*	*	*	*	*	*
2001.....	1.3	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*
1999.....	1.6	*	*	*	*	*	*	*	*

Black, male

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	1.2	*	*	*	*	*	*	*	*
2000.....	1.3	*	*	*	*	*	*	*	*
1999.....	1.5	*	*	*	*	*	*	*	*

Black, female

2003.....	1.4	*	*	*	*	*	*	*	*
2002.....	1.6	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.8	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malnutrition (E40-E46)

White, both sexes

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	1.3	*	*	*	*	*	*	*	*
2001.....	1.3	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	0.1
1999.....	1.5	*	*	*	*	*	*	*	*

White, male

2003.....	0.8	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*
1999.....	1.0	*	*	*	*	*	*	*	*

White, female

2003.....	1.5	*	*	*	*	*	*	*	*
2002.....	1.6	*	*	*	*	*	*	*	*
2001.....	1.7	*	*	*	*	*	*	*	*
2000.....	2.0	*	*	*	*	*	*	*	*
1999.....	2.0	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

All races 3/, male

All races 3/, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningitis (G00,G03)										
White, both sexes										
2003.....	0.2	1.7	*	*	*	*	*	*	*	*
2002.....	0.2	1.7	*	*	*	*	*	*	*	*
2001.....	0.2	1.5	*	*	*	*	*	*	0.1	*
2000.....	0.2	1.8	0.2	*	*	*	*	*	*	0.1
1999.....	0.3	2.3	0.3	*	*	*	*	*	*	0.1
White, male										
2003.....	0.3	2.0	*	*	*	*	*	*	*	*
2002.....	0.2	1.8	*	*	*	*	*	*	*	*
2001.....	0.2	1.4	*	*	*	*	*	*	*	*
2000.....	0.3	2.0	*	*	*	*	*	*	*	*
1999.....	0.3	3.0	0.3	*	*	*	*	*	*	*
White, female										
2003.....	0.2	1.3	*	*	*	*	*	*	*	*
2002.....	0.2	1.5	*	*	*	*	*	*	*	*
2001.....	0.2	1.6	*	*	*	*	*	*	*	*
2000.....	0.2	1.6	*	*	*	*	*	*	*	*
1999.....	0.3	1.5	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningitis (G00,G03)										
Black, both sexes										
2003.....	0.5	3.5	*	*	*	*	*	*	*	*
2002.....	0.5	3.0	*	*	*	*	*	*	*	*
2001.....	0.5	4.5	*	*	*	*	*	*	*	*
2000.....	0.6	6.0	*	*	*	*	*	*	*	*
1999.....	0.6	6.9	*	*	*	*	*	*	*	*
Black, male										
2003.....	0.5	*	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*	*
Black, female										
2003.....	0.5	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	*
1999.....	0.5	9.7	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	9.7	*	*	*	*	*	*	*	*
2002.....	8.9	*	*	*	*	*	*	*	*
2001.....	8.4	*	*	*	*	*	*	*	*
2000.....	7.7	*	*	*	*	*	*	*	*
1999.....	6.5	*	*	*	*	*	*	*	*

Black, male

2003.....	6.0	*	*	*	*	*	*	*	*
2002.....	5.3	*	*	*	*	*	*	*	*
2001.....	5.1	*	*	*	*	*	*	*	*
2000.....	4.6	*	*	*	*	*	*	*	*
1999.....	4.0	*	*	*	*	*	*	*	*

Black, female

2003.....	13.2	*	*	*	*	*	*	*	*
2002.....	12.0	*	*	*	*	*	*	*	*
2001.....	11.3	*	*	*	*	*	*	*	*
2000.....	10.5	*	*	*	*	*	*	*	*
1999.....	8.7	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

2003.....	310.3	13.9	1.5	0.7	1.0	2.5	4.5	7.1	13.2	25.1
2002.....	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3
2001.....	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0
2000.....	340.4	16.5	1.5	0.7	1.1	2.5	4.4	7.3	12.8	24.6
1999.....	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9

All races 3/, male

2003.....	296.3	15.2	1.4	0.7	1.2	3.1	5.3	8.5	17.0	33.4
2002.....	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9
2001.....	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1
2000.....	323.8	17.4	1.7	0.8	1.3	2.9	5.2	9.0	16.3	31.9
1999.....	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2

All races 3/, female

2003.....	323.9	12.5	1.5	0.6	0.8	2.0	3.6	5.6	9.3	16.8
2002.....	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7
2001.....	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0
2000.....	356.2	15.7	1.4	0.6	1.0	2.0	3.6	5.5	9.3	17.4
1999.....	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2003.....	330.0	11.0	1.2	0.6	0.9	2.2	3.7	5.7	11.0	21.4
2002.....	339.4	13.4	1.3	0.6	0.9	2.1	3.3	5.7	10.2	21.6
2001.....	344.8	12.2	1.5	0.6	1.0	1.9	3.4	5.9	10.9	21.4
2000.....	360.8	14.3	1.3	0.6	0.9	2.1	3.6	6.0	10.6	20.9
1999.....	362.7	14.6	1.3	0.7	0.9	2.4	3.4	5.4	10.6	22.0

White, male

2003.....	312.6	11.5	1.1	0.6	1.1	2.7	4.4	7.0	14.4	29.3
2002.....	320.9	14.3	1.5	0.5	1.1	2.7	4.3	7.5	13.3	28.5
2001.....	323.5	13.2	1.6	0.6	1.0	2.3	4.3	7.3	14.0	28.8
2000.....	340.9	15.0	1.6	0.7	1.1	2.5	4.3	7.6	14.0	28.1
1999.....	341.2	15.3	1.5	0.8	0.9	2.9	3.9	6.5	14.0	29.8

White, female

2003.....	347.0	10.4	1.4	0.5	0.7	1.7	2.8	4.3	7.4	13.2
2002.....	357.5	12.5	1.1	0.7	0.8	1.5	2.2	3.9	7.1	14.5
2001.....	365.5	11.1	1.5	0.7	0.9	1.5	2.5	4.3	7.6	13.9
2000.....	379.9	13.6	1.1	0.5	0.7	1.7	2.8	4.3	7.1	13.7
1999.....	383.6	13.8	1.2	0.6	0.9	1.9	2.9	4.2	7.2	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2003.....	273.1	26.2	2.8	1.2	1.6	4.4	9.6	16.1	29.3	52.7
2002.....	276.1	27.6	2.0	0.8	2.0	4.0	7.8	16.6	29.6	52.9
2001.....	280.4	28.6	3.5	1.0	1.6	4.0	9.4	16.4	28.5	51.5
2000.....	295.4	29.4	2.8	1.1	2.0	4.7	9.2	15.7	28.8	52.2
1999.....	289.4	29.2	2.4	1.0	1.8	4.8	9.0	14.4	26.8	55.2

Black, male

2003.....	268.6	31.2	3.3	1.4	1.7	5.6	11.6	19.5	36.3	65.7
2002.....	268.5	28.2	1.9	*	2.3	5.1	9.4	20.6	36.9	66.8
2001.....	272.4	28.1	3.6	*	1.5	5.1	11.3	20.3	35.2	65.6
2000.....	286.0	30.7	2.4	*	1.9	5.1	10.1	18.7	34.2	62.7
1999.....	282.5	30.6	2.5	*	2.1	6.3	10.9	17.7	33.0	69.6

Black, female

2003.....	277.2	21.0	2.4	*	1.5	3.1	7.6	13.0	23.0	41.0
2002.....	283.0	26.9	2.1	*	1.6	2.8	6.3	13.0	23.0	40.6
2001.....	287.8	29.0	3.4	1.3	1.6	2.8	7.5	12.9	22.4	38.9
2000.....	303.9	28.0	3.3	*	2.1	4.3	8.3	13.1	24.0	42.8
1999.....	295.7	27.9	2.4	*	1.5	3.3	7.1	11.5	21.3	42.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes

2003.....	235.6	11.0	1.2	0.5	0.8	1.9	3.6	5.6	10.5	20.1
2002.....	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3
2001.....	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9
2000.....	258.2	12.8	1.2	0.5	0.8	2.0	3.4	5.7	9.9	19.5
1999.....	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6

All races 3/, male

2003.....	235.0	12.1	1.1	0.6	0.9	2.4	4.4	6.9	13.9	27.8
2002.....	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8
2001.....	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4
2000.....	256.1	13.2	1.4	0.6	1.0	2.4	4.1	7.2	13.2	26.6
1999.....	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5

All races 3/, female

2003.....	236.2	9.8	1.3	0.5	0.6	1.4	2.7	4.2	7.1	12.5
2002.....	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8
2001.....	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4
2000.....	260.2	12.4	1.0	0.4	0.7	1.6	2.6	4.2	6.7	12.4
1999.....	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

2003.....	251.7	8.8	1.0	0.4	0.7	1.6	2.9	4.5	8.8	17.7
2002.....	258.5	10.6	0.9	0.4	0.7	1.7	2.5	4.6	8.1	17.7
2001.....	262.8	9.9	1.2	0.4	0.7	1.5	2.6	4.5	8.5	17.3
2000.....	274.8	11.3	1.0	0.5	0.7	1.7	2.7	4.5	8.3	16.9
1999.....	277.7	11.8	1.0	0.5	0.7	2.0	2.6	4.1	8.5	17.8

White, male

2003.....	249.5	9.4	0.8	0.4	0.8	2.0	3.5	5.7	12.0	25.1
2002.....	256.0	11.3	1.0	0.4	0.8	2.2	3.3	6.1	11.2	24.6
2001.....	257.8	10.6	1.2	0.4	0.8	1.8	3.4	5.9	11.3	24.4
2000.....	271.2	11.7	1.3	0.5	0.9	2.0	3.3	5.9	11.4	23.9
1999.....	272.9	12.0	1.2	0.6	0.7	2.4	3.0	5.2	11.4	25.2

White, female

2003.....	253.8	8.1	1.1	0.4	0.5	1.3	2.2	3.2	5.4	10.2
2002.....	261.0	9.9	0.8	0.5	0.5	1.1	1.6	3.0	5.0	10.7
2001.....	267.7	9.2	1.1	0.5	0.7	1.1	1.8	3.1	5.6	10.0
2000.....	278.3	10.9	0.8	0.4	0.5	1.3	2.0	3.1	5.1	9.9
1999.....	282.4	11.6	0.9	0.5	0.6	1.6	2.1	3.0	5.4	10.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2003.....	202.8	20.2	2.2	1.0	1.2	3.5	7.8	13.0	23.0	39.6
2002.....	205.6	21.3	1.6	*	1.3	3.4	6.5	13.4	23.2	40.8
2001.....	208.8	21.6	3.0	0.8	1.3	3.3	7.5	12.9	22.6	39.8
2000.....	219.6	20.9	2.2	1.0	1.5	4.1	7.5	13.4	22.3	38.9
1999.....	217.2	23.8	2.0	0.9	1.4	4.2	7.7	11.7	21.5	42.7

Black, male

2003.....	206.0	24.1	2.5	*	1.3	4.5	10.2	15.7	28.6	51.1
2002.....	206.3	22.1	*	*	1.5	4.5	8.0	17.5	30.0	54.4
2001.....	209.0	19.5	3.2	*	1.4	4.4	9.7	16.0	29.3	52.1
2000.....	219.0	20.9	2.1	*	1.5	4.8	8.6	16.2	27.8	49.5
1999.....	218.5	23.4	2.1	*	1.6	5.8	9.3	14.8	27.1	56.3

Black, female

2003.....	200.0	16.1	2.0	*	*	2.3	5.5	10.5	18.1	29.3
2002.....	205.0	20.6	1.8	*	*	2.4	5.0	9.7	17.0	28.6
2001.....	208.6	23.9	2.7	*	1.2	2.3	5.5	10.1	16.7	28.9
2000.....	220.1	21.0	2.3	*	1.4	3.3	6.4	10.8	17.4	29.6
1999.....	216.0	24.2	1.9	*	*	2.5	6.1	9.0	16.5	30.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	1.2	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	0.1
2001.....	1.2	*	*	*	*	*	*	0.1
2000.....	1.3	*	*	*	*	*	*	0.1
1999.....	1.3	*	*	*	*	*	*	0.1

2003.....	0.8	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	0.2
1999.....	0.8	*	*	*	*	*	*	*	*

2003.....	1.7	*	*	*	*	*	*	*	*
2002.....	1.7	*	*	*	*	*	*	*	*
2001.....	1.7	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.8	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*	*

2003.....	0.6	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*
1999.....	1.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Hypertensive heart disease (I11)

All races 3/, both sexes

2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.6	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5

All races 3/, male

2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.3	*	*	*	*	*	0.2	0.5	1.0	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0

All races 3/, female

2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.9	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Hypertensive heart disease (I11)

Black, both sexes

2003.....	17.9	*	*	*	*	*	0.6	1.5	3.5	6.0
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.7	*	*	*	*	*	*	1.4	2.9	5.0
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7

Black, male

2003.....	18.1	*	*	*	*	*	*	1.7	4.3	7.3
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	16.2	*	*	*	*	*	*	1.6	3.7	5.9
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3

Black, female

2003.....	17.8	*	*	*	*	*	*	*	2.7	4.8
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	17.1	*	*	*	*	*	*	*	2.2	4.2
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	187.2	0.8	*	*	*	0.2	0.5	1.3	3.6	10.0
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4

All races 3/, male

2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	193.6	*	*	*	*	0.2	0.6	2.0	5.4	15.0
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8

All races 3/, female

2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	181.1	*	*	*	*	*	0.3	0.6	1.8	5.1
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Ischemic heart diseases (I20-I25)

White, both sexes

2003.....	178.3	*	*	*	*	0.2	0.5	1.3	3.7	9.8
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	201.3	0.8	*	*	*	0.2	0.4	1.1	3.3	9.6
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9

White, male

2003.....	185.5	*	*	*	*	*	0.7	1.8	5.6	15.0
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	207.7	*	*	*	*	*	0.6	1.7	5.1	14.6
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1

White, female

2003.....	171.4	*	*	*	*	*	0.4	0.7	1.7	4.4
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	195.1	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Ischemic heart diseases (I20-I25)

Black, both sexes

2003.....	129.2	*	*	*	*	*	1.2	2.4	6.1	14.0
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	145.2	*	*	*	*	*	0.7	2.6	6.3	14.9
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1

Black, male

2003.....	133.4	*	*	*	*	*	1.6	3.1	9.2	19.6
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	146.8	*	*	*	*	*	*	3.9	8.3	20.6
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3

Black, female

2003.....	125.3	*	*	*	*	*	*	1.7	3.3	9.1
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	143.8	*	*	*	*	*	*	*	4.4	9.9
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	70.1	0.5	*	*	*	*	0.2	0.6	1.4	4.1
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1

All races 3/, male

2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	74.5	*	*	*	*	*	0.2	0.8	2.0	5.9
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2

All races 3/, female

2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	65.8	*	*	*	*	*	*	0.4	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Acute myocardial infarction (I21-I22)

White, both sexes

2003.....	63.4	*	*	*	*	*	0.2	0.5	1.4	4.1
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	75.4	*	*	*	*	*	0.2	0.5	1.3	4.1
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0

White, male

2003.....	67.7	*	*	*	*	*	0.3	0.7	2.0	6.3
2002.....	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	80.4	*	*	*	*	*	*	0.7	2.0	6.1
1999.....	82.7	*	*	*	*	*	*	0.6	2.1	6.2

White, female

2003.....	59.2	*	*	*	*	*	*	*	0.8	1.9
2002.....	62.8	*	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	70.6	*	*	*	*	*	*	0.3	0.6	2.1
1999.....	72.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2003.....	45.7	*	*	*	*	*	*	0.9	2.3	5.6
2002.....	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000.....	54.1	*	*	*	*	*	*	1.2	2.3	5.0
1999.....	53.3	*	*	*	*	*	*	0.8	2.4	5.5

Black, male

2003.....	46.6	*	*	*	*	*	*	*	3.5	7.5
2002.....	48.3	*	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	*	3.4	8.0
2000.....	53.9	*	*	*	*	*	*	1.7	2.8	6.6
1999.....	53.4	*	*	*	*	*	*	*	3.4	7.7

Black, female

2003.....	44.9	*	*	*	*	*	*	*	*	3.9
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	54.3	*	*	*	*	*	*	*	1.9	3.7
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*
2001.....	1.2	*	*	*	*	*	*	*	*
2000.....	1.2	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*

2003.....	1.1	*	*	*	*	*	*	*	*	0.2
2002.....	1.2	*	*	*	*	*	*	*	*	*
2001.....	1.2	*	*	*	*	*	*	*	*	*
2000.....	1.3	*	*	*	*	*	*	*	*	0.2
1999.....	1.2	*	*	*	*	*	*	*	*	*

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*
2001.....	1.2	*	*	*	*	*	*	*	*
2000.....	1.3	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*

2003.....	1.4	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*
1999.....	1.4	*	*	*	*	*	*	*	*

2003.....	0.9	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*
1999.....	1.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	115.9	*	*	*	*	*	0.3	0.7	2.1	5.8
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2

All races 3/, male

2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	117.7	*	*	*	*	*	0.4	1.2	3.3	8.9
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4

All races 3/, female

2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	114.1	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2003.....	113.8	*	*	*	*	*	0.3	0.8	2.2	5.6
2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	124.6	*	*	*	*	*	0.2	0.6	2.0	5.4
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8

White, male

2003.....	116.6	*	*	*	*	*	0.4	1.1	3.5	8.5
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	126.0	*	*	*	*	*	0.3	1.0	3.1	8.3
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8

White, female

2003.....	111.0	*	*	*	*	*	*	0.4	0.9	2.5
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	123.3	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, both sexes

2003.....	82.3	*	*	*	*	*	*	1.5	3.7	8.2
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	89.8	*	*	*	*	*	*	1.3	3.8	9.6
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1

Black, male

2003.....	85.5	*	*	*	*	*	*	2.1	5.7	11.7
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	91.3	*	*	*	*	*	*	2.2	5.2	13.7
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9

Black, female

2003.....	79.4	*	*	*	*	*	*	*	2.0	5.0
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	88.4	*	*	*	*	*	*	*	2.5	6.1
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	25.3	*	*	*	*	*	*	0.2	0.6	1.8
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0

All races 3/, male

2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.7	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1

All races 3/, female

2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.9	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

White, male

White, female

2003.....	21.9	*	*	*	*	*	*	*	*
2002.....	22.7	*	*	*	*	*	*	*	*
2001.....	23.3	*	*	*	*	*	*	0.3	0.7
2000.....	24.6	*	*	*	*	*	*	*	0.6
1999.....	25.6	*	*	*	*	*	*	*	0.7

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	90.6	*	*	*	*	*	0.2	0.5	1.6	4.1
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3

All races 3/, male

2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	91.1	*	*	*	*	*	0.2	0.9	2.4	6.1
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3

All races 3/, female

2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	90.1	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2003.....	90.2	*	*	*	*	*	0.2	0.5	1.5	3.7
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	99.1	*	*	*	*	*	0.1	0.5	1.4	3.8
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0

White, male

2003.....	91.3	*	*	*	*	*	0.3	0.7	2.2	5.7
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	99.5	*	*	*	*	*	*	0.8	2.2	5.7
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0

White, female

2003.....	89.1	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	98.7	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

Black, male

Black, female

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other heart diseases (I26-I51)

All races 3/, both sexes

2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	60.1	11.9	1.1	0.5	0.8	1.8	2.7	3.8	5.4	7.7
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2

All races 3/, male

2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	53.6	12.1	1.3	0.6	0.9	2.1	3.2	4.6	6.6	9.3
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3

All races 3/, female

2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	66.3	11.6	0.9	0.4	0.6	1.4	2.2	3.1	4.3	6.0
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other heart diseases (I26-I51)

White, both sexes

2003.....	62.2	8.2	0.9	0.4	0.6	1.5	2.2	2.8	4.4	6.4
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	63.5	10.4	0.9	0.4	0.6	1.5	2.1	3.1	4.5	6.2
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7

White, male

2003.....	54.7	8.8	0.8	0.4	0.8	1.8	2.5	3.2	5.5	7.9
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	55.7	10.6	1.2	0.5	0.8	1.7	2.6	3.8	5.5	7.6
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5

White, female

2003.....	69.6	7.6	1.0	0.3	0.4	1.1	1.8	2.4	3.2	5.0
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	71.1	10.2	0.7	0.3	0.4	1.2	1.6	2.4	3.4	4.7
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other heart diseases (I26-I51)

Black, both sexes

2003.....	52.5	19.6	2.0	0.9	1.0	3.0	5.9	8.9	12.8	18.1
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	54.7	20.1	2.2	0.9	1.4	3.7	6.0	9.1	12.4	18.2
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5

Black, male

2003.....	51.5	23.2	2.2	*	1.2	3.9	7.5	10.5	14.3	22.5
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	53.4	19.9	2.1	*	1.5	4.4	6.6	10.4	14.9	22.0
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3

Black, female

2003.....	53.4	15.8	1.8	*	*	2.1	4.2	7.3	11.4	14.3
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	55.8	20.3	2.2	*	1.3	3.0	5.4	7.9	10.1	14.7
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.6	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	0.7

2003.....	0.7	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*

2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.3	0.8	0.1	*	*	0.1	0.2	0.1	0.1	0.1
2002.....	0.3	0.6	0.1	*	*	*	*	*	0.1	0.2
2001.....	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2
2000.....	0.3	0.5	0.2	*	*	*	0.1	0.2	0.1	0.2
1999.....	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2

[illegible]

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	0.2

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

[illegible]

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	0.3

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes

2003.....	38.0	8.6	0.9	0.4	0.6	1.5	2.4	3.3	4.9	7.0
2002.....	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3
2001.....	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0
2000.....	39.2	10.5	0.9	0.4	0.6	1.6	2.5	3.5	5.0	6.9
1999.....	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4

All races 3/, male

2003.....	35.8	9.2	0.9	0.4	0.7	1.8	2.9	3.7	5.8	8.6
2002.....	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8
2001.....	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8
2000.....	37.0	10.4	1.0	0.5	0.8	1.9	2.9	4.1	6.1	8.5
1999.....	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3

All races 3/, female

2003.....	40.1	7.9	1.0	0.3	0.4	1.1	1.8	2.8	3.9	5.5
2002.....	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9
2001.....	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3
2000.....	41.2	10.6	0.8	0.3	0.5	1.2	2.0	2.8	3.9	5.4
1999.....	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2003.....	39.8	7.1	0.7	0.3	0.5	1.3	1.9	2.5	4.0	5.9
2002.....	40.4	9.1	0.7	0.3	0.5	1.4	1.7	2.7	3.6	6.1
2001.....	39.6	8.2	0.9	0.3	0.6	1.1	1.8	2.6	3.7	5.7
2000.....	40.7	9.4	0.8	0.3	0.5	1.3	2.0	2.7	4.1	5.6
1999.....	40.6	10.3	0.8	0.4	0.6	1.6	1.9	2.5	4.0	6.0

White, male

2003.....	36.9	7.4	0.7	0.3	0.7	1.6	2.2	2.9	5.0	7.2
2002.....	37.5	9.5	0.8	0.3	0.6	1.9	2.3	3.3	4.6	7.3
2001.....	36.5	8.2	0.9	0.3	0.6	1.3	2.3	3.1	4.4	7.2
2000.....	37.9	9.2	0.9	0.4	0.7	1.6	2.4	3.4	5.1	7.0
1999.....	37.9	10.7	1.0	0.4	0.6	1.9	2.2	3.0	4.9	7.7

White, female

2003.....	42.6	6.8	0.8	*	0.4	1.0	1.5	2.2	2.9	4.4
2002.....	43.2	8.7	0.6	0.3	0.4	0.9	1.1	2.1	2.6	4.8
2001.....	42.6	8.2	0.8	0.3	0.5	0.9	1.3	2.1	3.0	4.2
2000.....	43.4	9.5	0.6	*	0.3	1.0	1.5	2.1	3.1	4.2
1999.....	43.2	10.0	0.7	0.4	0.5	1.4	1.7	2.0	3.1	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2003.....	37.5	15.0	1.6	0.7	0.8	2.6	5.2	8.0	11.3	15.9
2002.....	37.8	17.2	1.1	*	0.9	2.9	4.8	8.7	11.4	17.0
2001.....	38.0	17.2	2.3	0.6	1.1	2.6	4.9	7.9	11.2	16.3
2000.....	39.6	16.5	1.6	0.8	1.1	3.3	5.5	8.3	11.2	16.2
1999.....	39.6	20.2	1.4	0.6	1.1	3.3	5.2	7.8	11.4	17.5

Black, male

2003.....	38.6	17.8	1.8	*	*	3.4	6.7	9.6	12.2	19.9
2002.....	38.3	18.9	*	*	1.2	3.7	5.8	10.9	14.5	20.5
2001.....	38.9	15.0	2.6	*	1.1	3.4	6.3	9.3	13.5	20.3
2000.....	40.7	15.5	1.8	*	1.2	3.9	6.0	9.5	13.8	19.7
1999.....	41.5	19.8	*	*	1.5	4.6	6.2	9.2	13.8	21.8

Black, female

2003.....	36.5	12.1	*	*	*	1.8	3.6	6.6	10.4	12.4
2002.....	37.4	15.4	*	*	*	2.0	3.8	6.7	8.7	13.9
2001.....	37.2	19.5	2.0	*	*	1.7	3.5	6.6	9.2	12.8
2000.....	38.6	17.5	*	*	*	2.7	4.9	7.3	9.0	13.1
1999.....	37.9	20.5	*	*	*	2.0	4.2	6.5	9.2	13.7

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	7.5	*	*	*	*	*	*	0.1	0.3	0.6
2002.....	7.0	*	*	*	*	*	*	0.2	0.2	0.6
2001.....	6.8	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	6.6	*	*	*	*	*	*	0.1	0.3	0.5
1999.....	6.1	*	*	*	*	*	*	0.1	0.3	0.5

All races 3/, male

[illegible]

All races 3/, female

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

[illegible]

Black, male

[illegible]

Black, female

2003.....	13.0	*	*	*	*	*	*	*	*
2002.....	12.5	*	*	*	*	*	*	*	*
2001.....	11.9	*	*	*	*	*	*	*	1.4
2000.....	12.4	*	*	*	*	*	*	*	*
1999.....	11.4	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	60.9	3.2	0.3	0.1	0.3	0.3	0.7	1.2	2.0	4.0
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0

All races 3/, male

2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	48.1	3.8	*	*	0.2	0.4	0.7	1.2	2.0	3.9
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9

All races 3/, female

2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	73.2	2.7	0.4	*	0.3	0.3	0.7	1.1	2.0	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Cerebrovascular diseases (I60-I69)

White, both sexes

2003.....	57.0	2.0	0.2	0.1	0.2	0.4	0.5	0.8	1.6	2.7
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	63.9	2.5	0.2	0.1	0.2	0.3	0.7	1.1	1.7	3.1
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2

White, male

2003.....	44.2	2.0	*	*	*	0.4	0.6	0.8	1.7	2.8
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	49.4	3.0	*	*	*	0.4	0.7	1.3	1.7	3.1
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2

White, female

2003.....	69.5	2.0	*	*	*	0.3	0.5	0.8	1.5	2.5
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	77.9	2.0	*	*	0.3	*	0.6	1.0	1.6	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Cerebrovascular diseases (I60-I69)

Black, both sexes

2003.....	49.3	4.9	*	*	*	0.7	1.2	2.0	4.1	8.8
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	54.4	7.9	*	*	*	*	1.1	1.6	4.5	9.8
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1

Black, male

2003.....	43.2	5.9	*	*	*	*	*	2.6	4.1	9.7
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	47.8	9.1	*	*	*	*	*	*	4.3	9.3
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9

Black, female

2003.....	54.8	*	*	*	*	*	1.6	1.5	4.1	8.0
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	60.4	*	*	*	*	*	*	1.7	4.7	10.2
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2003.....	8.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2002.....	8.6	*	*	*	*	0.2	0.2	0.3	0.5	0.7
2001.....	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	9.5	*	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8

All races 3/, male

2003.....	9.2	*	*	*	*	*	0.3	0.4	0.6	1.1
2002.....	9.4	*	*	*	*	*	0.2	0.3	0.7	0.7
2001.....	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2
2000.....	10.5	*	*	*	*	*	0.3	0.4	0.7	0.8
1999.....	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0

All races 3/, female

2003.....	7.8	*	*	*	*	*	*	*	0.3	0.5
2002.....	7.9	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	8.2	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	8.5	*	*	*	*	*	*	*	0.4	0.6
1999.....	8.4	*	*	*	*	*	*	*	0.2	0.5

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	0.1	*	0.2	0.4	0.5
2001.....	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6
2000.....	5.7	*	*	*	*	*	0.1	0.2	0.4	0.5
1999.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6

All races 3/, male

2003.....	6.3	*	*	*	*	*	0.2	0.3	0.5	0.9
2002.....	6.4	*	*	*	*	*	*	0.3	0.5	0.6
2001.....	6.5	*	*	*	*	*	*	0.4	0.5	0.9
2000.....	7.2	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	7.0	*	*	*	*	*	*	0.3	0.5	0.9

All races 3/, female

2003.....	3.9	*	*	*	*	*	*	*
2002.....	4.0	*	*	*	*	*	*	* 0.2
2001.....	4.2	*	*	*	*	*	*	* 0.3
2000.....	4.4	*	*	*	*	*	*	* 0.2
1999.....	4.4	*	*	*	*	*	*	* 0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

2003.....	3.8	*	*	*	*	*	*	*	*
2002.....	4.0	*	*	*	*	*	*	*	*
2001.....	4.0	*	*	*	*	*	*	*	*
2000.....	4.1	*	*	*	*	*	*	*	*
1999.....	4.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	3.4	*	*	*	*	*	*	*	*	*
2002.....	3.5	*	*	*	*	*	*	*	*	*
2001.....	3.7	*	*	*	*	*	*	*	*	*
2000.....	3.7	*	*	*	*	*	*	*	*	*
1999.....	3.5	*	*	*	*	*	*	*	*	*

2003.....	2.9	*	*	*	*	*	*	*	*
2002.....	3.0	*	*	*	*	*	*	*	*
2001.....	3.2	*	*	*	*	*	*	*	*
2000.....	3.2	*	*	*	*	*	*	*	*
1999.....	2.9	*	*	*	*	*	*	*	*

2003.....	3.8	*	*	*	*	*	*	*	*
2002.....	3.9	*	*	*	*	*	*	*	*
2001.....	4.2	*	*	*	*	*	*	*	*
2000.....	4.2	*	*	*	*	*	*	*	*
1999.....	4.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.7	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4

All races 3/, male

2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.5	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5

All races 3/, female

2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.9	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2003.....	1.6	0.9	*	*	*	*	0.1	0.2	0.3	0.5
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.7	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

White, male

2003.....	1.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	1.3	*	*	*	*	*	*	*	0.4	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.5	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.4

White, female

2003.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	1.9	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.3
2000.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

2003.....	22.4	8.0	1.0	0.4	0.3	0.4	0.7	0.8	1.1	1.8
2002.....	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7
2001.....	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8
2000.....	23.7	7.5	0.7	0.2	0.2	0.3	0.7	0.8	1.1	1.8
1999.....	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0

All races 3/, male

2003.....	20.1	9.4	1.1	0.4	0.3	0.5	0.7	0.9	1.3	1.9
2002.....	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0
2000.....	21.3	8.3	0.7	0.3	0.2	0.3	0.7	0.9	1.3	1.9
1999.....	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2

All races 3/, female

2003.....	24.6	6.6	0.9	0.4	0.4	0.3	0.7	0.6	0.8	1.6
2002.....	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6
2001.....	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6
2000.....	26.1	6.7	0.7	0.2	*	0.3	0.6	0.7	1.0	1.7
1999.....	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Influenza and pneumonia (J10-J18)

White, both sexes

2003.....	24.4	6.7	0.9	0.3	0.3	0.4	0.7	0.7	1.0	1.6
2002.....	24.9	5.2	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.5
2001.....	23.6	5.3	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2000.....	25.6	5.7	0.6	0.2	0.2	0.3	0.6	0.7	1.0	1.7
1999.....	24.8	6.2	0.6	0.2	0.2	0.3	0.5	0.5	0.8	1.7

White, male

2003.....	21.4	8.0	1.1	0.3	0.3	0.4	0.7	0.9	1.2	1.8
2002.....	21.9	6.6	0.6	*	*	0.4	0.5	0.7	1.0	1.5
2001.....	20.7	6.2	0.7	*	*	0.3	0.7	0.5	1.0	1.7
2000.....	22.5	6.3	0.5	0.3	*	0.3	0.7	0.7	1.1	1.8
1999.....	21.5	6.7	0.5	0.3	0.3	0.3	0.5	0.6	1.0	1.9

White, female

2003.....	27.3	5.4	0.8	0.4	0.4	0.3	0.7	0.5	0.7	1.4
2002.....	27.8	3.7	0.5	*	0.3	0.3	0.4	0.7	0.7	1.4
2001.....	26.4	4.4	0.6	*	*	0.3	0.3	0.6	0.8	1.4
2000.....	28.6	5.0	0.7	*	*	0.3	0.5	0.7	1.0	1.6
1999.....	27.9	5.6	0.7	*	*	0.4	0.4	0.5	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Influenza and pneumonia (J10-J18)

Black, both sexes

2003.....	15.4	14.1	1.1	0.6	*	*	0.8	1.6	2.1	3.3
2002.....	15.6	12.9	1.6	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.1	3.7
2000.....	17.0	17.7	1.3	*	*	*	1.1	1.5	2.1	2.9
1999.....	16.2	19.0	1.7	*	*	*	1.3	1.7	2.2	4.1

Black, male

2003.....	16.0	16.1	*	*	*	*	*	1.7	2.8	3.5
2002.....	15.4	13.9	1.7	*	*	*	*	*	2.4	4.0
2001.....	15.9	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	17.4	18.9	1.9	*	*	*	*	2.2	2.6	3.1
1999.....	16.4	18.9	2.0	*	*	*	*	1.9	2.5	4.5

Black, female

2003.....	14.9	12.1	*	*	*	*	*	1.6	1.4	3.1
2002.....	15.7	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.2	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	16.6	16.4	*	*	*	*	*	*	1.6	2.8
1999.....	16.1	19.1	*	*	*	*	*	1.5	1.9	3.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.6	0.8	0.4	0.1	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*

2003.....	0.5	*	0.4	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*

2003.....	0.7	*	0.3	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonia (J12-J18)

All races 3/, both sexes

2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	23.1	7.3	0.6	0.2	0.2	0.3	0.6	0.8	1.1	1.8
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0

All races 3/, male

2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.8	8.0	0.6	0.3	*	0.3	0.7	0.9	1.2	1.9
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2

All races 3/, female

2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	25.3	6.5	0.6	*	*	0.3	0.6	0.6	1.0	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonia (J12-J18)

White, both sexes

2003.....	23.7	6.0	0.6	0.2	0.3	0.3	0.7	0.6	0.9	1.5
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.9	5.4	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7

White, male

2003.....	20.9	7.1	0.7	*	0.3	0.4	0.7	0.8	1.1	1.7
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.9	6.0	0.4	0.3	*	0.3	0.7	0.7	1.1	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9

White, female

2003.....	26.5	4.8	0.5	*	0.3	*	0.6	0.5	0.7	1.3
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.7	4.9	0.6	*	*	*	0.5	0.6	0.9	1.6
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonia (J12-J18)

Black, both sexes

2003.....	15.1	13.0	0.9	*	*	*	0.7	1.6	2.0	3.3
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.7	17.3	1.3	*	*	*	1.0	1.4	2.1	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0

Black, male

2003.....	15.8	15.2	*	*	*	*	*	1.6	2.8	3.5
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	17.2	18.9	1.9	*	*	*	*	2.1	2.6	3.1
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4

Black, female

2003.....	14.6	10.8	*	*	*	*	*	1.5	1.4	3.1
2002.....	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	16.3	15.7	*	*	*	*	*	*	1.6	2.8
1999.....	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.2	4.0	*	*	*	*	*	*	*	*

Black, male

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Black, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Black, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.2	4.0	*	*	*	*	*	*	*	*

Black, male

2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

Black, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	44.3	0.9	0.3	0.2	0.5	0.4	0.6	0.7	0.9	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3

All races 3/, male

2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	44.6	1.2	0.4	0.3	0.5	0.6	0.7	0.8	1.0	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2

All races 3/, female

2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	44.1	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.5
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2003.....	49.5	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.9	0.8	0.2	0.1	0.3	0.3	0.4	0.5	0.6	1.1
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1

White, male

2003.....	47.4	1.3	*	*	0.3	0.2	0.5	0.4	0.5	1.1
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	49.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0

White, female

2003.....	51.5	*	*	*	*	*	0.3	0.5	0.7	1.3
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	50.4	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2003.....	20.2	*	1.0	0.7	1.1	1.3	1.3	1.7	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	21.5	*	0.9	0.8	1.5	1.4	1.3	2.0	2.5	3.6
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2

Black, male

2003.....	23.3	*	1.6	*	1.5	1.6	1.5	2.4	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	25.3	*	*	*	1.8	2.0	1.5	2.5	3.3	3.8
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Black, female

2003.....	17.3	*	*	*	*	*	*	*	*	2.9
2002.....	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000.....	18.2	*	*	*	*	*	*	1.5	1.7	3.4
1999.....	18.0	*	*	*	*	*	*	2.4	2.8	3.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.3	0.5	*	*	*	*	*	*	*	*
2002.....	0.3	0.6	*	*	*	*	*	*	*	*
2001.....	0.3	0.5	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.4	0.5	*	*	*	*	*	*	*	*

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Emphysema (J43)

White, both sexes

2003.....	5.9	*	*	*	*	*	*	*	*
2002.....	6.2	*	*	*	*	*	*	*	*
2001.....	6.6	*	*	*	*	*	*	*	0.1
2000.....	6.9	*	*	*	*	*	*	*	*
1999.....	7.3	*	*	*	*	*	*	*	*

White, male

2003.....	5.9	*	*	*	*	*	*	*	*	*
2002.....	6.4	*	*	*	*	*	*	*	*	*
2001.....	6.8	*	*	*	*	*	*	*	*	*
2000.....	7.2	*	*	*	*	*	*	*	*	*
1999.....	7.8	*	*	*	*	*	*	*	*	*

White, female

2003.....	5.8	*	*	*	*	*	*	*	*
2002.....	5.9	*	*	*	*	*	*	*	*
2001.....	6.3	*	*	*	*	*	*	*	*
2000.....	6.6	*	*	*	*	*	*	*	*
1999.....	6.8	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	2.1	*	*	*	*	*	*	*	*
2002.....	2.3	*	*	*	*	*	*	*	*
2001.....	2.2	*	*	*	*	*	*	*	*
2000.....	2.6	*	*	*	*	*	*	*	*
1999.....	2.6	*	*	*	*	*	*	*	*

2003.....	2.7	*	*	*	*	*	*	*	*
2002.....	3.0	*	*	*	*	*	*	*	*
2001.....	2.7	*	*	*	*	*	*	*	*
2000.....	3.3	*	*	*	*	*	*	*	*
1999.....	3.5	*	*	*	*	*	*	*	*

2003.....	1.6	*	*	*	*	*	*	*	*
2002.....	1.6	*	*	*	*	*	*	*	*
2001.....	1.7	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.7	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Asthma (J45-J46)

All races 3/, both sexes

2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.6	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9

All races 3/, male

2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8

All races 3/, female

2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.4	0.3	0.4	0.5	0.7	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Asthma (J45-J46)

White, both sexes

2003.....	1.2	*	*	*	0.2	0.2	0.3	0.4	0.4	0.5
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7

White, male

2003.....	0.8	*	*	*	0.3	*	0.3	0.4	0.4	0.4
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	1.0	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6

White, female

2003.....	1.6	*	*	*	*	*	*	0.4	0.5	0.7
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Asthma (J45-J46)

Black, both sexes

2003.....	2.7	*	0.9	0.7	1.0	1.1	1.3	1.4	1.9	2.0
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.3	*	*	0.8	1.4	1.3	1.2	1.9	2.1	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6

Black, male

2003.....	2.4	*	1.5	*	1.4	1.3	1.4	2.1	2.7	1.9
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.9	*	*	*	1.7	1.7	*	2.4	2.7	2.8
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6

Black, female

2003.....	2.9	*	*	*	*	*	*	*	*	2.1
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.7	*	*	*	*	*	*	*	1.5	2.7
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	15.2	*	*	*	*	*	*	*	*
2002.....	15.4	*	*	*	*	*	*	*	0.8
2001.....	15.1	*	*	*	*	*	*	*	1.1
2000.....	15.5	*	*	*	*	*	*	*	0.7
1999.....	15.9	*	*	*	*	*	*	*	*

Black, male

2003.....	18.0	*	*	*	*	*	*	*	*
2002.....	18.2	*	*	*	*	*	*	*	*
2001.....	18.1	*	*	*	*	*	*	*	*
2000.....	18.9	*	*	*	*	*	*	*	*
1999.....	19.6	*	*	*	*	*	*	*	*

Black, female

2003.....	12.6	*	*	*	*	*	*	*	*
2002.....	12.9	*	*	*	*	*	*	*	*
2001.....	12.3	*	*	*	*	*	*	*	*
2000.....	12.4	*	*	*	*	*	*	*	*
1999.....	12.5	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.7	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	6.0	*	*	*	*	*	0.2	0.1	0.2	0.3
2002.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	6.0	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3

All races 3/, male

2003.....	6.1	*	*	*	*	*	0.2	*	0.2	0.4
2002.....	6.3	*	*	*	*	*	*	*	0.2	0.4
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	6.2	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.7	*	*	*	*	*	*	0.2	0.3	0.3

All races 3/, female

2003.....	5.8	*	*	*	*	*	*	*	*
2002.....	5.9	*	*	*	*	*	*	*	*
2001.....	6.0	*	*	*	*	*	*	*	0.2
2000.....	5.9	*	*	*	*	*	*	*	*
1999.....	5.3	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	6.5	*	*	*	*	*	0.1	*	0.2	0.3
2002.....	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	6.6	*	*	*	*	*	*	*	0.2	0.2
1999.....	5.9	*	*	*	*	*	*	0.1	0.2	0.2

White, male

[illegible]

White, female

2003.....	6.4	*	*	*	*	*	*	*	*
2002.....	6.6	*	*	*	*	*	*	*	*
2001.....	6.6	*	*	*	*	*	*	*	*
2000.....	6.4	*	*	*	*	*	*	*	*
1999.....	5.8	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2003.....	8.8	7.0	0.5	0.1	0.2	0.3	0.4	0.4	0.6	0.9
2002.....	8.7	6.9	0.5	0.2	0.2	0.3	0.4	0.4	0.5	0.9
2001.....	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0
2000.....	9.3	7.1	0.6	0.2	0.2	0.3	0.4	0.6	0.8	1.0
1999.....	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0

All races 3/, male

2003.....	8.7	8.0	0.6	*	0.2	0.3	0.4	0.5	0.6	0.9
2002.....	8.5	7.9	0.7	0.2	*	0.3	0.4	0.3	0.5	0.9
2001.....	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9
2000.....	9.2	7.8	0.6	0.2	*	0.3	0.5	0.5	0.8	1.2
1999.....	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1

All races 3/, female

2003.....	8.9	5.9	0.5	*	*	0.3	0.3	0.3	0.6	0.9
2002.....	8.8	5.8	0.4	0.2	0.2	0.2	0.4	0.4	0.6	0.8
2001.....	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0
2000.....	9.3	6.3	0.6	*	0.2	0.3	0.3	0.6	0.8	0.9
1999.....	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2003.....	9.6	5.4	0.4	0.1	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	9.5	4.8	0.5	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	10.0	5.0	0.5	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	10.0	5.5	0.5	0.2	0.2	0.2	0.3	0.5	0.7	0.8
1999.....	9.7	4.9	0.5	0.2	0.2	0.3	0.3	0.4	0.7	0.8

White, male

2003.....	9.4	5.8	0.4	*	*	*	0.3	0.4	0.6	0.8
2002.....	9.2	5.4	0.6	*	*	*	0.4	0.3	0.5	0.8
2001.....	9.7	5.9	0.5	*	0.2	0.3	0.4	0.3	0.5	0.8
2000.....	9.9	6.4	0.5	*	*	*	0.4	0.5	0.7	1.0
1999.....	9.6	5.6	0.6	*	*	0.4	0.4	0.5	0.7	0.9

White, female

2003.....	9.8	5.0	0.4	*	*	0.3	*	0.3	0.5	0.7
2002.....	9.7	4.2	*	*	*	*	*	0.4	0.5	0.7
2001.....	10.2	4.0	0.6	*	*	*	0.4	0.5	0.5	0.7
2000.....	10.2	4.5	0.6	*	*	0.3	*	0.5	0.7	0.7
1999.....	9.8	4.3	0.5	*	*	*	*	0.3	0.7	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2003.....	5.9	15.0	1.1	*	*	*	0.8	0.8	1.2	1.8
2002.....	6.0	16.6	1.1	*	*	*	0.9	*	1.0	1.8
2001.....	6.7	18.0	1.0	*	*	*	0.8	1.1	1.4	2.5
2000.....	6.7	17.7	0.9	*	*	*	0.8	0.9	1.4	2.4
1999.....	6.5	15.4	1.0	*	*	*	*	1.1	1.6	2.3

Black, male

2003.....	6.0	18.4	*	*	*	*	*	*	*	1.9
2002.....	6.0	19.5	*	*	*	*	*	*	*	1.9
2001.....	6.4	23.1	*	*	*	*	*	*	1.5	2.0
2000.....	6.8	17.2	*	*	*	*	*	*	*	3.0
1999.....	6.4	17.6	*	*	*	*	*	*	1.7	2.5

Black, female

2003.....	5.7	11.4	*	*	*	*	*	*	*	1.7
2002.....	6.0	13.6	*	*	*	*	*	*	*	1.6
2001.....	7.0	12.6	*	*	*	*	*	*	*	2.9
2000.....	6.6	18.2	*	*	*	*	*	*	*	1.9
1999.....	6.7	13.1	*	*	*	*	*	1.7	1.6	2.2

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	1.4	*	*	*	*	*	*	*	*
2002.....	1.5	*	*	*	*	*	*	*	*
2001.....	1.7	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.8	*	*	*	*	*	*	*	*

2003.....	1.4	*	*	*	*	*	*	*	*	*
2002.....	1.5	*	*	*	*	*	*	*	*	0.2
2001.....	1.5	*	*	*	*	*	*	*	*	*
2000.....	1.7	*	*	*	*	*	*	*	*	0.2
1999.....	1.7	*	*	*	*	*	*	*	*	*

2003.....	1.5	*	*	*	*	*	*	*	*
2002.....	1.6	*	*	*	*	*	*	*	*
2001.....	1.9	*	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*
2000.....	1.2	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*

2003.....	1.2	*	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*	*
2000.....	1.4	*	*	*	*	*	*	*	*	*
1999.....	1.4	*	*	*	*	*	*	*	*	*

2003.....	0.8	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*
2001.....	0.9	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

All races 3/, male

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

All races 3/, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

White, male

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

White, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

Black, male

Black, female

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.6	0.9	*	*	*	*	*	*	*	*
2002.....	0.6	0.7	*	*	*	*	*	*	*	*
2001.....	0.5	1.1	*	*	*	*	*	*	*	*
2000.....	0.6	1.2	*	*	*	*	*	*	*	*
1999.....	0.5	1.2	*	*	*	*	*	*	*	*

2003.....	0.5	1.0	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	1.4	*	*	*	*	*	*	*	*
2000.....	0.4	1.4	*	*	*	*	*	*	*	*
1999.....	0.4	1.3	*	*	*	*	*	*	*	*

2003.....	0.6	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

2003.....	9.5	*	*	*	*	*	*	0.4	1.4	4.1
2002.....	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2
2001.....	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0
2000.....	9.6	*	*	*	*	*	0.2	0.5	1.7	4.9
1999.....	9.4	*	*	*	*	*	*	0.4	1.6	4.8

All races 3/, male

2003.....	12.5	*	*	*	*	*	*	0.6	1.9	5.4
2002.....	12.3	*	*	*	*	*	*	0.5	1.8	5.5
2001.....	12.4	*	*	*	*	*	*	0.6	2.0	6.4
2000.....	12.8	*	*	*	*	*	*	0.7	2.2	6.5
1999.....	12.5	*	*	*	*	*	*	0.4	2.1	6.3

All races 3/, female

2003.....	6.5	*	*	*	*	*	*	0.3	0.8	2.9
2002.....	6.7	*	*	*	*	*	*	0.3	1.0	3.0
2001.....	6.7	*	*	*	*	*	*	0.3	1.0	3.5
2000.....	6.6	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	6.4	*	*	*	*	*	*	0.3	1.2	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2003.....	10.2	*	*	*	*	*	*	0.4	1.4	4.2
2002.....	10.1	*	*	*	*	*	*	0.4	1.4	4.3
2001.....	10.1	*	*	*	*	*	*	0.3	1.5	5.1
2000.....	10.2	*	*	*	*	*	*	0.4	1.6	4.8
1999.....	9.9	*	*	*	*	*	*	0.3	1.6	4.8

White, male

2003.....	13.4	*	*	*	*	*	*	0.5	2.0	5.6
2002.....	13.2	*	*	*	*	*	*	0.4	1.9	5.6
2001.....	13.1	*	*	*	*	*	*	0.5	2.1	6.5
2000.....	13.5	*	*	*	*	*	*	0.6	2.3	6.4
1999.....	13.1	*	*	*	*	*	*	0.4	2.1	6.4

White, female

2003.....	7.0	*	*	*	*	*	*	*	0.8	2.9
2002.....	7.2	*	*	*	*	*	*	0.3	0.9	3.0
2001.....	7.1	*	*	*	*	*	*	*	1.0	3.7
2000.....	6.9	*	*	*	*	*	*	*	1.0	3.1
1999.....	6.8	*	*	*	*	*	*	*	1.1	3.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

[illegible]

2003.....	9.5	*	*	*	*	*	*	*		
2002.....	9.0	*	*	*	*	*	*	*	*	
2001.....	10.4	*	*	*	*	*	*	*	*	
2000.....	10.3	*	*	*	*	*	*	*	*	
1999.....	10.7	*	*	*	*	*	*	*	1.8	

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcoholic liver disease (K70)

All races 3/, both sexes

2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.4	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3

All races 3/, male

2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.7	*	*	*	*	*	*	0.5	1.6	4.4
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3

All races 3/, female

2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.7	2.3
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcoholic liver disease (K70)

White, both sexes

2003.....	4.5	*	*	*	*	*	*	0.2	1.0	2.9
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3

White, male

2003.....	6.8	*	*	*	*	*	*	0.4	1.4	3.9
2002.....	6.6	*	*	*	*	*	*	0.3	1.3	3.7
2001.....	6.7	*	*	*	*	*	*	0.3	1.5	4.5
2000.....	7.0	*	*	*	*	*	*	0.5	1.7	4.3
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	4.3

White, female

2003.....	2.3	*	*	*	*	*	*	*	0.5	1.9
2002.....	2.3	*	*	*	*	*	*	*	0.6	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.7	2.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

Black, male

Black, female

2003.....	1.8	*	*	*	*	*	*	*	*
2002.....	2.0	*	*	*	*	*	*	*	*
2001.....	2.1	*	*	*	*	*	*	*	*
2000.....	2.5	*	*	*	*	*	*	*	*
1999.....	2.4	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes

2003.....	5.2	*	*	*	*	*	*	0.2	0.4	1.3
2002.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5
2001.....	5.2	*	*	*	*	*	*	0.2	0.4	1.6
2000.....	5.2	*	*	*	*	*	0.1	0.2	0.5	1.5
1999.....	5.1	*	*	*	*	*	*	0.2	0.6	1.5

All races 3/, male

2003.....	6.2	*	*	*	*	*	*	*	0.5	1.6
2002.....	6.1	*	*	*	*	*	*	*	0.6	1.9
2001.....	6.1	*	*	*	*	*	*	0.2	0.6	2.0
2000.....	6.1	*	*	*	*	*	*	*	0.6	2.1
1999.....	6.0	*	*	*	*	*	*	*	0.7	2.0

All races 3/, female

2003.....	4.3	*	*	*	*	*	*	*	0.3	0.9
2002.....	4.5	*	*	*	*	*	*	0.2	0.4	1.0
2001.....	4.4	*	*	*	*	*	*	*	0.3	1.3
2000.....	4.4	*	*	*	*	*	*	*	0.4	1.0
1999.....	4.3	*	*	*	*	*	*	*	0.4	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2003.....	5.7	*	*	*	*	*	*	*	0.4	1.3
2002.....	5.7	*	*	*	*	*	*	0.2	0.4	1.5
2001.....	5.6	*	*	*	*	*	*	*	0.5	1.7
2000.....	5.6	*	*	*	*	*	*	*	0.5	1.5
1999.....	5.5	*	*	*	*	*	*	0.1	0.6	1.5

White, male

2003.....	6.6	*	*	*	*	*	*	*	0.6	1.7
2002.....	6.5	*	*	*	*	*	*	*	0.6	2.0
2001.....	6.4	*	*	*	*	*	*	*	0.6	2.0
2000.....	6.5	*	*	*	*	*	*	*	0.6	2.1
1999.....	6.4	*	*	*	*	*	*	*	0.7	2.0

White, female

2003.....	4.7	*	*	*	*	*	*	*	0.3	1.0
2002.....	4.9	*	*	*	*	*	*	*	0.3	1.0
2001.....	4.8	*	*	*	*	*	*	*	0.4	1.4
2000.....	4.8	*	*	*	*	*	*	*	0.4	0.9
1999.....	4.6	*	*	*	*	*	*	*	0.4	1.0

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	1.0	*	*	*	*	*	*	*	0.1
2000.....	1.0	*	*	*	*	*	*	*	0.1
1999.....	1.0	*	*	*	*	*	*	*	0.1

All races 3/, male

2003.....	0.9	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	*
2001.....	1.0	*	*	*	*	*	*	*	*
2000.....	0.9	*	*	*	*	*	*	*	*
1999.....	0.9	*	*	*	*	*	*	*	*

All races 3/, female

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.7	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*
2000.....	0.8	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*

Black, male

2003.....	0.7	*	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*	*

Black, female

2003.....	0.8	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*
2001.....	0.9	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes

2003.....	14.6	4.5	*	*	*	0.1	0.3	0.5	0.9	1.3
2002.....	14.2	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3
2001.....	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3
2000.....	13.5	4.2	*	*	*	0.2	0.3	0.4	0.8	1.2
1999.....	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2

All races 3/, male

2003.....	14.3	5.2	*	*	*	*	0.4	0.5	0.9	1.4
2002.....	13.9	5.3	*	*	*	*	0.3	0.5	0.8	1.4
2001.....	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4
2000.....	13.2	5.2	*	*	*	*	0.3	0.5	0.9	1.4
1999.....	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3

All races 3/, female

2003.....	14.9	3.8	*	*	*	*	0.3	0.5	0.8	1.1
2002.....	14.5	3.2	*	*	*	*	0.3	0.7	0.7	1.1
2001.....	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3
2000.....	13.8	3.1	*	*	*	*	0.3	0.4	0.8	1.1
1999.....	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2003.....	14.3	3.8	*	*	*	*	0.2	0.4	0.6	0.8
2002.....	13.9	3.6	*	*	*	*	0.2	0.4	0.6	0.8
2001.....	13.5	2.4	*	*	*	*	0.3	0.3	0.5	0.9
2000.....	13.1	2.9	*	*	*	0.1	0.3	0.3	0.6	0.8
1999.....	12.3	3.4	*	*	*	*	0.2	0.3	0.5	0.8

White, male

2003.....	14.0	4.5	*	*	*	*	0.3	0.4	0.6	0.8
2002.....	13.7	4.4	*	*	*	*	*	0.3	0.6	0.9
2001.....	13.3	2.8	*	*	*	*	*	0.3	0.4	1.1
2000.....	12.9	3.7	*	*	*	*	0.3	0.4	0.6	0.9
1999.....	12.1	4.1	*	*	*	*	*	0.3	0.5	0.9

White, female

2003.....	14.5	3.0	*	*	*	*	*	0.4	0.6	0.8
2002.....	14.1	2.7	*	*	*	*	*	0.5	0.5	0.8
2001.....	13.7	2.1	*	*	*	*	0.4	0.3	0.6	0.7
2000.....	13.2	2.0	*	*	*	*	0.3	0.3	0.5	0.7
1999.....	12.4	2.8	*	*	*	*	*	0.4	0.5	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2003.....	20.6	9.1	*	*	*	*	0.7	1.4	2.9	4.4
2002.....	19.8	8.2	*	*	*	*	0.9	1.9	2.1	4.0
2001.....	19.6	8.1	*	*	*	*	0.7	1.3	2.2	4.0
2000.....	19.6	12.2	*	*	*	*	*	1.1	2.7	4.5
1999.....	18.6	9.6	*	*	*	*	0.8	1.5	2.3	4.0

Black, male

2003.....	20.0	9.5	*	*	*	*	*	*	3.5	5.4
2002.....	19.1	10.2	*	*	*	*	*	1.6	2.2	4.4
2001.....	18.0	8.7	*	*	*	*	*	1.7	2.6	3.4
2000.....	18.3	14.2	*	*	*	*	*	*	3.0	5.2
1999.....	17.5	8.1	*	*	*	*	*	*	2.3	4.8

Black, female

2003.....	21.1	8.7	*	*	*	*	*	*	2.5	3.5
2002.....	20.5	6.1	*	*	*	*	*	2.1	2.1	3.6
2001.....	21.0	7.6	*	*	*	*	*	*	1.8	4.5
2000.....	20.7	10.1	*	*	*	*	*	*	2.4	3.9
1999.....	19.5	11.1	*	*	*	*	*	1.6	2.3	3.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

White, male

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

White, female

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

Black, male

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

Black, female

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Renal failure (N17-N19)

All races 3/, both sexes

2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.2	4.0	*	*	*	0.2	0.3	0.4	0.8	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1

All races 3/, male

2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.9	5.1	*	*	*	*	0.2	0.4	0.8	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3

All races 3/, female

2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.5	2.8	*	*	*	*	0.3	0.4	0.7	1.1
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Renal failure (N17-N19)

White, both sexes

2003.....	14.0	3.5	*	*	*	*	0.2	0.4	0.5	0.8
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.8	2.6	*	*	*	*	0.2	0.3	0.5	0.8
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7

White, male

2003.....	13.8	4.1	*	*	*	*	*	0.4	0.5	0.8
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.7	3.5	*	*	*	*	0.3	0.4	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8

White, female

2003.....	14.2	2.8	*	*	*	*	*	0.4	0.5	0.7
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	13.0	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Renal failure (N17-N19)

Black, both sexes

2003.....	20.3	8.9	*	*	*	*	0.6	1.3	2.8	4.4
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	19.2	11.7	*	*	*	*	*	1.0	2.6	4.2
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8

Black, male

2003.....	19.7	9.2	*	*	*	*	*	*	3.4	5.4
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	18.0	14.2	*	*	*	*	*	*	2.8	5.0
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5

Black, female

2003.....	20.9	8.7	*	*	*	*	*	*	2.3	3.5
2002.....	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000.....	20.3	9.1	*	*	*	*	*	*	2.3	3.5
1999.....	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

White, male

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

White, female

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

White, male

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

White, female

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

[illegible]

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

White, male

[illegible]

White, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.5	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	1.0	1.0	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2003.....	0.1	*	*	0.3	0.5	0.4	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.4	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	0.3	*	*	0.6	1.0	0.7	0.7
2002.....	0.2	*	*	0.4	0.6	0.7	0.5
2001.....	0.2	*	0.3	0.6	0.7	0.8	0.4
2000.....	0.2	*	*	0.6	0.8	0.7	0.6
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2003.....	0.5	*	*	1.0	1.2	1.8	1.4
2002.....	0.4	*	*	1.0	1.4	1.0	1.2
2001.....	0.4	*	*	1.1	1.4	1.1	1.0
2000.....	0.4	*	*	1.2	1.2	1.3	0.9
1999.....	0.4	*	*	1.1	1.4	1.2	0.9

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	1.0	*	*	2.1	2.3	3.5	2.7
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.8	*	*	2.3	2.3	2.4	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
All races 3/, both sexes										
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
White, both sexes										
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2003.....
2002.....
2001.....
2000.....
1999.....
White, female										
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
Black, both sexes										
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes

2003.....	0.2	*	0.1	0.4	0.5	0.5	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.3
2001.....	0.1	*	0.2	0.3	0.5	0.4	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.3
1999.....	0.1	*	0.1	0.3	0.4	0.4	0.4

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	0.3	*	0.2	0.8	1.1	1.1	0.9
2002.....	0.2	*	0.3	0.6	0.8	0.8	0.7
2001.....	0.3	*	0.3	0.7	0.9	0.9	0.6
2000.....	0.3	*	*	0.8	0.8	0.9	0.7
1999.....	0.3	*	0.3	0.6	0.9	0.8	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

White, both sexes

2003.....	0.1	*	*	0.3	0.4	0.3	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	0.3	*	*	0.6	0.9	0.7	0.6
2002.....	0.2	*	*	0.4	0.5	0.6	0.5
2001.....	0.2	*	*	0.5	0.6	0.7	0.3
2000.....	0.2	*	*	0.6	0.7	0.7	0.6
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, both sexes

2003.....	0.5	*	*	0.9	1.2	1.6	1.3
2002.....	0.4	*	*	0.9	1.4	0.9	1.0
2001.....	0.4	*	*	1.0	1.3	1.0	0.9
2000.....	0.3	*	*	1.0	1.0	1.1	0.8
1999.....	0.4	*	*	1.0	1.3	1.0	0.8

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	0.9	*	*	1.9	2.3	3.1	2.5
2002.....	0.7	*	*	1.8	2.6	1.8	2.0
2001.....	0.7	*	*	1.9	2.5	1.8	1.7
2000.....	0.7	*	*	1.9	1.9	2.1	1.4
1999.....	0.8	*	*	1.9	2.4	2.0	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2003.....	4.9	356.0	0.5	*	*	*	*	*	*	*
2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.1	361.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2003.....	5.7	397.4	0.5	*	*	*	*	*	*	*
2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.9	400.6	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2003.....	4.2	312.8	0.5	*	*	*	*	*	*	*
2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	321.2	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

White, both sexes

2003.....	3.8	285.1	0.4	*	*	*	*	*	*	*
2002.....	3.7	277.2	0.3	*	*	*	*	*	*	*
2001.....	3.7	268.5	0.4	*	*	*	*	*	*	*
2000.....	3.8	278.1	0.4	*	*	*	*	*	*	*
1999.....	3.8	285.9	0.5	*	*	*	*	*	*	*

White, male

2003.....	4.4	319.1	0.4	*	*	*	*	*	*	*
2002.....	4.3	309.4	*	*	*	*	*	*	*	*
2001.....	4.2	297.1	0.5	*	*	*	*	*	*	*
2000.....	4.3	307.0	0.5	*	*	*	*	*	*	*
1999.....	4.3	315.0	0.5	*	*	*	*	*	*	*

White, female

2003.....	3.2	249.5	0.3	*	*	*	*	*	*	*
2002.....	3.2	243.5	*	*	*	*	*	*	*	*
2001.....	3.1	238.6	*	*	*	*	*	*	*	*
2000.....	3.2	247.7	*	*	*	*	*	*	*	*
1999.....	3.2	255.3	0.4	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	149.3	3.3	1.0	1.0	1.1	1.2	1.2	1.4	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2

All races 3/, male

2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.1	154.1	3.3	1.0	1.0	1.3	1.4	1.3	1.6	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3

All races 3/, female

2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.6	144.2	3.2	1.0	1.0	0.9	0.9	1.0	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2003.....	3.5	137.6	3.2	0.9	1.0	1.1	1.0	1.1	1.1	1.2
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.7	144.2	2.9	0.9	0.9	1.0	1.2	1.2	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

White, male

2003.....	3.7	143.2	3.1	0.8	0.9	1.3	1.1	1.2	1.1	1.2
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	149.3	2.9	0.9	0.9	1.1	1.4	1.3	1.7	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

White, female

2003.....	3.4	131.8	3.2	0.9	1.0	0.8	1.0	0.9	1.1	1.1
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	138.7	2.8	1.0	0.9	0.9	0.9	1.0	1.2	1.1
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2003.....	4.5	157.5	4.7	1.1	1.1	1.2	1.7	1.2	1.5	1.2
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	5.1	189.2	5.7	1.4	1.5	1.8	1.3	1.5	1.5	1.7
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4

Black, male

2003.....	5.1	170.7	4.8	1.2	*	1.7	1.8	*	*	*
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.6	191.3	6.2	1.8	1.4	2.5	*	1.9	1.8	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*

Black, female

2003.....	3.9	143.8	4.6	*	*	*	1.5	*	1.6	*
2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.6	187.0	5.1	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.6	94.6	1.6	0.4	0.5	1.6	3.0	3.7	4.6	5.8
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.8	109.7	1.8	0.5	0.5	1.9	4.1	5.1	6.2	7.7
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.4	78.8	1.5	0.4	0.4	1.3	1.8	2.3	2.9	3.8
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2003.....	11.1	68.1	1.4	0.4	0.4	1.5	2.5	3.1	3.4	4.7
2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.6	75.7	1.5	0.4	0.4	1.5	2.8	3.5	4.4	5.7
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

White, male

2003.....	10.1	82.4	1.4	0.3	0.4	2.0	3.4	4.2	4.5	6.1
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.5	87.9	1.7	0.4	0.4	1.9	4.0	4.9	6.1	7.8
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

White, female

2003.....	12.0	53.3	1.3	0.4	0.3	1.0	1.6	1.9	2.1	3.2
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.7	62.9	1.3	0.3	0.4	1.1	1.6	2.1	2.7	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2003.....	12.3	159.3	3.1	*	1.0	1.5	3.4	4.9	5.0	7.1
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	14.2	208.1	2.6	0.7	0.8	2.2	4.3	5.5	6.6	7.4
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8

Black, male

2003.....	13.5	170.7	3.4	*	1.3	1.8	4.2	7.3	6.2	9.6
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	16.1	239.5	2.6	*	*	2.1	5.5	7.4	8.6	8.4
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4

Black, female

2003.....	11.3	147.5	2.9	*	*	*	2.7	2.7	3.9	4.9
2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.5	175.5	2.6	*	*	2.3	3.1	3.8	4.8	6.5
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2003.....	69.3	29.7	4.0	2.1	2.4	4.1	5.8	7.4	9.3	14.5
2002.....	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1
2001.....	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1
2000.....	64.8	28.7	4.0	1.9	2.5	3.7	5.6	7.0	9.4	14.2
1999.....	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0

All races 3/, male

2003.....	57.3	34.8	4.2	2.3	2.8	4.6	6.6	8.2	10.4	16.5
2002.....	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4
2001.....	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8
2000.....	53.7	32.0	4.4	2.0	2.7	4.2	6.6	8.3	11.1	16.9
1999.....	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8

All races 3/, female

2003.....	81.0	24.5	3.8	1.8	2.1	3.4	4.9	6.5	8.1	12.5
2002.....	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9
2001.....	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5
2000.....	75.5	25.3	3.7	1.8	2.3	3.1	4.5	5.8	7.8	11.5
1999.....	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2003.....	73.9	25.5	3.7	2.0	2.3	4.0	5.5	6.7	8.3	13.0
2002.....	71.8	25.2	3.9	1.9	2.2	3.4	5.0	6.0	8.1	12.8
2001.....	69.4	23.7	3.9	2.0	2.2	3.6	4.9	5.8	7.9	12.6
2000.....	68.1	24.4	3.5	1.8	2.4	3.5	5.2	6.2	8.4	12.5
1999.....	65.0	25.2	4.0	2.0	2.3	3.8	4.8	5.9	8.1	12.1

White, male

2003.....	59.9	31.5	3.8	2.4	2.6	4.7	6.6	7.8	9.6	15.2
2002.....	58.2	30.8	4.1	2.2	2.3	4.1	5.9	6.8	9.9	15.5
2001.....	55.8	25.9	4.4	1.9	2.4	4.4	5.8	6.6	8.9	15.5
2000.....	55.2	26.4	3.9	1.9	2.7	4.2	6.4	7.8	10.2	15.3
1999.....	52.7	29.2	4.3	2.0	2.2	4.8	5.5	7.0	9.8	15.1

White, female

2003.....	87.6	19.3	3.7	1.6	2.0	3.1	4.2	5.4	6.8	10.7
2002.....	85.2	19.3	3.6	1.6	2.1	2.8	4.1	5.0	6.1	9.9
2001.....	82.7	21.4	3.4	2.0	2.0	2.8	3.9	4.8	6.8	9.7
2000.....	80.6	22.3	3.1	1.6	2.1	2.9	4.0	4.7	6.5	9.6
1999.....	77.0	20.9	3.7	1.9	2.3	2.8	4.1	4.7	6.3	9.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,
N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2003.....	60.9	51.8	5.3	2.2	3.2	4.9	8.1	13.3	16.9	27.2
2002.....	59.7	45.2	5.9	2.2	2.9	4.3	9.4	12.2	16.6	25.7
2001.....	59.1	51.3	5.4	2.0	2.9	5.3	7.7	11.5	16.6	26.6
2000.....	60.3	52.4	6.7	2.6	3.5	4.9	8.4	13.3	17.8	27.4
1999.....	56.2	56.7	5.3	2.9	3.1	4.7	8.9	10.4	15.4	28.7

Black, male

2003.....	55.7	55.3	6.1	1.8	3.9	4.7	7.2	12.9	17.1	28.7
2002.....	54.8	48.8	6.4	2.3	3.4	4.2	9.4	12.7	17.9	26.4
2001.....	54.6	58.4	5.1	1.5	3.3	5.6	7.7	12.3	17.5	29.7
2000.....	56.4	63.1	6.9	2.2	3.2	5.1	9.0	13.3	19.2	30.5
1999.....	52.8	64.7	5.4	3.2	3.8	4.8	9.7	10.5	16.2	31.5

Black, female

2003.....	65.6	48.2	4.5	2.7	2.5	5.2	9.0	13.6	16.7	25.9
2002.....	64.1	41.5	5.4	2.2	2.5	4.3	9.5	11.8	15.5	25.0
2001.....	63.1	43.8	5.6	2.5	2.5	5.1	7.6	10.8	15.7	24.0
2000.....	63.8	41.2	6.4	2.9	3.7	4.7	7.8	13.3	16.6	24.7
1999.....	59.3	48.3	5.1	2.6	2.4	4.5	8.1	10.3	14.6	26.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	35.6	22.9	12.1	7.0	8.0	34.0	39.8	33.0	30.0	33.5
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2

All races 3/, male

2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	47.4	25.5	14.2	8.1	10.2	46.6	60.6	51.8	45.9	50.0
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4

All races 3/, female

2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	24.2	20.2	9.8	5.9	5.7	20.7	18.3	14.4	14.4	17.3
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2003.....	39.5	20.0	10.6	5.2	7.0	35.9	44.3	34.6	31.6	36.2
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	36.5	18.8	11.1	6.4	7.8	36.4	41.4	33.5	30.4	33.9
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

White, male

2003.....	51.3	22.5	12.8	6.0	9.0	48.4	66.2	51.9	46.8	51.4
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	48.0	21.3	13.1	7.4	9.9	49.3	63.1	52.1	46.5	50.4
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5

White, female

2003.....	28.0	17.4	8.3	4.4	4.8	22.7	20.7	16.1	15.7	20.6
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.4	16.1	9.0	5.4	5.7	22.6	18.6	14.7	14.2	17.3
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2003.....	32.4	43.5	13.4	7.3	8.1	20.7	30.6	29.8	29.9	35.5
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	34.8	47.0	18.2	10.9	9.2	24.1	34.4	34.5	33.1	36.5
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6

Black, male

2003.....	46.1	47.6	15.7	8.7	10.6	30.0	47.6	46.6	46.9	54.9
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	50.9	50.3	21.1	13.3	12.7	34.9	52.2	56.0	51.9	56.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4

Black, female

2003.....	19.9	39.3	11.0	5.9	5.5	11.2	13.7	14.3	14.7	18.3
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	20.2	43.7	15.2	8.5	5.6	13.0	17.0	14.8	16.4	18.7
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	17.0	4.5	4.6	4.3	5.4	27.5	30.0	22.2	17.6	17.5
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9

All races 3/, male

2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.9	4.8	5.1	4.6	6.6	36.1	44.6	33.9	26.3	25.3
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0

All races 3/, female

2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.4	4.2	4.1	3.9	4.2	18.4	15.0	10.5	9.0	9.7
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Transport accidents (V01-V99,Y85)

White, both sexes

2003.....	17.0	3.6	4.0	3.5	5.2	29.2	30.4	20.7	16.8	17.0
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.3	4.0	4.3	4.1	5.6	29.9	31.0	22.1	17.6	17.6
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8

White, male

2003.....	23.9	4.0	4.5	3.9	6.5	37.6	44.4	30.6	25.1	24.2
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	24.1	4.4	4.7	4.5	6.7	38.9	46.0	33.4	26.2	25.3
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7

White, female

2003.....	10.3	3.1	3.4	3.1	3.9	20.2	15.3	10.2	8.2	9.6
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.7	3.6	3.8	3.8	4.4	20.4	15.2	10.6	9.0	9.9
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Transport accidents (V01-V99,Y85)

Black, both sexes

2003.....	15.1	4.2	5.0	3.6	4.5	16.2	23.2	20.3	18.3	17.9
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.5	6.9	6.3	5.1	5.0	17.7	26.7	24.5	20.2	18.2
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9

Black, male

2003.....	22.6	*	5.1	4.5	5.7	22.7	36.3	31.8	29.4	28.4
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	25.0	7.1	7.0	5.5	6.8	24.5	39.8	39.5	32.2	29.1
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7

Black, female

2003.....	8.3	*	5.0	2.8	3.4	9.5	10.2	9.8	8.2	8.4
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.9	*	5.7	4.6	3.2	10.6	13.9	10.8	9.5	8.5
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.7	4.4	4.3	3.9	5.0	26.4	28.7	20.8	16.2	15.9
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1

All races 3/, male

2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.9	4.6	4.8	4.2	6.0	34.5	42.4	31.5	24.0	22.6
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2

All races 3/, female

2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.9	4.1	3.8	3.6	3.9	17.8	14.6	10.2	8.5	9.3
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2003.....	15.8	3.5	3.8	3.3	4.8	28.1	29.1	19.5	15.7	15.6
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	16.0	3.8	4.0	3.8	5.1	28.8	29.6	20.6	16.2	16.0
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

White, male

2003.....	22.0	3.9	4.3	3.6	6.0	36.1	42.3	28.8	23.3	21.9
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	22.0	4.1	4.5	4.1	6.1	37.3	43.6	30.8	23.8	22.5
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

White, female

2003.....	9.8	3.0	3.2	2.9	3.6	19.6	14.8	9.6	7.7	9.0
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.2	3.4	3.6	3.5	4.1	19.7	14.8	10.2	8.5	9.5
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2003.....	14.2	4.2	4.9	3.4	4.3	15.5	22.3	19.5	17.3	16.7
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.6	6.9	6.0	4.8	4.8	16.8	25.9	23.7	18.7	16.8
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2

Black, male

2003.....	21.1	*	5.0	4.1	5.2	21.6	34.6	30.5	27.7	26.3
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	23.4	7.1	6.5	5.1	6.5	23.0	38.3	38.0	29.9	26.5
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Black, female

2003.....	8.0	*	4.8	2.7	3.4	9.2	10.1	9.4	8.0	8.2
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.5	*	5.5	4.5	3.1	10.3	13.7	10.7	8.9	8.1
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2003.....	0.5	*	*	0.1	0.2	0.6	0.7	0.5	0.5	0.5
2002.....	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6
2000.....	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

All races 3/, male

2003.....	0.7	*	*	*	0.2	0.9	1.0	0.8	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8
2001.....	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1
2000.....	0.8	*	*	0.2	0.4	1.0	1.0	0.9	0.9	1.1
1999.....	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4

All races 3/, female

2003.....	0.2	*	*	*	*	0.3	0.2	0.2	*	0.2
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	0.2	*	*	*	0.2
2000.....	0.3	*	*	*	*	0.3	0.2	*	0.3	0.3
1999.....	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2003.....	0.5	*	*	*	0.2	0.7	0.7	0.6	0.5	0.5
2002.....	0.4	*	*	*	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	*	0.2	0.5	0.6	0.5	0.4	0.5
2000.....	0.5	*	0.2	0.2	0.3	0.7	0.6	0.5	0.6	0.6
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

White, male

2003.....	0.7	*	*	*	0.3	1.0	1.1	0.9	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.9	0.5	0.6	0.8
2001.....	0.7	*	*	*	0.3	0.7	0.9	0.8	0.7	0.9
2000.....	0.8	*	*	*	0.4	1.0	1.0	0.9	0.9	1.1
1999.....	1.1	*	*	0.4	0.5	1.2	1.7	1.3	1.1	1.4

White, female

2003.....	0.2	*	*	*	*	0.4	0.3	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	0.3	*	*	0.3	0.2
1999.....	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2003.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.6	0.9
2002.....	0.7	*	*	0.1	0.2	0.4	0.6	0.5	0.7	0.9
2001.....	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9
2000.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9

All races 3/, male

2003.....	1.1	*	*	*	0.2	0.6	1.0	0.8	1.0	1.4
2002.....	1.1	*	*	*	0.2	0.6	0.9	0.9	1.2	1.5
2001.....	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4
2000.....	1.2	*	*	*	0.2	0.7	1.2	1.5	1.4	1.6
1999.....	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4

All races 3/, female

2003.....	0.2	*	*	*	*	*	0.2	0.3	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2001.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2
1999.....	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2003.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.7	0.9
2002.....	0.7	*	*	*	0.2	0.5	0.6	0.6	0.8	0.9
2001.....	0.8	*	0.2	0.1	0.2	0.5	0.7	0.9	1.0	1.0
2000.....	0.8	*	*	*	0.2	0.5	0.8	1.0	0.9	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.7	1.0	0.9	1.0

White, male

2003.....	1.2	*	*	*	*	0.6	1.0	1.0	1.1	1.5
2002.....	1.2	*	*	*	*	0.6	1.1	1.0	1.4	1.6
2001.....	1.3	*	*	*	*	0.8	1.1	1.4	1.6	1.5
2000.....	1.3	*	*	*	*	0.6	1.4	1.7	1.5	1.7
1999.....	1.2	*	*	*	0.3	0.6	1.1	1.5	1.4	1.5

White, female

2003.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	0.2	*	*	*	*	0.3	*	*	*	0.3
2001.....	0.3	*	*	*	*	*	0.4	0.5	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	*	*
1999.....	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	0.7
2001.....	0.5	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	0.7

Black, male

2003.....	0.7	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*
2001.....	0.8	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*

Black, female

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

2003.....	21.0	19.9	6.8	2.0	2.1	6.3	12.3	12.7	13.7	18.3
2002.....	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8
2001.....	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7
2000.....	18.6	18.4	7.5	2.8	2.6	6.4	9.8	10.9	12.4	16.1
1999.....	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3

All races 3/, male

2003.....	26.0	22.4	8.5	2.5	3.0	10.0	19.4	19.6	20.2	26.2
2002.....	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9
2001.....	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8
2000.....	23.5	20.7	9.1	3.6	3.6	10.4	16.0	17.9	19.6	24.7
1999.....	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4

All races 3/, female

2003.....	16.2	17.3	5.0	1.6	1.2	2.3	4.9	5.5	7.0	10.4
2002.....	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7
2001.....	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5
2000.....	13.9	15.9	5.8	2.0	1.5	2.2	3.3	3.9	5.4	7.6
1999.....	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nontransport accidents (W00-X59,Y86)

White, both sexes

2003.....	22.5	16.4	6.6	1.7	1.8	6.8	13.9	13.9	14.8	19.2
2002.....	21.4	15.9	5.9	1.8	2.0	6.9	12.3	12.7	14.6	18.3
2001.....	20.0	16.9	6.7	2.2	2.1	6.3	11.3	11.0	12.2	17.2
2000.....	19.2	14.8	6.8	2.2	2.2	6.5	10.4	11.4	12.8	16.3
1999.....	19.0	14.3	7.3	2.2	2.4	6.2	9.7	10.4	12.1	15.6

White, male

2003.....	27.4	18.6	8.3	2.1	2.5	10.8	21.7	21.3	21.7	27.2
2002.....	26.2	19.2	7.4	2.4	2.8	10.9	19.7	19.7	22.0	26.4
2001.....	24.5	18.9	8.4	2.7	2.8	10.0	18.4	17.3	18.6	25.5
2000.....	23.8	17.0	8.4	2.9	3.2	10.5	17.1	18.7	20.3	25.1
1999.....	23.5	16.5	8.8	2.7	3.4	10.0	15.8	16.9	19.2	23.8

White, female

2003.....	17.7	14.2	4.9	1.3	0.9	2.5	5.4	5.9	7.6	11.0
2002.....	16.8	12.5	4.4	1.2	1.2	2.7	4.3	5.3	6.9	10.1
2001.....	15.6	14.8	4.8	1.7	1.3	2.3	3.8	4.4	5.6	8.6
2000.....	14.8	12.5	5.2	1.6	1.3	2.2	3.3	4.1	5.3	7.4
1999.....	14.6	11.8	5.8	1.6	1.3	2.2	3.2	3.7	4.8	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2003.....	17.2	39.3	8.4	3.7	3.6	4.5	7.4	9.5	11.6	17.7
2002.....	17.8	43.6	9.2	4.7	4.4	6.0	7.4	9.8	12.1	18.9
2001.....	17.6	40.2	8.4	4.1	4.1	5.9	7.2	9.4	11.1	17.7
2000.....	18.2	40.2	11.9	5.9	4.2	6.5	7.7	10.0	12.9	18.3
1999.....	18.9	34.4	10.6	5.6	4.6	6.3	8.8	10.2	14.4	16.7

Black, male

2003.....	23.5	44.0	10.6	4.2	5.0	7.3	11.3	14.8	17.5	26.4
2002.....	24.8	45.9	12.1	6.1	6.5	10.1	12.4	16.2	18.8	29.1
2001.....	24.4	46.4	9.7	6.0	5.7	9.6	11.8	15.1	16.5	26.1
2000.....	25.8	43.2	14.2	7.8	5.9	10.4	12.4	16.5	19.7	27.6
1999.....	26.8	40.0	13.0	7.4	6.6	10.8	15.5	16.7	22.3	25.7

Black, female

2003.....	11.6	34.3	6.0	3.1	2.1	1.7	3.5	4.5	6.4	9.8
2002.....	11.4	41.2	6.2	3.3	2.3	1.7	2.4	4.0	6.2	9.8
2001.....	11.5	33.7	7.1	2.2	2.5	2.1	2.7	4.3	6.2	10.2
2000.....	11.4	37.1	9.5	3.9	2.4	2.4	3.1	4.0	6.9	10.1
1999.....	11.7	28.5	8.1	3.7	2.5	1.6	2.5	4.3	7.3	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Falls (W00-W19)

All races 3/, both sexes

2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.8	*	0.2	*	0.1	0.5	0.7	0.8	0.8	1.1
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2

All races 3/, male

2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.3	*	0.3	*	*	0.9	1.2	1.4	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0

All races 3/, female

2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.4	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Falls (W00-W19)

White, both sexes

2003.....	6.7	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.4	*	0.2	*	*	0.6	0.9	0.9	0.9	1.1
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3

White, male

2003.....	6.9	*	0.5	*	*	0.8	1.3	1.4	1.2	2.1
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.8	*	*	*	*	1.0	1.4	1.6	1.6	1.8
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0

White, female

2003.....	6.5	*	*	*	*	*	0.3	*	0.3	0.3
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.2	0.3
2002.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.2
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4
2000.....	0.3	*	*	*	0.3	0.5	0.4	0.4	0.3	0.3
1999.....	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3

2003.....	0.4	*	*	*	0.3	0.8	0.8	0.6	0.4	0.5
2002.....	0.4	*	*	*	0.3	0.8	0.8	0.6	0.5	0.4
2001.....	0.5	*	*	*	0.4	0.9	0.9	0.6	0.5	0.6
2000.....	0.5	*	*	*	0.5	0.9	0.8	0.6	0.4	0.5
1999.....	0.5	*	*	*	0.5	1.0	1.0	0.6	0.6	0.5

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.3	1.9	3.3	1.0	0.9	1.9	1.5	1.2	1.1	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1

All races 3/, male

2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.1	1.4	1.3	3.3	2.6	2.1	1.9	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9

All races 3/, female

2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.7	0.5	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

White, both sexes

2003.....	1.1	1.4	3.1	0.6	0.5	1.3	1.3	1.0	0.8	1.0
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1

White, male

2003.....	1.8	1.5	4.0	0.8	0.7	2.3	2.4	1.7	1.3	1.7
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.1	0.8	2.8	2.4	2.0	1.8	1.9
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8

White, female

2003.....	0.5	1.3	2.2	0.3	*	*	*	*	*	0.3
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

Black, both sexes

2003.....	1.3	*	2.1	1.0	1.6	1.8	1.4	1.1	1.1	0.8
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.7	4.5	2.5	2.2	2.2	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1

Black, male

2003.....	2.2	*	3.0	1.4	2.4	3.4	2.6	2.1	2.2	1.7
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.8	*	3.1	3.1	3.3	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2

Black, female

2003.....	0.4	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	1.6	*	1.3	*	*	*	*	*
2000.....	0.6	*	1.9	*	*	*	*	*	*	*
1999.....	0.5	*	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.7	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9

All races 3/, male

2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.1	0.4	0.4	0.8	0.9	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2

All races 3/, female

2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	1.0	*	1.6	0.8	0.5	0.4	0.4	0.5	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, both sexes

2003.....	1.1	0.7	1.2	0.5	0.2	0.3	0.6	0.6	0.7	0.8
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.1	0.8	1.3	0.7	0.3	0.4	0.6	0.6	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8

White, male

2003.....	1.3	*	1.4	0.5	*	0.4	0.8	0.7	0.8	1.1
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.8	0.3	0.4	0.8	0.8	0.7	1.1
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0

White, female

2003.....	0.9	*	1.0	0.5	0.3	0.3	0.5	0.5	0.5	0.5
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2003.....	2.0	*	3.0	1.9	1.0	0.6	*	0.9	0.8	1.0
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.6	*	5.4	2.4	0.9	*	1.0	1.2	1.3	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9

Black, male

2003.....	2.4	*	3.2	1.9	*	*	*	*	*	*
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.3	*	6.3	2.9	*	*	*	*	1.6	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8

Black, female

2003.....	1.7	*	2.7	1.8	*	*	*	*	*	*
2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.6	1.8	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.6	*	0.2	*	0.1	1.8	4.4	5.5	7.2	9.7
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1

All races 3/, male

2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.8	*	0.3	*	*	2.7	6.9	8.8	10.7	14.0
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3

All races 3/, female

2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.6	*	*	*	*	0.8	1.8	2.2	3.6	5.4
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2003.....	7.1	*	0.3	*	0.1	3.0	8.5	9.1	10.2	13.3
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	5.0	6.0	7.5	10.0
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3

White, male

2003.....	9.7	*	*	*	*	4.6	12.9	13.4	14.4	17.7
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.8	*	*	*	*	3.1	8.0	9.5	11.3	14.5
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7

White, female

2003.....	4.5	*	*	*	*	1.4	3.8	4.5	5.8	8.7
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.6	*	*	*	*	0.9	1.8	2.5	3.7	5.3
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2003.....	6.3	*	*	*	*	*	2.3	4.0	7.3	11.8
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.7	*	*	*	*	0.7	2.1	4.4	7.1	11.0
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3

Black, male

2003.....	8.8	*	*	*	*	*	3.5	5.7	10.2	16.7
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.5	*	*	*	*	*	2.7	7.4	10.0	15.1
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9

Black, female

2003.....	3.9	*	*	*	*	*	*	2.4	4.8	7.4
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.2	*	*	*	*	*	1.5	1.8	4.5	7.2
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2003.....	5.9	16.9	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.6
2002.....	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6
2001.....	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9
2000.....	6.3	14.9	1.8	0.6	0.8	1.4	2.0	2.3	2.4	3.0
1999.....	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7

All races 3/, male

2003.....	6.9	19.5	2.2	0.6	0.9	1.8	3.3	3.3	3.4	4.0
2002.....	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3
2001.....	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8
2000.....	7.4	16.6	2.1	0.7	1.2	2.2	3.6	4.1	4.2	5.2
1999.....	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5

All races 3/, female

2003.....	4.9	14.1	1.3	0.3	0.2	0.3	0.7	0.7	0.6	1.1
2002.....	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0
2001.....	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1
2000.....	5.3	13.0	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2003.....	6.3	13.6	1.7	0.5	0.6	1.2	2.1	2.1	2.2	2.6
2002.....	6.3	12.9	1.4	0.5	0.8	1.4	2.0	2.2	2.3	2.7
2001.....	6.6	13.4	1.7	0.6	0.7	1.4	2.2	1.9	2.2	3.0
2000.....	6.7	11.9	1.6	0.6	0.8	1.4	2.1	2.4	2.6	3.1
1999.....	6.9	11.2	1.8	0.7	0.9	1.6	2.2	2.2	2.3	2.9

White, male

2003.....	7.3	15.7	2.2	0.6	0.9	2.0	3.6	3.6	3.6	4.1
2002.....	7.3	15.3	1.8	0.7	1.0	2.4	3.4	3.6	3.8	4.3
2001.....	7.5	15.2	2.0	0.7	1.0	2.1	3.7	3.2	3.7	5.0
2000.....	7.6	13.7	2.0	0.7	1.2	2.3	3.7	4.2	4.4	5.3
1999.....	7.8	12.9	2.1	1.0	1.3	2.5	3.7	3.9	3.8	4.8

White, female

2003.....	5.3	11.4	1.2	0.3	*	0.4	0.6	0.6	0.7	1.1
2002.....	5.4	10.5	1.1	0.4	0.5	0.5	0.5	0.7	0.6	1.0
2001.....	5.8	11.5	1.4	0.5	0.3	0.6	0.6	0.6	0.6	1.0
2000.....	5.8	10.1	1.2	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	6.1	9.5	1.4	0.4	0.5	0.6	0.5	0.6	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2003.....	4.9	34.9	2.0	*	*	0.7	1.9	2.0	1.6	2.7
2002.....	5.2	37.5	2.4	0.7	0.9	1.0	1.2	1.5	1.9	3.0
2001.....	5.4	34.2	2.5	*	0.7	0.8	1.3	1.8	1.9	2.6
2000.....	5.6	32.3	2.6	0.8	0.8	1.3	1.8	2.1	2.5	3.2
1999.....	6.1	27.9	2.5	0.7	0.7	1.0	1.6	2.2	2.6	2.2

Black, male

2003.....	6.4	40.5	2.8	*	*	*	2.6	3.0	2.9	4.6
2002.....	6.6	38.9	3.3	*	1.5	1.5	1.9	2.3	3.4	5.2
2001.....	6.8	40.4	2.8	*	*	*	2.0	2.8	3.6	3.7
2000.....	7.6	33.7	2.8	*	1.2	1.9	3.2	3.6	4.0	5.6
1999.....	8.0	33.2	3.4	*	1.2	1.8	3.0	3.5	4.4	3.5

Black, female

2003.....	3.6	29.1	*	*	*	*	1.3	*	*	*
2002.....	4.0	36.0	*	*	*	*	*	*	*	*
2001.....	4.1	27.7	2.2	*	*	*	*	*	*	1.5
2000.....	3.9	30.8	2.3	*	*	*	*	*	*	*
1999.....	4.4	22.5	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 4/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.7	*	1.5	8.2	12.8	13.1	12.5	14.0
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4

All races 3/, male

2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 4/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.5	*	2.3	13.2	22.0	22.0	20.4	22.2
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7

All races 3/, female

2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 4/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.1	*	0.6	2.8	3.3	4.3	4.8	5.9
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2003.....	12.1	*	1.1	7.9	12.5	12.6	14.7	15.3
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 4/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.7	*	1.6	8.6	13.3	13.7	13.6	15.4
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7

White, male

2003.....	19.5	*	1.7	12.7	21.1	20.3	23.3	23.9
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 4/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	19.1	*	2.4	13.9	22.8	22.9	22.0	24.0
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3

White, female

2003.....	4.7	*	0.5	2.9	3.3	4.3	5.7	6.5
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 4/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.5	*	0.7	2.9	3.3	4.4	5.2	6.6
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Black, both sexes

2003.....	5.1	*	1.2	3.8	10.5	9.5	9.1	7.5
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 4/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.6	*	1.3	5.6	11.1	10.3	7.9	7.4
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4

Black, male

2003.....	8.8	*	1.9	6.6	18.0	17.4	15.7	12.9
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 4/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.8	*	2.2	9.7	19.2	18.1	14.3	13.2
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6

Black, female

2003.....	1.8	*	*	*	3.1	2.2	3.1	2.6
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 4/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.8	*	*	1.5	3.1	3.1	2.3	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	6.0	*	0.6	4.5	7.4	6.9	6.3	6.8
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1

All races 3/, male

2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.7	*	0.9	7.8	13.3	12.2	10.7	11.3
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8

All races 3/, female

2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.6	1.9	2.4
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2003.....	6.6	*	0.4	4.0	6.7	6.1	6.7	6.8
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.7	*	0.7	4.7	7.7	7.2	6.9	7.6
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8

White, male

2003.....	11.7	*	0.7	6.9	11.8	10.5	11.3	11.3
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.9	*	1.1	8.0	13.8	12.6	11.7	12.4
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7

White, female

2003.....	1.6	*	*	0.9	1.2	1.5	1.8	2.1
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.2	1.4	1.7	2.1	2.8
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, both sexes

2003.....	2.6	*	*	2.1	6.5	5.5	4.8	2.8
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.1	*	*	3.8	6.9	6.2	4.2	3.5
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2

Black, male

2003.....	5.0	*	*	3.9	11.9	11.0	9.1	5.5
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.8	*	*	7.0	12.5	11.7	7.6	6.5
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1

Black, female

2003.....	0.5	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2003.....	5.0	*	0.8	3.7	5.6	6.1	7.4	7.8
2002.....	5.0	*	0.8	3.8	5.7	6.4	6.9	7.7
2001 4/.....	4.8	*	0.9	3.8	5.4	5.9	6.7	7.9
2000.....	4.6	*	1.0	3.6	5.4	6.3	6.2	7.1
1999.....	4.5	*	0.7	3.2	5.1	5.7	6.7	7.3

All races 3/, male

2003.....	7.3	*	1.2	5.4	8.8	9.3	11.0	11.7
2002.....	7.3	*	1.2	5.8	8.9	9.7	10.6	11.4
2001 4/.....	7.1	*	1.3	5.8	8.7	9.3	10.2	11.9
2000.....	6.8	*	1.5	5.5	8.7	9.8	9.7	10.8
1999.....	6.6	*	1.1	4.6	8.0	8.8	10.2	10.9

All races 3/, female

2003.....	2.8	*	0.4	1.9	2.2	2.7	3.7	4.0
2002.....	2.9	*	0.4	1.6	2.3	2.9	3.1	4.0
2001 4/.....	2.6	*	0.4	1.7	2.0	2.5	3.1	3.9
2000.....	2.6	*	0.4	1.7	2.0	2.7	2.8	3.5
1999.....	2.5	*	0.3	1.6	2.0	2.5	3.2	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2003.....	5.5	*	0.8	4.0	5.8	6.4	8.0	8.5
2002.....	5.5	*	0.8	4.2	6.2	6.8	7.4	8.5
2001 4/.....	5.3	*	0.9	4.2	5.9	6.3	7.2	8.6
2000.....	5.0	*	0.9	3.9	5.6	6.6	6.7	7.8
1999.....	4.9	*	0.7	3.3	5.4	5.9	7.1	7.9

White, male

2003.....	7.9	*	1.1	5.8	9.3	9.8	11.9	12.6
2002.....	8.0	*	1.2	6.5	9.7	10.2	11.2	12.3
2001 4/.....	7.7	*	1.3	6.4	9.4	9.8	10.8	12.8
2000.....	7.3	*	1.4	5.9	9.0	10.4	10.3	11.6
1999.....	7.0	*	1.1	4.9	8.5	9.1	10.7	11.6

White, female

2003.....	3.1	*	0.4	2.0	2.2	2.8	4.0	4.3
2002.....	3.2	*	0.5	1.8	2.5	3.1	3.4	4.5
2001 4/.....	2.9	*	0.4	1.8	2.2	2.6	3.4	4.3
2000.....	2.8	*	0.5	1.8	2.0	2.7	3.1	3.8
1999.....	2.8	*	0.3	1.7	2.1	2.5	3.3	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes

2003.....	2.5	*	1.0	1.7	4.0	4.0	4.3	4.6
2002.....	2.3	*	0.7	1.6	3.2	4.2	4.2	4.1
2001 4/.....	2.3	*	0.8	1.5	3.0	3.8	4.5	4.1
2000.....	2.5	*	1.1	1.8	4.2	4.1	3.7	3.9
1999.....	2.3	*	0.7	1.9	3.1	3.9	4.6	4.2

Black, male

2003.....	3.8	*	1.6	2.6	6.1	6.4	6.7	7.4
2002.....	3.7	*	1.1	2.6	5.0	7.0	7.5	7.0
2001 4/.....	3.7	*	1.1	2.3	5.5	6.8	7.4	6.7
2000.....	4.0	*	1.9	2.6	6.8	6.4	6.7	6.7
1999.....	3.7	*	*	2.9	5.3	6.7	7.6	7.5

Black, female

2003.....	1.3	*	*	*	1.9	1.8	2.2	2.2
2002.....	1.1	*	*	*	1.5	1.6	1.4	1.6
2001 4/.....	1.1	*	*	*	*	*	1.8	1.8
2000.....	1.1	*	*	*	1.7	1.9	*	1.4
1999.....	1.0	*	*	*	*	*	1.9	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races 3/, both sexes

2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 4/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.1	9.1	2.3	0.7	1.2	9.6	16.4	13.0	9.5	7.8
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7

All races 3/, male

2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 4/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.5	10.3	2.6	0.8	1.5	15.8	27.5	21.6	14.9	11.1
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0

All races 3/, female

2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 4/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.8	2.1	0.7	0.8	3.1	4.8	4.4	4.1	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2003.....	3.7	6.9	1.7	0.4	0.6	5.0	8.4	6.9	5.3	5.0
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 4/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.7	6.5	1.7	0.6	0.9	5.2	7.9	6.7	5.5	5.0
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1

White, male

2003.....	5.4	8.1	1.5	0.4	0.9	8.0	13.3	10.7	7.8	6.9
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 4/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.3	8.1	1.9	0.7	1.1	8.1	12.2	10.1	8.1	6.7
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0

White, female

2003.....	2.0	5.7	1.8	0.4	0.3	1.8	3.2	2.9	2.6	3.0
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 4/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	4.9	1.4	0.5	0.7	2.1	3.3	3.2	3.0	3.3
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2003.....	22.0	16.1	6.1	1.7	2.7	33.2	62.4	54.5	36.3	25.4
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 4/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	22.3	24.0	6.5	1.3	2.6	33.7	64.4	50.3	34.8	26.4
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1

Black, male

2003.....	38.9	17.8	7.4	2.0	3.7	58.9	111.6	101.6	63.9	43.7
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 4/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	38.6	24.6	6.6	*	3.5	57.9	116.3	92.6	61.4	43.3
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1

Black, female

2003.....	6.6	14.2	4.8	1.3	1.7	6.7	13.6	10.9	11.4	9.0
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 4/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.5	23.4	6.4	1.6	1.8	8.6	13.5	11.5	11.2	11.4
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.9	*	0.2	0.3	0.7	7.8	13.1	10.0	6.6	4.8
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6

All races 3/, male

2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.7	*	*	0.3	1.1	13.5	23.0	17.7	11.1	7.6
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1

All races 3/, female

2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.4	2.1	2.0
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2003.....	2.2	*	0.2	0.1	0.4	3.8	6.3	4.9	3.6	3.0
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.3	0.5	3.9	5.6	4.6	3.5	2.9
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0

White, male

2003.....	3.5	*	*	*	0.6	6.4	10.6	8.2	5.7	4.4
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.3	*	*	0.3	0.7	6.5	9.2	7.4	5.4	4.2
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3

White, female

2003.....	0.9	*	*	*	*	1.0	1.7	1.4	1.4	1.6
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.7	1.5	1.6
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2003.....	16.8	*	*	0.7	2.0	29.0	55.3	47.9	30.1	18.1
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	16.1	*	*	*	1.8	29.1	55.6	42.4	27.0	17.6
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9

Black, male

2003.....	31.6	*	*	*	2.9	53.2	101.7	92.4	56.4	33.9
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	30.3	*	*	*	2.9	52.3	103.9	81.5	50.4	32.2
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3

Black, female

2003.....	3.3	*	*	*	1.1	4.0	9.1	6.8	6.3	4.1
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.3	*	*	*	*	5.1	8.2	6.5	6.2	4.6
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2003.....	2.0	8.3	2.1	0.4	0.3	1.7	2.9	2.6	2.3	2.7
2002.....	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7
2001 4/.....	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2
2000.....	2.2	8.8	2.2	0.5	0.5	1.8	3.3	2.9	2.9	3.0
1999.....	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0

All races 3/, male

2003.....	2.6	9.7	2.3	0.4	0.4	2.3	3.8	3.3	2.8	3.4
2002.....	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4
2001 4/.....	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3
2000.....	2.8	9.9	2.3	0.5	0.4	2.3	4.5	3.9	3.8	3.5
1999.....	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9

All races 3/, female

2003.....	1.4	6.9	2.0	0.3	0.2	1.1	2.0	1.8	1.8	1.9
2002.....	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1
2001 4/.....	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2
2000.....	1.5	7.6	2.0	0.4	0.6	1.4	2.1	2.0	2.0	2.5
1999.....	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2003.....	1.5	6.9	1.5	0.3	0.2	1.2	2.1	2.0	1.7	2.0
2002.....	1.5	5.3	1.7	0.3	0.2	1.1	2.2	1.9	1.7	2.0
2001 4/.....	2.6	6.0	1.6	0.4	0.3	1.4	3.0	3.9	4.3	4.6
2000.....	1.6	6.3	1.5	0.4	0.4	1.3	2.3	2.1	2.0	2.1
1999.....	1.6	6.6	1.5	0.5	0.4	1.4	2.4	2.1	2.2	2.1

White, male

2003.....	1.9	8.0	1.4	0.3	0.3	1.6	2.6	2.4	2.2	2.5
2002.....	1.9	6.2	1.8	*	*	1.5	2.9	2.5	2.0	2.5
2001 4/.....	3.7	7.0	1.9	0.3	*	1.8	3.9	5.3	5.9	6.8
2000.....	2.0	7.9	1.7	0.5	0.4	1.6	3.1	2.7	2.6	2.5
1999.....	2.1	7.7	1.5	0.4	0.3	1.7	3.0	2.7	2.6	2.7

White, female

2003.....	1.1	5.6	1.6	*	*	0.8	1.5	1.5	1.2	1.4
2002.....	1.1	4.5	1.6	0.4	0.3	0.7	1.5	1.2	1.4	1.5
2001 4/.....	1.6	4.9	1.4	0.4	0.3	0.9	2.0	2.5	2.6	2.3
2000.....	1.1	4.7	1.3	0.3	0.5	1.0	1.5	1.5	1.5	1.7
1999.....	1.2	5.4	1.5	0.5	0.4	1.1	1.8	1.5	1.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2003.....	5.2	15.2	5.3	1.0	0.6	4.2	7.2	6.6	6.2	7.2
2002.....	5.3	16.5	5.8	1.0	0.6	3.7	6.4	7.3	7.0	7.4
2001 4/.....	6.3	18.6	6.1	*	0.6	4.0	7.1	9.0	8.4	9.8
2000.....	6.1	22.8	6.2	0.9	0.8	4.6	8.8	7.9	7.8	8.8
1999.....	5.7	19.2	5.4	1.3	0.6	4.5	7.0	7.7	8.1	9.2

Black, male

2003.....	7.4	16.1	6.5	*	*	5.7	9.9	9.2	7.6	9.8
2002.....	7.3	15.7	6.4	*	*	4.8	9.4	9.7	9.1	9.5
2001 4/.....	8.5	20.7	7.3	*	*	5.4	9.8	12.8	11.0	11.3
2000.....	8.3	22.3	6.2	*	*	5.6	12.4	11.1	11.0	11.1
1999.....	7.7	19.2	5.4	1.2	*	6.1	9.8	9.5	10.7	12.8

Black, female

2003.....	3.3	14.2	4.1	*	*	2.7	4.5	4.1	5.0	5.0
2002.....	3.4	17.3	5.1	*	*	2.6	3.5	5.0	5.0	5.6
2001 4/.....	4.3	16.4	4.8	*	*	2.5	4.5	5.5	6.1	8.4
2000.....	4.2	23.4	6.1	*	*	3.5	5.3	5.0	5.1	6.8
1999.....	3.9	19.1	5.4	1.3	*	2.9	4.3	6.1	5.7	6.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.4	1.7	1.9	2.7
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8

All races 3/, male

2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	2.0	1.4	0.3	*	0.4	0.9	2.0	2.5	2.7	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9

All races 3/, female

2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	1.8	1.7	0.2	*	0.2	0.9	2.1	2.0	2.6	3.4
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.7	1.4	1.8	1.9	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8

White, male

2003.....	2.3	1.9	*	*	0.3	1.3	2.9	2.7	3.3	4.4
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.9	*	*	*	0.3	1.0	2.0	2.5	2.7	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7

White, female

2003.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.3
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	1.0	1.1	1.7
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	1.8	3.2	*	*	*	*	1.1	1.4	2.1	3.2
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.9	*	*	*	*	*	1.4	1.7	2.3	3.4
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9

Black, male

2003.....	2.7	*	*	*	*	*	1.9	2.5	3.4	4.4
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.9	*	*	*	*	*	2.5	2.8	3.6	4.9
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3

Black, female

2003.....	0.9	*	*	*	*	*	*	*	*	2.2
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	1.0	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All races 3/, both sexes

2003.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2002.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	0.1	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	0.1	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1

All races 3/, male

2003.....	0.1	*	*	*	*	0.3	0.3	*	*	*
2002.....	0.1	*	*	*	*	0.3	0.4	0.2	*	0.2
2001.....	0.1	*	*	*	*	0.2	0.4	0.3	*	0.2
2000.....	0.1	*	*	*	*	0.2	0.4	0.3	*	*
1999.....	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2

All races 3/, female

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	0.1	*	*	*	*	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	*	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.1	*	0.1

White, male

2003.....	0.1	*	*	*	*	0.3	*	*	*	*
2002.....	0.1	*	*	*	*	0.3	0.3	*	*	*
2001.....	0.1	*	*	*	*	*	0.3	0.3	*	*
2000.....	0.1	*	*	*	*	*	0.3	*	*	*
1999.....	0.2	*	*	*	*	0.5	0.5	*	*	0.2

White, female

2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2003.....	1.7	1.9	0.2	*	0.1	0.7	1.7	1.7	2.3	3.1
2002.....	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2
2001.....	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9
2000.....	1.3	1.7	0.3	*	0.2	0.5	1.1	1.5	1.8	2.6
1999.....	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7

All races 3/, male

2003.....	2.2	2.4	*	*	0.2	0.9	2.4	2.4	3.0	4.1
2002.....	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3
2001.....	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8
2000.....	1.8	1.4	0.3	*	0.3	0.7	1.6	2.2	2.6	3.6
1999.....	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7

All races 3/, female

2003.....	1.2	1.3	*	*	*	0.4	1.0	1.1	1.5	2.1
2002.....	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0
2001.....	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	1.7	1.7	0.2	*	0.1	0.8	1.9	1.9	2.5	3.3
2002.....	1.6	1.8	0.4	*	0.2	0.6	1.7	1.8	2.2	3.3
2001.....	1.4	1.4	0.3	*	0.2	0.6	1.3	1.6	2.0	2.9
2000.....	1.3	1.4	0.2	*	0.2	0.5	1.2	1.6	1.8	2.6
1999.....	1.3	1.2	0.2	*	0.1	0.5	1.2	1.5	2.0	2.6

White, male

2003.....	2.2	1.9	*	*	*	1.1	2.7	2.5	3.2	4.2
2002.....	2.0	1.8	0.4	*	*	0.8	2.3	2.5	3.0	4.3
2001.....	1.9	1.7	*	*	0.3	0.9	1.9	2.2	2.7	3.6
2000.....	1.7	*	*	*	0.3	0.7	1.7	2.3	2.6	3.7
1999.....	1.7	1.4	*	*	*	0.8	1.8	2.0	2.7	3.5

White, female

2003.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.2
2002.....	1.2	1.8	0.4	*	*	0.4	1.0	1.1	1.5	2.2
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.1
2000.....	0.8	1.6	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.8	*	*	*	*	*	0.5	0.9	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	1.7	3.2	*	*	*	*	0.8	1.3	2.0	3.2
2002.....	1.9	5.5	*	*	*	*	1.1	1.8	2.4	3.9
2001.....	1.7	5.5	*	*	*	*	0.8	1.3	1.9	3.8
2000.....	1.7	*	*	*	*	*	0.9	1.4	2.0	3.2
1999.....	1.7	*	*	*	*	*	1.1	1.5	2.4	3.8

Black, male

2003.....	2.6	*	*	*	*	*	1.4	2.2	3.2	4.3
2002.....	2.7	*	*	*	*	*	1.4	2.7	3.4	6.3
2001.....	2.5	6.0	*	*	*	*	1.5	2.2	2.7	5.9
2000.....	2.6	*	*	*	*	*	1.5	2.3	3.2	4.6
1999.....	2.7	*	*	*	*	*	1.8	2.7	3.5	6.1

Black, female

2003.....	0.9	*	*	*	*	*	*	*	*	2.1
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.8
2001.....	0.8	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3

All races 3/, male

2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.1	*	*	*	*	*	*	0.3	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3

All races 3/, female

2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.2	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

White, both sexes

2003.....	1.0	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.0	*	*	*	*	*	*	0.1	0.2	0.3
2001.....	1.1	*	*	*	*	*	0.1	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.2	0.2	*	*	*	*	*	0.2	0.3

2003.....	1.0	*	*	*	*	*	*	*	
2002.....	0.9	*	*	*	*	*	*	*	0.3
2001.....	0.9	*	*	*	*	*	*	*	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.3
1999.....	1.0	1.4	*	*	*	*	*	*	0.3

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*
2001.....	1.2	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Black, both sexes

2003.....	1.1	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	0.7
2001.....	1.4	*	*	*	*	*	*	*	0.8
2000.....	1.4	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*

Black, male

2003.....	1.0	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*
2000.....	1.2	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*

Black, female

2003.....	1.3	*	*	*	*	*	*	*	*
2002.....	1.3	*	*	*	*	*	*	*	*
2001.....	1.4	*	*	*	*	*	*	*	*
2000.....	1.6	*	*	*	*	*	*	*	*
1999.....	1.3	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

All races 3/, both sexes

2003.....	7.1	*	*	*	*	0.2	0.4	0.9	2.1	5.3
2002 6/.....	7.0	*	*	*	*	0.2	0.4	0.8	2.1	5.4
2001 6/.....	7.1	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000 6/.....	7.1	*	*	*	*	0.2	0.3	1.0	2.4	6.1
1999 6/.....	7.0	*	*	*	*	0.1	0.4	0.8	2.4	6.2

All races 3/, male

2003.....	10.9	*	*	*	*	0.4	0.6	1.4	3.2	7.4
2002 6/.....	10.8	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001 6/.....	10.8	*	*	*	*	0.3	0.6	1.3	3.3	8.5
2000 6/.....	11.1	*	*	*	*	0.3	0.5	1.6	3.7	8.9
1999 6/.....	10.9	*	*	*	*	0.2	0.6	1.2	3.3	8.7

All races 3/, female

2003.....	3.4	*	*	*	*	*	*	0.3	0.9	3.2
2002 6/.....	3.4	*	*	*	*	*	*	0.3	1.1	3.3
2001 6/.....	3.4	*	*	*	*	*	*	0.5	1.3	3.5
2000 6/.....	3.3	*	*	*	*	*	*	0.3	1.2	3.3
1999 6/.....	3.2	*	*	*	*	*	*	0.3	1.5	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

White, both sexes

2003.....	7.4	*	*	*	*	0.2	0.4	0.9	2.1	5.4
2002 6/.....	7.2	*	*	*	*	0.2	0.3	0.8	2.1	5.5
2001 6/.....	7.2	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000 6/.....	7.2	*	*	*	*	0.2	0.3	1.0	2.3	6.0
1999 6/.....	7.0	*	*	*	*	*	0.4	0.8	2.4	6.0

White, male

2003.....	11.3	*	*	*	*	0.4	0.6	1.5	3.2	7.6
2002 6/.....	11.1	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001 6/.....	11.0	*	*	*	*	0.3	0.5	1.3	3.3	8.4
2000 6/.....	11.2	*	*	*	*	0.3	0.5	1.7	3.6	8.9
1999 6/.....	10.9	*	*	*	*	*	0.7	1.3	3.2	8.5

White, female

2003.....	3.5	*	*	*	*	*	*	0.3	1.0	3.2
2002 6/.....	3.4	*	*	*	*	*	*	*	1.0	3.3
2001 6/.....	3.4	*	*	*	*	*	*	0.4	1.2	3.6
2000 6/.....	3.2	*	*	*	*	*	*	*	1.1	3.2
1999 6/.....	3.1	*	*	*	*	*	*	*	1.4	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

Black, both sexes

2003.....	6.3	*	*	*	*	*	*	0.9	1.3	4.7
2002 6/.....	6.4	*	*	*	*	*	*	*	1.9	4.4
2001 6/.....	7.3	*	*	*	*	*	*	0.8	1.9	5.2
2000 6/.....	7.7	*	*	*	*	*	*	0.9	2.6	5.6
1999 6/.....	7.8	*	*	*	*	*	*	*	2.2	7.1

Black, male

2003.....	10.0	*	*	*	*	*	*	*	2.2	6.5
2002 6/.....	10.0	*	*	*	*	*	*	*	2.9	6.2
2001 6/.....	11.6	*	*	*	*	*	*	*	2.5	8.4
2000 6/.....	11.9	*	*	*	*	*	*	1.6	3.6	8.3
1999 6/.....	12.2	*	*	*	*	*	*	*	2.9	10.2

Black, female

2003.....	2.9	*	*	*	*	*	*	*	*	3.2
2002 6/.....	3.2	*	*	*	*	*	*	*	*	2.9
2001 6/.....	3.5	*	*	*	*	*	*	*	*	2.4
2000 6/.....	3.9	*	*	*	*	*	*	*	1.8	3.3
1999 6/.....	3.9	*	*	*	*	*	*	*	1.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ year 2/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

All races 3/, both sexes

2003.....	9.9	0.6	0.2	*	0.2	3.3	9.4	10.9	13.3	18.3
2002 6/.....	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5
2001 6/.....	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5
2000 6/.....	7.2	*	*	*	0.2	2.3	6.0	8.1	10.6	14.7
1999 6/.....	6.9	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2

All races 3/, male

2003.....	12.9	*	0.3	*	*	4.5	13.9	15.4	17.7	23.8
2002 6/.....	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1
2001 6/.....	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7
2000 6/.....	9.8	*	*	*	*	3.1	9.0	11.9	15.0	20.2
1999 6/.....	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6

All races 3/, female

2003.....	7.0	*	*	*	*	1.9	4.7	6.2	8.8	12.8
2002 6/.....	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9
2001 6/.....	5.1	*	*	*	*	1.6	3.2	4.4	6.7	10.3
2000 6/.....	4.7	*	*	*	*	1.4	2.9	4.4	6.3	9.2
1999 6/.....	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

White, both sexes

2003.....	10.4	*	0.2	*	0.1	3.9	11.2	12.4	14.7	19.4
2002 6/.....	9.4	*	*	*	0.2	3.6	9.2	10.9	13.5	18.5
2001 6/.....	7.8	*	*	*	0.2	3.1	7.5	9.1	11.4	16.1
2000 6/.....	7.2	*	*	*	0.2	2.6	6.8	8.9	11.2	15.0
1999 6/.....	6.9	*	*	*	0.1	2.1	5.9	7.9	10.7	14.4

White, male

2003.....	13.5	*	*	*	*	5.5	16.4	17.6	19.5	25.1
2002 6/.....	12.2	*	*	*	*	5.0	13.2	15.5	18.5	24.2
2001 6/.....	10.4	*	*	*	*	4.4	10.9	13.1	15.6	21.6
2000 6/.....	9.8	*	*	*	*	3.7	10.2	12.8	15.8	20.8
1999 6/.....	9.3	*	*	*	*	2.8	8.5	11.3	15.2	19.9

White, female

2003.....	7.4	*	*	*	*	2.2	5.5	6.8	9.7	13.6
2002 6/.....	6.7	*	*	*	*	2.1	4.9	6.1	8.3	12.6
2001 6/.....	5.4	*	*	*	*	1.6	3.8	5.0	7.0	10.6
2000 6/.....	4.8	*	*	*	*	1.5	3.2	4.9	6.5	9.2
1999 6/.....	4.5	*	*	*	*	1.4	3.0	4.3	6.1	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Black, both sexes

2003.....	9.2	*	*	*	*	0.7	2.9	5.8	10.4	17.9
2002 6/.....	9.2	*	*	*	*	1.0	3.2	6.6	10.2	17.3
2001 6/.....	8.5	*	*	*	*	0.9	3.0	5.8	9.5	16.7
2000 6/.....	8.6	*	*	*	*	1.0	3.2	6.4	10.7	16.9
1999 6/.....	8.6	*	*	*	*	0.7	3.7	6.8	11.2	16.5

Black, male

2003.....	12.7	*	*	*	*	*	4.4	7.7	13.6	24.3
2002 6/.....	12.8	*	*	*	*	1.4	4.6	9.9	13.9	24.2
2001 6/.....	12.2	*	*	*	*	*	5.1	9.3	12.3	22.4
2000 6/.....	12.5	*	*	*	*	*	4.6	10.1	14.9	23.2
1999 6/.....	12.7	*	*	*	*	*	5.6	11.0	15.7	23.9

Black, female

2003.....	6.1	*	*	*	*	*	1.5	4.0	7.6	12.2
2002 6/.....	5.8	*	*	*	*	*	1.9	3.6	6.8	11.2
2001 6/.....	5.1	*	*	*	*	*	*	2.6	7.0	11.6
2000 6/.....	5.1	*	*	*	*	*	1.9	3.0	7.0	11.2
1999 6/.....	4.8	*	*	*	*	*	1.9	3.1	7.3	10.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

All races 3/, both sexes

2003.....	10.4	*	0.3	0.3	1.2	12.1	21.1	17.4	13.8	11.6
2002.....	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4
2001.....	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2
2000.....	10.4	*	0.3	0.4	1.5	13.1	21.5	17.7	13.4	12.2
1999.....	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3

All races 3/, male

2003.....	18.3	*	0.3	0.4	2.0	21.2	37.1	30.7	23.7	19.3
2002.....	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6
2001.....	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0
2000.....	18.3	*	0.4	0.5	2.5	22.7	38.0	31.4	22.7	19.9
1999.....	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0

All races 3/, female

2003.....	2.7	*	0.3	0.2	0.4	2.4	4.2	3.6	3.7	3.8
2002.....	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2
2001.....	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6
2000.....	2.9	*	*	0.2	0.5	2.9	4.3	4.1	4.2	4.6
1999.....	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

White, both sexes

2003.....	9.2	*	0.2	0.2	1.0	8.4	13.9	11.7	10.8	10.4
2002.....	9.3	*	0.3	0.3	1.1	8.7	13.9	12.3	10.2	10.9
2001.....	9.4	*	0.4	0.4	1.1	8.9	14.0	12.9	11.1	11.2
2000.....	9.3	*	0.2	0.3	1.4	9.3	14.1	12.5	10.9	11.0
1999.....	9.2	*	0.3	0.3	1.6	10.2	14.0	12.1	11.1	11.3

White, male

2003.....	16.0	*	*	*	1.6	14.4	23.9	19.8	17.9	16.8
2002.....	16.1	*	0.4	0.3	1.5	14.9	23.9	20.5	16.7	17.7
2001.....	16.2	*	0.4	0.5	1.7	15.0	24.1	21.8	18.2	17.9
2000.....	16.0	*	*	0.4	2.4	15.8	24.4	21.3	18.0	17.5
1999.....	15.9	*	*	0.3	2.3	17.3	23.8	20.3	18.1	18.1

White, female

2003.....	2.6	*	*	*	0.3	2.0	3.1	3.0	3.3	3.9
2002.....	2.7	*	*	0.3	0.6	2.1	3.1	3.7	3.4	4.0
2001.....	2.7	*	*	0.3	0.5	2.3	3.1	3.4	3.7	4.3
2000.....	2.8	*	*	*	0.5	2.4	3.3	3.5	3.8	4.5
1999.....	2.8	*	*	*	0.7	2.6	3.5	3.4	3.8	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
Black, both sexes										
2003.....	20.1	*	0.9	0.8	2.5	32.2	63.8	55.5	35.8	21.5
2002.....	20.2	*	*	*	2.4	31.4	64.7	54.8	35.1	24.4
2001.....	19.3	*	0.9	*	1.9	32.9	64.4	50.4	31.5	22.6
2000.....	20.0	*	*	*	2.3	34.5	64.3	50.1	32.1	22.1
1999.....	19.4	*	1.1	0.9	2.3	38.0	63.5	46.5	29.0	20.9
Black, male										
2003.....	37.8	*	*	*	3.7	59.3	117.4	107.6	67.4	40.3
2002.....	37.8	*	*	*	3.5	56.0	120.6	106.3	65.9	44.8
2001.....	36.4	*	*	*	3.0	60.5	123.4	96.0	60.1	39.5
2000.....	37.5	*	*	*	3.8	62.2	119.8	96.2	59.7	40.6
1999.....	36.0	*	*	1.3	3.4	67.1	119.9	88.6	54.3	37.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
Black, female										
2003.....	3.9	*	*	*	1.3	4.2	10.5	7.4	7.2	4.7
2002.....	4.2	*	*	*	1.3	6.0	10.2	7.7	7.4	6.3
2001.....	3.8	*	*	*	*	4.5	7.8	8.9	5.8	7.5
2000.....	4.1	*	*	*	*	5.8	9.8	7.9	7.6	5.6
1999.....	4.4	*	*	*	*	8.2	9.9	8.4	6.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All causes

All races 3/, both sexes

2003.....	241.3	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	237.3	356.0	518.6	801.8	1,257.9	1,928.2	2,968.1	4,556.6	7,399.6	15,321.5
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6

All races 3/, male

2003.....	304.1	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	303.3	461.7	658.3	1,007.5	1,565.5	2,399.3	3,705.4	5,591.2	8,956.9	16,605.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2

All races 3/, female

2003.....	179.2	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	172.2	254.0	386.3	611.8	982.0	1,527.5	2,381.8	3,812.6	6,444.8	14,768.6
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All causes

White, both sexes

2003.....	222.6	329.4	474.5	710.6	1,127.8	1,739.2	2,728.6	4,329.5	6,992.1	14,792.3
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	213.3	319.6	473.5	749.8	1,197.7	1,871.7	2,906.9	4,497.2	7,379.4	15,532.5
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0

White, male

2003.....	284.0	423.5	606.9	884.2	1,390.6	2,146.2	3,359.5	5,349.4	8,522.6	16,037.9
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	275.9	418.9	598.5	936.0	1,484.5	2,328.7	3,632.1	5,521.0	8,956.4	16,897.7
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9

White, female

2003.....	160.7	235.8	344.9	543.9	884.5	1,379.6	2,212.3	3,592.1	6,048.6	14,240.6
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	150.4	221.6	352.3	574.0	934.5	1,475.3	2,321.8	3,754.5	6,408.5	14,948.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All causes

Black, both sexes

2003.....	410.4	632.9	940.3	1,295.1	1,889.9	2,595.8	3,754.9	5,421.2	7,745.3	13,976.7
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.1	677.5	971.3	1,366.1	1,950.9	2,662.8	3,984.2	5,803.9	8,515.6	14,752.1
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9

Black, male

2003.....	507.1	800.8	1,227.5	1,688.7	2,454.5	3,334.8	4,843.9	6,978.2	9,827.8	14,903.4
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	548.0	877.0	1,300.9	1,854.6	2,573.7	3,365.9	4,960.0	7,139.2	10,247.9	15,494.6
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5

Black, female

2003.....	325.1	486.7	697.2	971.3	1,449.4	2,047.3	3,020.5	4,495.8	6,654.1	13,616.7
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.2	508.2	702.4	986.1	1,487.3	2,140.4	3,292.0	4,943.7	7,562.6	14,441.9
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes

2003.....	*	0.1	0.2	0.4	0.9	1.7	3.3	7.0	12.3	24.8
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.7	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.8
2000.....	*	*	*	0.2	0.3	0.9	1.6	2.7	5.3	11.6
1999.....	*	*	*	*	0.3	0.7	1.2	2.3	4.5	10.3

All races 3/, male

2003.....	*	*	0.2	0.5	0.8	1.5	3.1	6.9	12.9	21.4
2002.....	*	*	*	0.4	0.8	1.5	3.1	5.1	10.1	20.4
2001.....	*	*	*	*	0.4	0.7	2.0	3.5	6.2	12.1
2000.....	*	*	*	0.3	*	1.0	1.6	2.8	5.7	10.8
1999.....	*	*	*	*	*	0.6	1.2	2.7	4.9	9.5

All races 3/, female

2003.....	*	*	0.2	0.4	0.9	1.8	3.5	7.1	12.0	26.3
2002.....	*	*	*	0.4	0.6	1.3	2.3	5.0	9.5	21.3
2001.....	*	*	0.2	*	0.5	0.8	1.8	3.0	6.5	13.1
2000.....	*	*	*	*	*	0.8	1.6	2.6	5.0	12.0
1999.....	*	*	*	*	*	0.8	1.3	2.0	4.3	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2003.....	*	*	0.2	0.5	0.9	1.8	3.5	7.2	13.2	26.1
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.7	5.3	10.3	22.0
2001.....	*	*	0.2	0.2	0.4	0.7	2.0	3.3	6.6	13.5
2000.....	*	*	*	0.3	0.3	0.9	1.6	2.9	5.6	12.1
1999.....	*	*	*	*	0.3	0.7	1.3	2.3	4.7	10.9

White, male

2003.....	*	*	*	0.4	0.8	1.6	3.2	7.2	13.7	22.7
2002.....	*	*	*	0.4	0.7	1.6	3.0	5.3	10.7	21.4
2001.....	*	*	*	*	*	0.7	2.0	3.6	6.2	12.7
2000.....	*	*	*	*	*	1.1	1.6	3.0	6.0	11.4
1999.....	*	*	*	*	*	0.7	1.3	2.8	5.2	9.9

White, female

2003.....	*	*	*	0.5	0.9	1.9	3.8	7.2	12.8	27.6
2002.....	*	*	*	0.4	0.6	1.4	2.4	5.3	10.1	22.3
2001.....	*	*	0.3	*	0.5	0.8	1.9	3.2	6.8	13.9
2000.....	*	*	*	*	*	0.8	1.6	2.8	5.4	12.4
1999.....	*	*	*	*	*	0.8	1.3	2.1	4.4	11.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Tuberculosis (A16-A19)

All races 3/, both sexes

2003.....	0.1	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2

All races 3/, male

2003.....	*	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.1	1.1	2.2	3.8	4.3
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1

All races 3/, female

2003.....	*	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.5	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Tuberculosis (A16-A19)

White, both sexes

2003.....	*	0.1	0.1	0.3	0.2	0.3	0.7	1.0	1.7	1.8
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2

White, male

2003.....	*	*	*	0.4	*	*	1.0	1.3	2.2	2.4
2002.....	*	*	*	*	0.5	0.8	1.0	1.3	2.5	3.2
2001.....	*	*	*	0.5	0.5	0.6	0.8	1.4	1.8	3.1
2000.....	*	*	0.3	*	*	0.7	0.8	1.4	2.2	3.6
1999.....	*	0.3	0.4	0.5	0.6	0.9	1.1	1.5	3.0	5.4

White, female

2003.....	*	*	*	*	*	*	*	0.9	1.3	1.6
2002.....	*	*	*	*	*	*	*	1.1	1.3	1.6
2001.....	*	*	*	*	*	*	0.5	0.9	1.2	1.8
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.1	1.4	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Respiratory tuberculosis (A16)

All races 3/, both sexes

2003.....	0.1	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.2	0.2	0.2	0.3	0.6	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7

All races 3/, male

2003.....	*	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.1	3.9
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4

All races 3/, female

2003.....	*	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.7	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Respiratory tuberculosis (A16)

White, both sexes

2003.....	*	*	*	0.2	*	*	0.5	0.9	1.4	1.4
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0

White, male

2003.....	*	*	*	0.3	*	*	0.9	1.1	1.8	1.8
2002.....	*	*	*	*	*	0.7	0.8	0.9	2.0	2.9
2001.....	*	*	*	0.4	0.5	*	0.7	1.2	1.4	2.6
2000.....	*	*	0.3	*	*	0.5	0.7	1.3	1.8	3.3
1999.....	*	*	*	0.4	0.5	0.7	1.0	1.3	2.6	5.2

White, female

2003.....	*	*	*	*	*	*	*	0.8	1.1	1.2
2002.....	*	*	*	*	*	*	*	0.9	1.1	1.3
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.4
2000.....	*	*	*	*	*	*	*	0.8	1.3	1.3
1999.....	*	*	*	*	*	*	*	0.9	1.1	2.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Septicemia (A40-A41)

All races 3/, both sexes

2003.....	2.6	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.4	3.9	6.2	9.5	15.4	24.4	38.9	64.0	105.9	212.9
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7

All races 3/, male

2003.....	2.5	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.5	7.1	10.3	17.0	27.3	43.5	72.5	119.8	211.6
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5

All races 3/, female

2003.....	2.7	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.9	22.0	35.2	57.9	97.4	213.4
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Septicemia (A40-A41)

White, both sexes

2003.....	2.1	3.4	5.2	8.8	14.3	21.7	36.9	63.7	102.5	191.1
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.8	21.0	35.0	58.0	98.0	200.7
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9

White, male

2003.....	2.0	3.8	5.7	9.3	15.3	24.6	40.8	71.3	116.7	203.3
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.1	3.5	5.6	8.5	14.2	23.6	39.8	65.5	112.3	200.1
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4

White, female

2003.....	2.2	3.0	4.7	8.2	13.4	19.2	33.7	58.2	93.7	185.8
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.4	18.9	31.1	52.5	89.3	201.0
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Septicemia (A40-A41)

Black, both sexes

2003.....	6.3	10.5	17.5	25.3	41.5	61.0	88.0	133.4	193.4	372.0
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.2	17.5	24.7	40.6	57.4	85.8	147.0	225.7	393.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3

Black, male

2003.....	6.4	11.8	19.7	31.8	46.1	74.9	100.2	160.2	235.5	404.5
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.8	22.5	29.1	47.2	66.0	92.4	176.2	246.6	394.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5

Black, female

2003.....	6.3	9.4	15.6	20.0	37.9	50.8	79.8	117.5	171.4	359.3
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.8	13.4	21.3	35.8	50.9	81.1	128.1	214.2	393.1
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

All races 3/, both sexes

2003.....	2.1	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.9	3.7	3.9	4.3	5.0	5.4	5.3	3.9
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7

All races 3/, male

2003.....	2.9	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.1	7.6	5.3	4.7	4.7	5.0	6.4	6.1	5.4
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9

All races 3/, female

2003.....	1.2	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.7	4.8	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

White, both sexes

2003.....	2.1	4.9	6.0	3.8	2.7	2.9	3.8	4.0	3.8	2.3
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	5.0	4.3	3.1	3.4	3.9	4.6	5.3	5.0	3.8
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5

White, male

2003.....	2.9	7.2	9.2	5.7	3.4	3.3	4.3	4.2	4.8	2.8
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.7	6.7	4.4	4.2	4.1	4.7	6.3	5.8	5.4
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5

White, female

2003.....	1.2	2.6	2.9	2.0	2.0	2.6	3.5	3.8	3.2	2.1
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.8	4.6	4.5	4.5	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

Black, both sexes

2003.....	2.7	5.2	10.1	9.4	6.3	5.1	5.7	6.5	5.3	*
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.4	9.1	7.0	4.6	5.0	6.2	5.7	7.4	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4

Black, male

2003.....	3.6	7.6	16.2	14.9	7.5	5.4	6.9	*	*	*
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.3	14.8	10.8	5.4	6.5	7.0	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*

Black, female

2003.....	1.9	3.2	5.0	4.8	5.3	4.9	4.9	5.7	*	*
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.1	4.4	4.0	4.0	3.9	5.7	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2003.....	12.8	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.4	12.8	9.2	6.0	4.1	2.7	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All races 3/, male

2003.....	19.1	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.2	20.5	15.3	10.2	7.1	4.6	2.9	1.7	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All races 3/, female

2003.....	6.6	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.6	5.4	3.4	2.1	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2003.....	7.4	6.7	4.9	3.5	2.1	1.3	1.0	0.6	*	*
2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.4	6.9	4.8	3.4	2.2	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*

White, male

2003.....	12.1	11.3	8.4	6.0	3.6	2.2	2.0	1.3	*	*
2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.1	11.8	8.5	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

White, female

2003.....	2.6	2.1	1.3	1.1	0.6	0.5	*	*	*	*
2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2003.....	52.3	53.4	47.2	33.9	23.8	17.2	11.0	4.9	*	*
2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	59.6	48.2	29.7	22.3	14.6	11.5	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*

Black, male

2003.....	74.0	79.5	75.7	57.6	41.3	30.0	20.0	*	*	*
2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	83.9	95.3	82.0	53.3	41.3	25.2	21.3	11.4	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*

Black, female

2003.....	33.1	30.7	23.1	14.3	10.2	7.7	4.9	*	*	*
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.4	20.7	11.4	8.1	6.7	4.5	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2003.....	1.3	2.4	3.5	3.7	4.5	5.9	8.9	12.3	14.8	22.5
2002.....	1.3	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001.....	0.9	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000.....	0.8	1.4	1.8	2.4	3.6	4.8	7.1	10.0	13.0	20.9
1999.....	1.4	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0

All races 3/, male

2003.....	1.6	3.3	4.7	4.9	5.4	7.0	10.3	13.9	18.8	26.5
2002.....	1.7	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001.....	1.2	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000.....	1.0	1.7	2.3	2.8	4.0	5.8	8.9	11.8	16.0	26.2
1999.....	1.7	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9

All races 3/, female

2003.....	0.9	1.6	2.2	2.6	3.7	5.0	7.8	11.1	12.4	20.8
2002.....	0.8	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001.....	0.6	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000.....	0.6	1.1	1.3	2.1	3.2	3.9	5.7	8.6	11.1	18.7
1999.....	1.1	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2003.....	1.2	2.3	3.3	3.5	4.2	5.9	8.9	11.8	14.7	23.0
2002.....	1.1	1.8	2.5	2.6	3.6	5.4	7.9	11.3	13.8	22.0
2001.....	0.8	1.4	1.7	2.2	3.6	5.4	6.7	9.3	12.1	20.1
2000.....	0.7	1.3	1.6	2.3	3.2	4.7	6.9	9.8	12.8	20.8
1999.....	1.1	1.6	2.2	2.9	3.9	6.4	8.5	12.1	16.1	26.5

White, male

2003.....	1.6	3.2	4.5	4.6	5.0	6.8	10.0	13.5	18.6	27.2
2002.....	1.5	2.5	3.5	3.2	4.3	6.6	9.4	13.6	17.6	27.2
2001.....	1.0	1.9	2.2	2.8	4.3	6.0	7.2	11.3	15.9	23.8
2000.....	0.9	1.5	2.0	2.7	3.7	5.6	8.7	11.4	15.9	25.6
1999.....	1.4	2.0	2.6	3.3	4.5	7.5	10.2	14.8	18.6	30.6

White, female

2003.....	0.8	1.5	2.1	2.4	3.5	5.0	8.0	10.6	12.3	21.2
2002.....	0.7	1.1	1.5	2.0	2.9	4.3	6.7	9.7	11.5	19.8
2001.....	0.6	0.8	1.3	1.6	3.1	5.0	6.2	7.8	9.9	18.5
2000.....	0.6	1.0	1.1	1.9	2.8	3.9	5.5	8.7	11.0	18.7
1999.....	0.7	1.2	1.9	2.6	3.4	5.6	7.2	10.2	14.6	24.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2003.....	2.1	3.6	5.3	6.0	6.2	5.9	8.3	16.2	16.2	19.5
2002.....	2.5	3.2	4.8	6.0	5.4	5.3	9.7	15.3	11.5	21.7
2001.....	1.8	2.5	3.4	4.4	6.0	7.1	9.8	9.7	13.3	20.6
2000.....	1.7	2.7	3.6	4.1	6.0	5.8	9.8	11.6	16.8	24.1
1999.....	3.6	5.1	5.9	6.3	8.2	10.7	13.1	18.7	23.4	34.1

Black, male

2003.....	2.6	4.6	7.1	7.7	8.3	8.9	13.1	16.4	22.1	22.9
2002.....	3.2	4.5	6.7	9.2	7.3	6.5	12.0	18.4	*	28.9
2001.....	2.7	3.6	4.3	5.5	7.3	9.7	10.4	10.2	16.1	*
2000.....	2.1	3.3	5.0	4.6	7.6	8.2	11.8	16.4	19.9	29.8
1999.....	3.6	5.8	6.5	5.8	10.4	12.5	15.9	21.8	41.9	45.7

Black, female

2003.....	1.6	2.6	3.8	4.6	4.5	3.7	5.1	16.0	13.2	18.2
2002.....	1.9	2.0	3.1	3.4	3.9	4.4	8.1	13.4	9.9	18.9
2001.....	*	1.6	2.7	3.6	5.0	5.3	9.4	9.4	11.9	20.2
2000.....	*	2.1	2.4	3.6	4.8	4.1	8.4	8.5	15.1	21.8
1999.....	3.5	4.5	5.3	6.8	6.5	9.4	11.3	16.9	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2003.....	46.1	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.4	94.5	169.4	292.4	469.7	698.6	964.2	1,223.2	1,518.6	1,795.6
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8

All races 3/, male

2003.....	41.7	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.0	94.6	178.3	327.0	546.0	847.9	1,199.9	1,557.6	2,019.2	2,571.9
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2

All races 3/, female

2003.....	50.5	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	52.8	94.5	160.9	260.4	401.4	571.6	776.7	982.8	1,211.6	1,461.3
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

White, both sexes

2003.....	44.0	84.8	151.3	261.2	434.5	644.8	908.5	1,195.1	1,450.7	1,701.4
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.2	88.7	161.6	283.4	462.0	695.1	957.3	1,214.3	1,512.4	1,795.6
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8

White, male

2003.....	39.8	85.1	159.7	285.4	494.7	763.2	1,108.1	1,518.9	1,930.1	2,412.1
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	40.9	88.3	166.8	310.7	528.9	835.4	1,179.3	1,531.7	1,996.5	2,569.2
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3

White, female

2003.....	48.3	84.6	143.1	237.8	378.9	540.3	745.2	961.0	1,155.2	1,386.5
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	49.5	89.1	156.5	257.6	400.7	573.5	778.2	984.1	1,214.4	1,464.7
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

Black, both sexes

2003.....	66.5	138.3	243.2	382.6	589.1	802.7	1,080.1	1,385.5	1,640.8	1,853.1
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	148.0	260.6	413.6	611.5	828.3	1,168.8	1,487.1	1,774.5	2,004.0
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4

Black, male

2003.....	61.2	145.6	283.0	464.8	747.3	1,045.9	1,455.8	1,975.5	2,417.1	2,825.5
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.1	156.1	311.3	535.4	804.1	1,089.4	1,577.3	2,086.9	2,642.5	3,012.7
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7

Black, female

2003.....	71.1	132.0	209.5	314.9	465.7	622.1	826.8	1,034.8	1,234.0	1,475.3
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	141.1	219.2	318.9	468.1	634.4	879.0	1,100.8	1,297.1	1,582.6
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2003.....	0.9	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.1	3.8	5.8	8.2	10.0	11.6	12.6	14.1	19.2
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3

All races 3/, male

2003.....	1.4	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.2	6.1	9.7	12.9	15.4	17.0	18.4	19.9	25.7
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1

All races 3/, female

2003.....	0.5	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.4	8.4	10.6	16.4
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, both sexes

2003.....	0.8	1.8	3.2	5.2	7.3	9.3	10.4	12.6	13.4	19.0
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.8	3.2	5.1	7.4	9.5	11.2	12.5	14.1	19.5
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8

White, male

2003.....	1.3	2.7	5.5	8.5	11.3	14.1	15.2	18.2	19.2	27.2
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	5.0	8.4	11.4	14.3	16.1	17.7	19.6	26.3
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3

White, female

2003.....	0.3	0.9	0.9	2.1	3.7	5.1	6.4	8.5	9.8	15.4
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.7	5.4	7.3	8.7	10.8	16.6
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, both sexes

2003.....	1.7	3.9	6.5	8.8	14.4	15.5	15.3	14.8	16.0	15.5
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.6	12.3	15.3	14.3	16.0	14.6	16.5	15.4
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3

Black, male

2003.....	1.9	7.0	11.4	16.3	27.8	27.6	30.2	26.5	30.3	25.0
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.5	16.9	22.9	28.6	26.4	27.7	27.8	28.2	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6

Black, female

2003.....	1.5	*	2.3	2.7	3.9	6.6	5.3	7.9	8.5	11.7
2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.1	5.5	5.4	7.7	6.2	10.0	14.7
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes

2003.....	0.9	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	5.0	8.9	13.4	18.2	22.4	24.5	27.5	25.6
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8

All races 3/, male

2003.....	1.5	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.7	4.2	8.9	15.5	23.4	31.1	38.3	43.1	48.1	45.3
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0

All races 3/, female

2003.....	0.3	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.8	11.1	14.8	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of esophagus (C15)

White, both sexes

2003.....	0.9	2.3	4.3	8.5	12.3	17.0	22.5	25.8	27.8	25.8
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.6	8.1	12.6	17.7	22.1	24.0	27.4	26.3
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2

White, male

2003.....	1.6	3.9	7.6	15.0	22.0	29.5	39.0	46.5	52.3	46.7
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.7	4.1	8.1	14.2	22.2	30.6	37.9	42.4	48.9	47.1
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0

White, female

2003.....	0.3	0.6	1.0	2.3	3.4	6.0	9.0	10.9	12.7	16.5
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.2	2.4	3.7	6.5	9.3	10.7	14.1	17.4
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of esophagus (C15)

Black, both sexes

2003.....	1.3	4.1	8.4	14.1	20.7	26.1	24.9	26.9	29.4	21.3
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	10.2	17.2	23.6	25.9	29.4	36.2	31.2	22.3
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9

Black, male

2003.....	2.1	7.1	14.0	23.3	36.2	46.5	41.4	47.5	43.6	38.6
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.3	6.5	18.5	31.8	40.3	42.9	47.5	63.3	44.8	35.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6

Black, female

2003.....	*	1.6	3.6	6.5	8.6	11.0	13.7	14.7	22.0	14.6
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.1	13.3	16.5	18.8	23.7	16.9
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2003.....	1.3	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.4	3.6	5.7	9.9	14.0	19.9	27.1	37.2	52.0
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1

All races 3/, male

2003.....	1.5	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.6	3.0	4.9	7.9	14.3	19.9	28.6	38.7	50.6	71.8
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3

All races 3/, female

2003.....	1.1	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.8	2.4	3.6	5.9	9.0	13.0	18.7	29.0	43.5
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of stomach (C16)

White, both sexes

2003.....	1.1	1.8	3.1	4.6	7.6	11.0	16.3	21.4	28.8	38.7
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.2	4.9	8.6	12.1	17.5	23.6	32.9	46.8
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8

White, male

2003.....	1.2	2.3	4.1	6.6	10.6	15.5	23.5	30.3	40.3	53.5
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.3	6.8	12.7	17.4	25.4	34.0	45.4	65.7
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4

White, female

2003.....	0.9	1.4	2.2	2.7	4.8	7.0	10.5	15.0	21.7	32.1
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.5	11.1	16.0	25.2	38.7
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of stomach (C16)

Black, both sexes

2003.....	2.4	4.3	7.4	9.6	16.7	22.3	33.1	48.8	62.4	85.8
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.2	11.2	17.5	25.8	37.8	57.1	78.4	101.9
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4

Black, male

2003.....	3.3	6.2	10.9	14.9	26.0	36.4	54.2	74.9	87.1	106.3
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.8	10.2	17.2	27.1	36.9	54.5	80.1	103.8	125.6
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6

Black, female

2003.....	1.5	2.7	4.5	5.3	9.5	11.9	18.8	33.4	49.5	77.8
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.3	17.6	26.0	42.2	64.4	92.0
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2003.....	3.8	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	3.9	8.3	15.0	25.5	42.6	62.7	93.7	127.0	176.3	266.8
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7

All races 3/, male

2003.....	4.2	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.3	9.1	17.5	31.1	52.0	78.6	116.7	156.5	209.7	299.5
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0

All races 3/, female

2003.....	3.4	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.6	20.4	34.2	49.2	75.4	105.7	155.8	252.8
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2003.....	3.4	7.3	13.0	21.5	36.9	58.0	80.7	115.1	158.2	238.5
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	14.0	23.9	41.0	61.2	91.9	125.1	173.2	266.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9

White, male

2003.....	3.8	8.3	15.2	26.2	45.0	73.6	100.8	140.7	196.0	272.9
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.4	16.4	28.9	49.8	76.7	114.7	154.9	205.5	298.8
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5

White, female

2003.....	2.9	6.3	10.8	17.0	29.5	44.1	64.2	96.6	134.8	223.2
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.7	11.6	19.2	32.9	47.8	73.5	103.5	153.4	252.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Black, both sexes

2003.....	6.6	13.2	25.5	39.4	59.7	81.2	124.6	154.7	197.7	281.5
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.5	15.5	25.7	40.5	63.5	83.6	125.0	168.0	236.0	310.8
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1

Black, male

2003.....	7.1	15.2	30.3	46.1	76.4	102.7	156.7	194.9	242.2	317.0
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.9	30.9	53.6	79.9	104.7	152.6	201.2	293.9	348.0
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4

Black, female

2003.....	6.1	11.4	21.4	33.8	46.8	65.2	103.0	130.8	174.5	267.8
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.4	30.2	51.2	68.0	105.3	146.6	204.1	295.3
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2003.....	1.2	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.5	7.8	11.4	16.2	21.9	27.1	31.6	32.3
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6

All races 3/, male

2003.....	1.8	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.1	8.7	11.9	16.8	24.0	31.2	38.6	45.9	42.4
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1

All races 3/, female

2003.....	0.7	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.5	2.4	4.0	6.6	9.6	14.5	18.9	22.7	27.9
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

White, both sexes

2003.....	1.0	3.1	6.3	7.5	10.3	15.2	21.7	28.3	31.0	32.1
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.0	3.1	4.5	6.6	10.1	14.8	20.5	25.8	30.9	31.6
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9

White, male

2003.....	1.4	4.9	10.2	11.4	15.2	21.4	31.4	41.8	44.8	45.6
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.9	6.9	9.7	14.6	21.8	29.2	37.5	45.2	41.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6

White, female

2003.....	0.6	1.3	2.5	3.8	5.7	9.7	13.9	18.6	22.5	26.2
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.1	3.6	5.9	8.6	13.4	17.3	22.1	27.6
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2003.....	1.9	6.2	14.2	14.7	18.4	18.9	27.9	30.3	32.7	41.7
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.1	11.0	12.9	16.0	20.1	26.7	28.9	29.8	31.7
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4

Black, male

2003.....	3.0	10.3	25.5	24.0	28.5	26.8	42.4	43.8	45.0	63.6
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.9	18.7	23.0	26.4	32.6	39.5	35.1	44.8	44.7
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5

Black, female

2003.....	*	2.7	4.5	7.0	10.5	13.0	18.2	22.2	26.3	33.2
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	4.9	8.2	10.7	17.6	24.9	21.5	26.2
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes

2003.....	1.7	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.9	15.2	25.1	37.6	52.1	70.5	86.4	95.1
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7

All races 3/, male

2003.....	2.0	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.4	18.9	30.6	45.4	57.4	78.9	93.5	97.6
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5

All races 3/, female

2003.....	1.4	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.5	11.9	20.2	31.0	47.9	64.4	82.1	94.1
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of pancreas (C25)

White, both sexes

2003.....	1.6	3.9	8.1	14.3	23.8	36.0	52.2	67.2	85.2	94.4
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.5	14.6	24.6	36.8	51.1	69.2	85.1	93.6
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6

White, male

2003.....	1.8	4.8	10.0	17.7	28.9	42.7	60.2	76.1	99.3	96.0
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.9	18.0	30.0	45.1	56.5	78.5	92.6	97.6
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4

White, female

2003.....	1.4	2.9	6.3	11.0	19.1	30.0	45.8	60.7	76.6	93.8
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.2	11.4	19.6	29.5	46.7	62.4	80.4	91.9
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of pancreas (C25)

Black, both sexes

2003.....	2.7	6.4	14.5	20.7	36.1	47.1	68.0	83.6	107.4	112.9
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.9	13.6	23.1	33.7	48.3	69.2	94.2	108.7	120.1
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6

Black, male

2003.....	3.5	9.0	18.6	26.2	42.8	50.0	80.8	89.0	112.2	104.3
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.5	18.6	30.6	42.2	53.3	75.2	97.9	114.6	101.1
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8

Black, female

2003.....	2.1	4.1	11.0	16.2	30.8	45.0	59.4	80.3	104.8	116.3
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.5	9.5	17.3	27.4	44.6	64.9	91.8	105.5	128.1
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of larynx (C32)

All races 3/, both sexes

2003.....	0.3	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.6	2.9	4.6	6.1	7.4	8.0	8.2	6.4
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0

All races 3/, male

2003.....	0.5	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.6	5.0	8.1	10.2	13.0	14.5	16.9	16.5
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8

All races 3/, female

2003.....	*	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.5	2.6	2.9	3.3	2.9	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of larynx (C32)

White, both sexes

2003.....	0.3	0.6	1.1	2.3	4.2	5.5	6.2	6.3	7.1	6.2
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	4.0	5.7	6.7	7.5	8.0	6.5
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7

White, male

2003.....	0.5	1.1	2.0	3.7	7.2	9.3	10.4	11.9	14.4	14.5
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.9	9.4	11.8	13.7	16.6	16.8
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8

White, female

2003.....	*	*	0.3	0.9	1.4	2.1	2.7	2.3	2.5	2.5
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.7	3.1	2.8	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of larynx (C32)

Black, both sexes

2003.....	*	1.7	3.6	5.8	9.7	10.6	12.7	11.1	12.7	11.1
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.2	7.6	10.9	10.2	15.2	15.7	12.1	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3

Black, male

2003.....	*	2.7	6.8	10.7	17.5	22.7	25.3	25.6	31.7	27.1
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.3	8.0	15.0	21.6	18.9	28.7	28.2	26.6	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6

Black, female

2003.....	*	*	*	*	3.6	*	*	*	*	*
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.7	5.7	7.6	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2003.....	8.5	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.2	44.6	93.5	162.0	248.7	329.7	368.4	378.0	298.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0

All races 3/, male

2003.....	9.5	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.4	26.0	54.6	118.0	206.2	327.5	444.0	507.3	549.6	494.5
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2

All races 3/, female

2003.....	7.5	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.6	35.1	70.9	122.3	181.6	238.7	268.6	272.8	213.5
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, both sexes

2003.....	8.2	19.9	39.1	80.7	149.9	233.5	319.7	381.3	384.5	302.4
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.5	19.6	42.7	92.3	162.9	251.8	332.6	370.8	383.1	301.1
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8

White, male

2003.....	8.9	22.8	46.9	97.3	180.4	291.1	412.1	513.0	541.9	478.5
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.2	23.6	50.4	112.8	203.2	325.2	441.6	502.6	551.6	498.3
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5

White, female

2003.....	7.5	17.0	31.5	64.8	121.6	182.6	244.0	286.0	287.4	224.4
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	7.9	15.7	35.3	72.8	126.0	188.2	244.7	275.2	279.4	216.8
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2003.....	12.4	35.4	62.0	114.8	182.9	256.6	328.9	387.1	377.2	280.9
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	36.0	71.7	124.9	190.8	269.2	356.0	396.3	358.3	288.8
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5

Black, male

2003.....	15.6	45.1	86.6	167.9	267.8	376.2	495.0	618.5	628.3	502.5
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	48.8	106.4	193.4	284.0	414.0	535.3	632.8	607.7	519.3
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2

Black, female

2003.....	9.6	26.9	41.2	71.2	116.7	167.8	216.9	249.6	245.6	194.8
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.2	43.3	71.5	121.5	161.7	228.7	244.0	221.0	192.5
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant melanoma of skin (C43)

All races 3/, both sexes

2003.....	1.6	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.9	4.7	6.3	8.0	10.1	12.0	15.0	18.7
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4

All races 3/, male

2003.....	1.9	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.3	5.3	6.6	8.8	11.4	14.8	18.0	24.4	30.2
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5

All races 3/, female

2003.....	1.3	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.1	6.4	7.6	9.3	13.8
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant melanoma of skin (C43)

White, both sexes

2003.....	1.9	2.9	3.9	5.6	7.3	8.3	11.6	15.2	17.0	19.1
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.2	4.5	5.3	7.2	9.0	11.2	13.1	16.1	20.1
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6

White, male

2003.....	2.3	3.6	5.3	7.9	10.1	11.7	17.1	24.4	27.1	32.9
2002.....	2.5	3.8	5.1	6.9	9.7	12.7	16.4	21.8	25.6	31.7
2001.....	2.6	3.8	5.5	7.2	10.0	13.3	18.0	21.5	26.1	32.7
2000.....	2.4	3.9	6.1	7.4	10.0	12.8	16.3	19.7	26.2	33.1
1999.....	2.7	4.1	5.3	7.7	9.5	13.4	16.2	21.5	24.5	29.7

White, female

2003.....	1.5	2.3	2.6	3.5	4.6	5.2	7.1	8.5	10.8	12.9
2002.....	1.5	1.7	2.9	3.2	4.2	5.5	6.7	8.8	10.7	13.8
2001.....	1.7	2.3	3.1	3.5	4.2	4.8	7.0	8.0	9.8	13.7
2000.....	1.6	2.5	2.9	3.4	4.6	5.8	7.0	8.3	9.9	14.5
1999.....	1.5	2.2	3.0	3.3	4.6	5.8	6.4	8.8	9.9	12.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2003.....	7.9	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.2	13.7	21.2	28.5	35.5	43.9	55.9	70.9	91.7	145.5
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4

All races 3/, male

2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.8	2.1	2.4	3.9
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5

All races 3/, female

2003.....	15.6	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.3	26.8	40.9	54.1	66.6	80.3	99.0	120.4	146.4	206.4
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

White, both sexes

2003.....	7.1	11.4	17.8	25.3	32.6	39.2	51.7	66.0	87.5	132.5
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.4	12.4	20.0	27.1	34.7	44.0	56.2	71.2	92.3	145.6
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0

White, male

2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.9	3.9
2002.....	*	*	0.3	0.4	0.6	0.6	1.3	1.7	2.7	3.3
2001.....	*	*	0.3	0.5	0.7	1.2	1.6	2.1	2.3	3.3
2000.....	*	*	0.3	0.6	0.8	1.1	1.7	2.1	2.2	3.7
1999.....	*	*	0.3	0.5	0.6	1.0	1.3	1.8	2.2	4.4

White, female

2003.....	14.2	22.7	34.8	49.2	62.0	72.9	93.0	112.7	139.7	189.4
2002.....	13.9	23.2	36.3	48.6	63.1	76.2	93.3	114.7	142.0	192.6
2001.....	14.7	24.5	38.1	50.1	63.4	75.7	95.8	117.0	140.0	189.9
2000.....	14.8	24.6	39.0	52.1	65.9	81.1	100.2	121.2	147.7	206.3
1999.....	14.2	23.9	38.9	51.7	64.4	79.2	98.6	117.4	146.3	202.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

Black, both sexes

2003.....	14.1	22.6	31.2	43.4	48.1	54.6	65.1	80.3	106.4	161.6
2002.....	15.3	22.9	32.1	40.5	46.3	54.5	67.2	78.5	117.5	152.6
2001.....	14.8	22.6	33.1	42.4	47.4	58.0	70.4	83.5	105.9	152.0
2000.....	14.8	24.3	34.0	44.5	47.8	50.7	65.1	84.5	104.9	176.9
1999.....	14.6	24.2	34.7	44.4	48.6	57.8	69.9	88.0	104.9	147.3

Black, male

2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2003.....	26.3	42.0	57.3	78.3	84.7	94.1	107.4	127.2	160.5	222.4
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.6	61.3	78.1	82.2	87.8	109.4	137.5	160.3	247.2
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2003.....	1.7	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.3	2.7	3.1	3.1	3.6	3.9	4.3	5.7	7.2
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	3.3	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.3	6.0	5.9	6.7	7.1	7.4	9.2	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2003.....	1.6	1.9	2.1	2.6	2.6	2.8	2.4	2.8	4.6	4.9
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.5	2.0	2.4	2.7	2.7	3.0	3.3	3.8	5.1	6.2
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	3.2	3.7	4.2	5.1	4.9	5.2	4.4	4.9	7.4	7.1
2002.....	3.1	3.7	4.3	4.9	4.5	5.2	5.4	6.2	5.7	7.3
2001.....	3.5	4.3	4.6	4.6	5.5	4.4	5.5	5.8	7.6	8.8
2000.....	3.1	4.0	4.7	5.3	5.1	5.6	6.0	6.6	8.2	8.9
1999.....	3.4	4.5	5.0	5.2	5.7	5.9	5.5	6.8	6.5	9.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2003.....	2.4	3.5	3.8	5.1	6.1	7.1	7.7	11.2	11.4	14.6
2002.....	3.0	3.7	4.6	4.2	6.0	8.2	10.1	9.6	12.6	17.5
2001.....	3.0	3.2	4.4	4.6	5.9	7.9	7.7	9.7	15.2	19.0
2000.....	3.1	3.8	5.1	6.2	6.5	9.0	9.7	10.9	14.1	19.8
1999.....	3.0	3.0	4.1	6.1	8.3	8.7	13.1	12.3	9.5	21.7

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	4.5	6.5	7.1	9.3	10.8	12.4	12.8	17.9	17.4	20.3
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.2	11.1	11.3	15.7	16.5	17.9	21.9	28.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2003.....	0.5	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	1.0	1.8	3.6	6.0	8.5	11.2	14.0	18.7	24.4
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	1.0	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.6	6.9	11.5	15.7	20.1	24.1	30.2	34.9
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2003.....	0.4	0.7	1.9	3.4	5.4	7.9	10.5	13.0	16.2	22.7
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	1.0	1.8	3.4	5.4	7.4	10.0	13.0	17.6	24.1
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	0.8	1.5	3.7	6.6	10.4	14.9	19.1	22.4	26.2	32.8
2002.....	0.9	1.7	3.7	6.3	10.1	13.6	19.6	22.4	26.3	34.2
2001.....	0.8	1.7	3.8	6.1	9.9	13.6	18.9	23.5	29.8	34.7
2000.....	0.8	1.9	3.5	6.7	10.4	13.8	18.0	22.5	28.5	34.4
1999.....	0.9	1.8	3.2	5.9	9.9	14.2	18.4	23.3	30.0	32.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2003.....	0.9	1.1	2.5	5.3	13.9	19.9	22.4	25.5	31.2	30.0
2002.....	*	1.1	2.9	6.6	10.8	20.0	25.2	27.8	29.0	38.9
2001.....	*	1.4	2.9	4.6	11.6	18.8	24.2	26.6	31.1	32.2
2000.....	*	1.0	2.4	5.3	12.1	18.8	26.2	27.8	36.5	33.2
1999.....	0.8	*	2.2	5.8	10.5	20.9	21.8	25.7	29.8	35.4

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	1.6	2.1	4.7	9.6	24.8	34.7	37.4	40.7	47.6	41.7
2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.5	21.1	32.8	44.8	45.7	56.6	47.1
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of ovary (C56)

All races 3/, both sexes

2003.....	1.4	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.4	3.1	5.6	8.4	12.4	17.7	22.1	31.0	36.8	40.4
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6

All races 3/, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2003.....	2.8	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.1	10.9	16.2	23.5	32.8	39.7	53.2	59.3	57.8
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of ovary (C56)

White, both sexes

2003.....	1.4	3.0	5.4	8.7	13.5	17.9	22.4	30.3	38.1	39.1
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.3	5.7	8.8	12.9	18.3	22.7	32.0	37.8	41.5
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	2.9	6.0	10.6	17.1	25.9	33.7	40.7	52.1	61.5	56.4
2002.....	2.9	6.6	11.4	16.8	25.9	33.9	41.6	53.4	62.5	55.9
2001.....	2.9	6.2	11.4	16.6	25.3	33.9	42.2	53.1	61.9	57.3
2000.....	3.0	6.5	11.2	17.1	24.7	34.1	41.0	55.2	61.1	59.3
1999.....	3.2	6.5	11.2	16.8	25.1	32.8	41.0	50.5	57.9	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of ovary (C56)

Black, both sexes

2003.....	1.2	2.8	5.1	6.8	11.0	15.3	20.6	26.2	29.4	36.8
2002.....	1.2	2.9	4.1	7.7	10.9	14.9	20.0	26.5	32.4	34.7
2001.....	1.4	1.9	4.4	7.9	12.6	16.3	20.7	26.2	27.9	36.5
2000.....	1.5	2.2	4.9	6.5	11.0	16.1	19.4	26.2	28.6	34.5
1999.....	1.1	2.4	5.0	7.2	11.4	15.0	22.7	28.9	28.3	28.7

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	2.3	5.3	9.4	12.5	19.6	26.7	34.6	41.8	44.9	51.0
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.1	8.9	11.5	19.2	28.1	33.2	43.1	44.3	48.9
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races 3/, both sexes

2003.....	1.0	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.3	4.6	7.7	11.0	15.2	19.6	24.5	29.8	33.3
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4

All races 3/, male

2003.....	1.5	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.2	6.8	11.4	15.9	21.7	27.6	34.0	43.5	52.3
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4

All races 3/, female

2003.....	0.6	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.5	4.3	6.5	9.6	13.3	17.7	21.5	25.2
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

White, both sexes

2003.....	1.0	2.0	4.2	7.2	11.6	15.0	20.2	25.4	30.8	33.8
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.7	7.9	11.3	15.6	20.0	24.9	30.2	34.1
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5

White, male

2003.....	1.5	3.1	5.9	10.7	17.5	20.8	28.2	36.5	46.2	50.0
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.8	11.5	16.3	21.9	28.0	34.5	43.9	53.9
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8

White, female

2003.....	0.6	0.9	2.5	3.8	6.2	9.7	13.7	17.4	21.3	26.6
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.6	10.1	13.5	17.9	21.8	25.6
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2003.....	1.5	2.5	4.1	8.2	11.2	13.4	16.1	22.3	26.1	33.8
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.8	7.5	10.5	14.2	18.4	23.5	30.3	29.8
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2

Black, male

2003.....	2.5	3.6	6.5	11.5	17.0	19.5	25.9	35.6	39.1	43.8
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.3	7.5	11.9	16.0	23.9	26.8	31.0	48.2	45.8
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2

Black, female

2003.....	*	1.5	2.1	5.5	6.6	9.0	9.5	14.4	19.3	30.0
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.1	6.4	7.0	12.4	18.8	20.6	23.1
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of bladder (C67)

All races 3/, both sexes

2003.....	0.3	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.7	6.9	11.9	19.9	29.1	45.9	68.7
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5

All races 3/, male

2003.....	0.4	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.8	10.9	19.3	32.4	49.7	81.1	132.2
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8

All races 3/, female

2003.....	0.2	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.9	14.2	24.4	41.4
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of bladder (C67)

White, both sexes

2003.....	0.3	0.8	1.8	3.7	7.3	12.2	19.6	31.1	46.2	66.6
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.3	12.5	20.5	29.7	46.7	71.2
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8

White, male

2003.....	0.4	1.4	2.8	5.7	11.7	19.8	31.6	53.8	83.5	131.0
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.6	20.6	33.9	51.7	83.4	138.9
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7

White, female

2003.....	*	0.2	0.8	1.7	3.3	5.5	9.8	14.6	23.3	38.1
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.7	13.7	24.1	42.3
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of bladder (C67)

Black, both sexes

2003.....	*	1.2	2.2	3.2	5.0	11.4	12.7	26.5	40.9	53.4
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.9	1.3	3.1	5.5	9.1	17.7	28.5	46.6	49.9
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3

Black, male

2003.....	*	1.7	2.8	4.5	6.6	18.2	16.4	41.1	61.3	80.3
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.8	7.6	12.0	22.9	39.6	72.2	81.9
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7

Black, female

2003.....	*	*	1.7	*	3.8	6.4	10.2	17.9	30.2	42.9
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.0	14.0	21.4	32.4	36.5
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2003.....	2.9	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.8	4.1	6.4	8.3	11.5	14.9	18.6	19.4	21.2	16.7
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All races 3/, male

2003.....	3.4	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.9	10.3	14.3	18.2	23.0	22.7	25.9	21.3
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All races 3/, female

2003.....	2.4	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	5.0	6.5	9.0	12.2	15.1	17.0	18.4	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2003.....	3.3	4.3	6.3	9.1	12.2	14.9	18.5	20.6	20.7	15.7
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.0	4.5	7.0	8.9	12.5	16.2	20.0	20.5	22.0	17.3
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6

White, male

2003.....	3.8	5.5	8.0	10.3	14.8	17.4	22.6	25.3	26.6	20.7
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	3.9	5.4	8.4	11.0	15.4	19.7	24.5	23.9	27.1	22.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8

White, female

2003.....	2.7	3.0	4.8	7.9	9.8	12.7	15.2	17.3	17.0	13.4
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.6	6.9	9.8	13.1	16.3	18.0	18.8	15.1
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2003.....	1.6	2.7	3.6	5.5	5.6	8.3	9.7	10.6	10.4	10.5
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.9	5.5	6.1	7.2	8.7	10.3	15.0	10.7
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9

Black, male

2003.....	2.4	3.1	5.1	6.6	6.7	11.8	12.2	16.0	*	*
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.3	6.0	6.9	8.0	7.5	11.8	11.4	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*

Black, female

2003.....	*	2.3	2.2	4.6	4.7	5.7	8.0	7.3	8.5	11.7
2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.4	4.7	7.0	6.6	9.7	15.1	11.1
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2003.....	4.7	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.4	13.6	23.7	38.7	62.9	92.8	131.5	177.6	201.9
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All races 3/, male

2003.....	5.8	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.4	16.5	28.9	46.7	79.5	118.6	166.7	227.1	267.4
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All races 3/, female

2003.....	3.5	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.9	18.9	31.5	48.8	72.2	106.1	147.2	173.7
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, both sexes

2003.....	4.6	7.0	11.3	20.1	35.8	56.1	88.7	130.3	169.5	204.5
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.6	8.1	13.3	23.5	38.9	64.4	94.7	134.4	182.6	206.3
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1

White, male

2003.....	5.7	8.7	13.9	25.1	46.3	73.6	115.2	168.7	224.3	279.5
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.7	10.0	15.9	28.6	46.7	81.9	120.3	171.0	234.4	276.8
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8

White, female

2003.....	3.4	5.4	8.8	15.3	26.1	40.7	67.0	102.6	135.8	171.3
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.8	18.7	31.7	49.2	74.0	107.9	150.7	176.2
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2003.....	6.3	9.5	17.3	23.5	41.1	59.4	90.5	117.4	139.1	150.5
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.7	18.1	29.6	44.5	59.8	89.7	114.7	137.3	175.9
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0

Black, male

2003.....	8.0	11.9	22.5	32.0	51.1	73.4	105.1	137.8	166.8	184.5
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	15.3	23.2	38.3	56.7	70.0	119.5	142.5	165.2	207.5
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3

Black, female

2003.....	4.8	7.4	12.9	16.6	33.2	49.0	80.6	105.3	124.6	137.3
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.5	14.1	22.7	35.5	52.2	68.5	96.8	121.9	162.7
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hodgkin's disease (C81)

All races 3/, both sexes

2003.....	0.3	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.5	2.4
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4

All races 3/, male

2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	2.0	2.3	2.7	3.5
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8

All races 3/, female

2003.....	0.2	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.5	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hodgkin's disease (C81)

White, both sexes

2003.....	0.4	0.4	0.5	0.6	0.9	0.8	1.7	2.2	2.4	2.5
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.1	1.6	2.0	2.6	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4

White, male

2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	3.0	2.7
2002.....	0.4	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.5	3.9
2001.....	0.5	0.5	0.7	0.6	0.9	1.6	1.8	2.5	3.0	2.7
2000.....	0.4	0.5	0.6	0.5	1.1	1.5	1.9	2.5	2.8	3.7
1999.....	0.5	0.5	0.6	0.9	1.3	1.7	2.4	2.5	3.0	3.8

White, female

2003.....	0.2	0.3	0.4	0.5	0.6	0.6	1.4	2.0	2.0	2.4
2002.....	0.3	0.3	0.5	*	0.7	0.6	1.2	1.6	1.7	2.5
2001.....	0.3	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.2
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.4	1.6	2.4	2.1
1999.....	0.3	0.4	0.4	0.5	0.6	0.9	1.6	2.0	2.2	1.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.0	3.7	5.9	10.8	16.3	25.3	36.7	53.8	72.4	79.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7

All races 3/, male

2003.....	2.2	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.8	7.7	13.5	19.5	31.0	45.5	65.1	86.7	100.4
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4

All races 3/, female

2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.2	8.2	13.4	20.5	29.8	45.6	63.7	70.0
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2003.....	1.7	2.9	4.7	8.4	14.9	22.8	34.2	51.8	69.0	83.3
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	6.0	11.3	17.1	26.7	38.6	56.5	76.3	83.4
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2

White, male

2003.....	2.1	3.7	6.0	10.6	19.4	29.6	43.1	63.9	86.5	109.7
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.7	13.9	20.2	32.9	47.3	68.5	91.3	106.7
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0

White, female

2003.....	1.2	2.1	3.5	6.4	10.7	16.7	27.0	43.0	58.2	71.6
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.2	21.4	31.6	47.8	67.1	73.4
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

Black, both sexes

2003.....	2.3	3.9	5.9	6.1	12.7	14.6	22.8	26.5	31.0	31.2
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.6	5.7	9.0	12.7	15.9	22.5	26.0	29.5	36.4
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5

Black, male

2003.....	3.0	5.7	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.3	8.4	14.1	16.4	18.4	30.6	29.6	31.5	40.4
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8

Black, female

2003.....	1.6	2.3	4.5	4.7	10.1	12.6	21.5	25.2	29.0	30.0
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	4.9	10.0	14.1	16.7	23.8	28.3	34.7
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Leukemia (C91-C95)

All races 3/, both sexes

2003.....	2.1	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.6	7.8	12.9	22.0	33.6	47.4	65.6	82.5
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4

All races 3/, male

2003.....	2.4	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.2	9.8	16.3	29.7	46.6	63.9	91.9	113.7
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2

All races 3/, female

2003.....	1.7	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	4.0	6.0	9.8	15.4	23.3	35.6	49.5	69.0
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Leukemia (C91-C95)

White, both sexes

2003.....	2.1	2.8	4.0	7.1	12.7	20.3	33.6	49.3	65.0	83.4
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.6	7.9	13.0	22.9	34.8	49.2	68.0	85.1
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4

White, male

2003.....	2.4	3.4	4.8	9.0	16.4	27.7	46.4	69.2	92.8	116.9
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.1	3.5	5.1	9.9	16.4	31.1	47.9	66.3	95.3	117.9
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1

White, female

2003.....	1.7	2.3	3.2	5.4	9.3	13.8	23.1	34.9	47.9	68.6
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.1	5.9	9.9	15.9	24.3	36.7	51.2	71.1
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Leukemia (C91-C95)

Black, both sexes

2003.....	2.5	2.5	5.1	7.2	12.8	17.9	29.6	37.3	48.0	59.2
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.6	5.4	8.0	14.2	17.2	28.2	34.8	45.7	66.2
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8

Black, male

2003.....	3.1	2.7	6.8	10.7	14.5	24.9	39.4	50.2	58.3	77.2
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	5.1	6.5	10.5	19.5	21.4	41.4	47.8	63.1	90.5
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3

Black, female

2003.....	2.1	2.4	3.8	4.3	11.4	12.8	23.0	29.6	42.6	52.3
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.3	14.1	18.8	26.4	36.1	56.0
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes

2003.....	0.5	1.1	2.5	4.4	7.9	13.1	20.4	28.5	34.5	36.3
2002.....	0.4	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001.....	0.5	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000.....	0.5	1.2	2.6	4.6	8.6	14.6	20.6	28.3	36.7	37.3
1999.....	0.5	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6

All races 3/, male

2003.....	0.7	1.3	3.1	5.4	10.1	16.0	24.3	33.9	43.3	50.4
2002.....	0.5	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001.....	0.7	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000.....	0.6	1.5	3.0	5.0	9.7	17.3	24.4	35.1	45.4	49.1
1999.....	0.6	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3

All races 3/, female

2003.....	0.4	0.9	1.9	3.5	6.0	10.6	17.2	24.7	29.2	30.1
2002.....	0.4	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001.....	0.4	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000.....	0.4	1.0	2.3	4.2	7.7	12.2	17.6	23.4	31.4	32.3
1999.....	0.4	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

White, both sexes

2003.....	0.5	0.9	2.1	3.9	7.3	12.1	19.1	26.9	33.1	35.0
2002.....	0.4	1.1	2.1	4.2	7.5	13.0	19.9	26.9	34.8	35.7
2001.....	0.4	1.0	2.1	4.4	7.5	12.9	19.9	27.1	33.0	35.0
2000.....	0.4	1.0	2.3	3.9	7.9	13.6	19.4	26.5	35.5	34.8
1999.....	0.4	1.1	2.1	4.7	7.9	13.2	19.2	25.7	34.9	34.8

White, male

2003.....	0.6	1.1	2.6	4.8	9.4	15.1	23.4	32.9	41.9	49.7
2002.....	0.4	1.3	2.7	4.9	9.4	15.8	24.9	34.8	45.3	50.3
2001.....	0.6	1.3	2.5	5.2	9.0	15.7	24.1	34.6	41.7	47.1
2000.....	0.5	1.2	2.5	4.3	8.9	16.4	23.0	33.3	44.7	47.9
1999.....	0.4	1.3	2.6	5.6	10.0	15.8	22.5	32.4	44.3	49.7

White, female

2003.....	0.3	0.7	1.6	3.1	5.4	9.5	15.6	22.6	27.7	28.5
2002.....	0.3	0.8	1.6	3.4	5.7	10.5	15.8	21.2	28.4	29.4
2001.....	*	0.7	1.6	3.7	6.1	10.4	16.5	21.7	27.8	29.9
2000.....	0.2	0.8	2.0	3.4	6.9	11.2	16.5	21.6	29.8	29.2
1999.....	0.3	1.0	1.5	3.8	5.9	10.9	16.6	21.0	29.4	28.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2003.....	1.2	2.8	5.9	9.4	15.1	25.6	37.3	52.6	59.4	59.8
2002.....	1.0	2.9	5.6	10.7	16.3	26.7	37.7	44.7	54.9	55.5
2001.....	1.5	2.6	4.8	11.8	19.4	25.8	38.2	47.9	61.2	58.3
2000.....	1.1	3.1	6.5	11.9	16.7	26.2	37.6	53.5	60.4	71.8
1999.....	1.4	2.4	6.6	13.8	20.0	24.2	37.6	51.1	56.3	63.1

Black, male

2003.....	1.5	3.2	7.7	12.2	19.7	29.3	39.4	57.5	73.1	73.0
2002.....	*	3.5	5.9	12.4	20.2	31.4	44.7	52.4	70.3	71.8
2001.....	1.6	3.4	5.3	14.5	23.4	30.2	45.1	57.1	80.3	71.5
2000.....	*	3.4	7.6	12.9	19.3	29.7	45.9	64.6	68.1	73.4
1999.....	1.9	2.6	6.7	16.1	21.6	26.2	46.8	59.4	74.1	80.9

Black, female

2003.....	*	2.4	4.4	7.0	11.6	22.8	35.9	49.6	52.2	54.7
2002.....	*	2.4	5.3	9.3	13.3	23.1	32.9	40.1	46.9	49.3
2001.....	1.5	2.0	4.3	9.5	16.3	22.5	33.6	42.4	51.5	53.5
2000.....	*	2.9	5.6	11.1	14.8	23.7	31.6	46.3	56.2	71.1
1999.....	*	2.3	6.6	11.9	18.8	22.7	31.5	46.1	47.5	56.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2003.....	5.3	10.3	17.7	28.1	44.4	64.6	92.9	131.5	167.7	224.0
2002.....	5.8	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001.....	5.9	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000.....	5.8	10.6	18.5	30.7	48.8	72.3	101.8	136.1	179.0	236.6
1999.....	6.7	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4

All races 3/, male

2003.....	5.9	11.9	20.1	33.0	52.9	78.3	111.3	162.7	208.2	275.0
2002.....	6.2	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001.....	6.4	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000.....	6.4	12.1	21.1	35.8	58.3	87.5	123.5	164.5	218.1	285.3
1999.....	7.6	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9

All races 3/, female

2003.....	4.8	8.7	15.5	23.6	36.7	52.8	78.1	109.2	142.9	201.4
2002.....	5.4	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001.....	5.3	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000.....	5.3	9.2	16.0	26.0	40.3	59.4	84.5	115.7	155.0	215.7
1999.....	5.8	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2003.....	5.0	9.7	17.0	27.3	44.0	64.7	93.5	132.7	170.1	227.9
2002.....	5.5	9.9	16.8	28.1	45.7	65.6	96.9	131.7	170.0	233.0
2001.....	5.6	9.9	17.2	29.0	45.3	66.5	97.4	134.1	171.4	238.1
2000.....	5.4	9.9	17.4	30.0	47.6	72.5	101.3	135.6	180.6	241.0
1999.....	6.3	10.8	18.3	31.1	51.2	76.3	107.1	142.3	189.4	252.4

White, male

2003.....	5.4	11.0	19.2	31.8	52.2	78.0	111.6	164.1	212.2	280.9
2002.....	5.7	11.1	20.0	32.0	55.1	80.2	118.8	162.2	210.9	296.5
2001.....	6.2	11.3	20.1	33.1	53.5	78.0	119.7	163.1	212.0	300.2
2000.....	5.8	11.2	19.9	34.5	55.4	87.9	123.1	162.8	221.8	292.8
1999.....	7.2	12.2	20.8	36.6	60.9	90.1	131.8	174.6	235.3	323.6

White, female

2003.....	4.6	8.4	14.9	23.1	36.3	52.9	78.8	110.0	144.1	204.4
2002.....	5.4	8.7	13.7	24.3	36.9	52.7	79.1	109.9	144.9	205.5
2001.....	5.1	8.5	14.3	25.1	37.7	56.3	79.3	113.7	146.8	212.1
2000.....	5.0	8.7	15.0	25.7	40.4	59.1	83.7	116.0	155.2	218.8
1999.....	5.3	9.5	15.8	26.0	42.3	64.3	87.5	119.6	162.3	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2003.....	7.8	16.0	26.5	41.1	57.7	77.0	102.4	139.4	166.0	206.3
2002.....	8.0	15.8	25.3	41.5	63.3	82.5	113.5	146.5	174.5	222.1
2001.....	7.7	16.7	27.8	44.1	64.1	85.2	120.3	140.7	182.3	217.1
2000.....	10.0	17.1	29.8	41.9	65.6	82.3	116.6	156.2	179.4	211.4
1999.....	10.1	18.3	29.0	48.3	73.5	85.4	128.3	157.3	193.0	242.6

Black, male

2003.....	9.4	19.8	31.4	53.0	72.3	98.5	128.4	176.2	206.0	241.9
2002.....	9.6	18.4	30.9	50.2	82.0	97.2	142.9	187.4	209.4	267.0
2001.....	8.2	19.0	35.9	57.1	80.9	110.0	151.5	178.9	219.1	283.7
2000.....	11.8	21.4	35.1	53.6	93.1	97.7	142.1	204.0	205.1	234.1
1999.....	11.7	23.2	36.5	61.9	94.0	109.8	163.8	191.4	233.7	292.9

Black, female

2003.....	6.4	12.6	22.3	31.3	46.3	61.0	84.8	117.5	145.1	192.4
2002.....	6.5	13.5	20.5	34.5	48.6	71.6	93.9	122.1	156.4	205.0
2001.....	7.3	14.8	20.9	33.5	50.9	66.8	99.5	117.9	163.5	192.4
2000.....	8.3	13.5	25.4	32.8	45.1	70.9	98.5	125.5	165.3	201.9
1999.....	8.8	14.0	22.7	37.2	57.4	67.3	104.6	136.7	172.9	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2003.....	0.9	1.4	2.0	3.2	6.0	9.8	16.3	28.5	47.4	80.5
2002.....	1.0	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000.....	0.9	1.5	2.3	3.9	6.7	10.7	17.6	28.9	51.1	83.8
1999.....	0.9	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9

All races 3/, male

2003.....	1.0	1.4	2.3	3.6	7.0	11.6	21.2	35.9	61.5	101.7
2002.....	1.0	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001.....	1.0	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000.....	0.9	1.6	2.7	4.2	7.7	13.6	21.4	36.1	64.3	103.3
1999.....	1.1	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2

All races 3/, female

2003.....	0.7	1.3	1.7	2.8	5.1	8.3	12.4	23.2	38.7	71.1
2002.....	0.9	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001.....	0.8	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000.....	1.0	1.3	1.9	3.5	5.8	8.2	14.6	23.7	42.9	75.4
1999.....	0.8	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2003.....	0.9	1.3	2.0	3.1	6.1	9.8	16.8	29.5	48.7	83.0
2002.....	0.9	1.6	2.0	3.7	5.6	9.9	16.7	29.7	47.2	80.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.7	18.1	30.4	48.9	83.9
2000.....	0.9	1.4	2.3	3.9	6.7	10.9	17.5	29.5	52.6	86.4
1999.....	0.9	1.3	2.2	3.5	6.2	10.7	19.0	29.7	48.7	85.3

White, male

2003.....	1.0	1.3	2.3	3.5	7.1	11.8	21.8	37.4	63.3	105.5
2002.....	1.1	1.7	2.1	4.4	6.4	12.4	20.7	38.4	60.1	104.2
2001.....	1.0	1.7	2.5	4.1	7.4	13.2	23.0	40.4	63.7	109.9
2000.....	0.8	1.5	2.7	4.3	7.4	13.7	21.2	36.9	67.0	107.4
1999.....	1.1	1.4	2.6	3.9	8.0	13.3	24.6	37.7	62.9	114.3

White, female

2003.....	0.7	1.2	1.7	2.8	5.3	8.0	12.7	23.9	39.8	73.0
2002.....	0.8	1.4	1.8	3.0	4.8	7.6	13.5	23.4	39.2	70.5
2001.....	0.8	1.2	2.0	3.8	5.2	8.6	14.2	23.3	40.0	73.0
2000.....	1.0	1.2	1.9	3.5	6.0	8.5	14.5	24.1	43.7	77.4
1999.....	0.7	1.2	1.8	3.0	4.6	8.3	14.7	24.0	40.4	73.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2003.....	1.0	2.2	2.7	4.2	6.6	11.2	15.3	22.6	39.9	57.8
2002.....	1.7	2.0	3.0	5.2	5.8	10.6	15.0	24.4	41.9	63.6
2001.....	1.2	1.4	2.7	4.5	7.6	10.9	18.6	24.5	42.5	73.4
2000.....	1.4	2.4	3.2	4.3	7.2	10.2	20.1	26.8	39.5	64.3
1999.....	1.3	2.2	3.2	5.0	7.8	13.7	18.1	29.5	38.7	61.9

Black, male

2003.....	*	2.1	2.9	5.5	8.5	12.3	21.4	26.5	53.2	66.7
2002.....	*	1.7	2.8	7.4	6.5	12.7	16.0	29.9	53.5	79.3
2001.....	*	*	2.4	4.5	9.2	13.0	22.9	31.6	58.6	88.5
2000.....	*	2.3	3.8	4.6	10.6	15.0	22.6	33.2	38.2	70.2
1999.....	*	*	3.3	6.1	8.3	16.9	21.7	36.1	42.7	63.3

Black, female

2003.....	*	2.3	2.6	3.1	5.1	10.4	11.3	20.3	32.9	54.3
2002.....	2.1	2.2	3.3	3.3	5.3	9.1	14.3	21.1	35.8	57.5
2001.....	*	*	3.0	4.5	6.4	9.4	15.7	20.2	34.3	67.8
2000.....	1.3	2.5	2.7	4.0	4.7	6.7	18.3	22.6	40.2	61.8
1999.....	1.4	2.5	3.0	4.0	7.5	11.4	15.8	25.5	36.7	61.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

All races 3/, both sexes

2003.....	0.4	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.6	0.7	1.0	1.2	2.2	3.5	6.5	14.1	41.8
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8

All races 3/, male

2003.....	0.5	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.3	2.2	3.6	6.4	14.3	38.0
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4

All races 3/, female

2003.....	0.4	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.3	3.4	6.6	13.9	43.4
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

White, both sexes

2003.....	0.1	0.3	0.3	0.4	0.9	1.6	3.2	5.7	12.4	40.6
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	2.0	3.2	6.2	13.9	41.6
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0

White, male

2003.....	*	0.3	0.3	0.5	1.1	2.0	3.3	6.3	13.4	37.9
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	1.0	2.1	3.4	6.0	14.0	37.3
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4

White, female

2003.....	*	*	0.3	0.3	0.8	1.3	3.1	5.3	11.8	41.8
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.1	6.4	13.8	43.4
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

Black, both sexes

2003.....	2.5	3.3	2.7	4.1	4.4	4.3	7.1	10.9	19.8	49.3
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	3.0	3.3	3.0	4.0	4.6	6.1	10.3	18.9	51.1
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9

Black, male

2003.....	2.9	3.9	2.6	4.9	6.2	5.7	10.8	12.8	18.5	49.0
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.3	3.6	*	*	*	*	10.5	19.1	51.1
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5

Black, female

2003.....	2.1	2.8	2.7	3.3	3.0	*	4.7	9.8	20.5	49.4
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	2.9	3.9	5.6	6.3	10.3	18.7	51.1
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2003.....	5.8	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.6	17.6	30.0	48.7	74.8	110.0	156.6	215.8	315.6
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2

All races 3/, male

2003.....	7.4	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.7	11.5	20.2	34.0	55.0	82.1	120.6	170.6	238.8	327.4
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7

All races 3/, female

2003.....	4.2	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	15.1	26.3	43.1	68.7	101.6	146.6	201.6	310.5
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

White, both sexes

2003.....	5.1	8.8	15.2	26.4	42.7	66.5	98.1	143.1	202.7	300.8
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.1	14.8	25.9	42.7	66.1	99.5	143.7	200.3	300.1
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9

White, male

2003.....	6.6	11.1	18.7	31.6	51.5	79.1	115.3	170.2	241.1	333.4
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.2	9.8	17.6	29.9	49.3	75.2	112.9	161.1	230.6	321.8
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4

White, female

2003.....	3.5	6.5	11.8	21.4	34.5	55.4	84.0	123.5	179.1	286.3
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.1	22.2	36.7	58.3	88.8	131.0	181.7	290.8
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

Black, both sexes

2003.....	11.7	22.1	39.9	69.8	106.9	156.1	221.2	300.0	389.9	527.5
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	22.3	42.2	67.2	103.6	153.1	221.7	317.7	422.5	526.3
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0

Black, male

2003.....	14.4	25.4	44.0	77.3	121.7	170.6	232.6	306.3	387.6	453.5
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.7	26.3	46.1	75.3	113.4	147.8	213.8	300.0	376.9	417.2
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2

Black, female

2003.....	9.3	19.2	36.4	63.6	95.4	145.3	213.6	296.3	391.1	556.2
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.9	39.0	61.0	96.2	157.0	227.4	329.0	447.6	571.8
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional deficiencies (E40-E64)

All races 3/, both sexes

2003.....	0.1	0.2	0.3	0.5	0.8	1.3	2.5	5.7	10.4	35.5
2002.....	0.1	0.2	0.3	0.5	0.8	1.6	2.7	6.1	12.8	41.5
2001.....	0.1	0.2	0.3	0.5	0.8	1.4	2.8	5.9	12.8	42.9
2000.....	0.1	0.2	0.3	0.5	1.0	2.0	3.1	6.8	15.9	49.4
1999.....	0.2	0.2	0.4	0.6	0.9	1.6	3.2	6.4	15.9	53.9

All races 3/, male

2003.....	*	*	0.3	0.5	0.7	1.4	2.7	5.6	10.9	30.6
2002.....	*	0.2	0.3	0.6	0.9	1.6	3.0	6.3	12.0	37.0
2001.....	*	0.3	0.3	0.5	0.9	1.4	2.9	5.6	13.2	35.2
2000.....	*	0.3	0.3	0.7	1.0	2.0	3.5	6.7	14.2	41.4
1999.....	0.2	0.2	0.4	0.7	1.1	1.6	3.8	6.8	16.0	48.4

All races 3/, female

2003.....	*	0.2	0.3	0.5	0.8	1.2	2.3	5.8	10.1	37.6
2002.....	*	*	0.4	0.4	0.7	1.6	2.4	6.0	13.3	43.5
2001.....	*	0.2	0.4	0.4	0.7	1.3	2.7	6.1	12.6	46.1
2000.....	*	0.2	0.4	0.4	0.9	2.1	2.8	6.9	17.0	52.8
1999.....	0.2	0.2	0.4	0.5	0.7	1.5	2.6	6.2	15.8	56.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional deficiencies (E40-E64)

White, both sexes

2003.....	*	0.2	0.2	0.4	0.6	1.2	2.3	5.6	9.8	34.2
2002.....	0.1	0.2	0.3	0.4	0.7	1.4	2.5	5.9	12.3	40.2
2001.....	*	0.2	0.3	0.4	0.8	1.2	2.6	5.5	12.0	42.2
2000.....	*	0.2	0.2	0.5	0.9	2.0	2.7	6.4	15.4	48.3
1999.....	0.2	0.2	0.3	0.4	0.8	1.4	2.7	5.9	14.8	52.8

White, male

2003.....	*	*	*	0.4	0.6	1.3	2.3	5.2	10.1	29.6
2002.....	*	*	0.3	0.4	0.7	1.4	2.8	6.1	11.0	36.3
2001.....	*	*	*	0.4	0.7	1.1	2.6	5.3	12.2	33.8
2000.....	*	0.2	*	0.5	0.8	1.8	3.0	6.1	13.7	40.9
1999.....	*	*	0.3	0.4	1.0	1.3	3.0	6.1	14.7	46.5

White, female

2003.....	*	*	*	0.5	0.7	1.1	2.3	5.9	9.7	36.3
2002.....	*	*	0.4	0.4	0.7	1.3	2.2	5.7	13.1	41.9
2001.....	*	0.2	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.7
2000.....	*	*	0.3	0.4	0.9	2.0	2.6	6.6	16.5	51.5
1999.....	*	*	0.3	0.4	0.6	1.4	2.4	5.9	14.9	55.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2003.....	*	*	1.0	*	1.9	*	4.6	9.0	19.5	55.1
2002.....	*	*	*	1.5	2.0	3.5	5.1	10.5	20.7	60.6
2001.....	*	*	1.0	*	*	3.0	4.4	10.8	23.8	54.7
2000.....	*	*	*	*	2.1	3.1	6.5	12.1	23.6	67.1
1999.....	*	*	*	2.0	2.0	3.2	8.3	12.3	30.9	70.8

Black, male

2003.....	*	*	*	*	*	*	7.2	13.7	24.4	48.0
2002.....	*	*	*	*	*	*	*	9.6	27.5	47.2
2001.....	*	*	*	*	*	*	6.7	11.6	24.9	57.9
2000.....	*	*	*	*	*	*	9.2	14.6	22.4	58.5
1999.....	*	*	*	*	*	*	12.9	16.1	34.9	75.0

Black, female

2003.....	*	*	*	*	*	*	*	6.2	17.0	57.9
2002.....	*	*	*	*	*	*	4.7	11.0	17.1	65.7
2001.....	*	*	*	*	*	*	*	10.3	23.3	53.5
2000.....	*	*	*	*	*	*	4.5	10.6	24.2	70.7
1999.....	*	*	*	*	*	*	5.2	10.0	29.0	69.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

All races 3/, both sexes

2003.....	0.1	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.7	45.7
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0

All races 3/, male

2003.....	*	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.3	12.9	38.6
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4

All races 3/, female

2003.....	*	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.6	15.8	48.8
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

White, both sexes

2003.....	*	0.1	0.2	0.4	0.6	1.1	2.2	5.4	9.3	32.1
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	44.7
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9

White, male

2003.....	*	*	*	0.4	0.5	1.2	2.2	4.9	9.5	27.2
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.7	2.8	5.8	12.3	37.9
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2

White, female

2003.....	*	*	*	0.5	0.7	1.0	2.2	5.7	9.3	34.2
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.2	6.3	15.4	47.6
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

Black, both sexes

2003.....	*	*	1.0	*	1.8	*	4.4	8.8	19.3	53.4
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.0	6.5	11.2	21.8	64.0
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0

Black, male

2003.....	*	*	*	*	*	*	6.9	13.2	23.6	45.9
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.2	13.7	22.4	56.4
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8

Black, female

2003.....	*	*	*	*	*	*	*	6.2	17.0	56.3
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.5	9.7	21.5	67.1
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	0.4	0.6	3.1
2001.....	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	0.3	0.4	1.2	3.6
1999.....	*	*	*	*	*	*	0.4	1.1	3.9

All races 3/, male

2003.....	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	1.3	2.9
1999.....	*	*	*	*	*	*	*	1.3	3.9

All races 3/, female

2003.....	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	0.6	3.6
2001.....	*	*	*	*	*	*	*	1.0	3.1
2000.....	*	*	*	*	*	*	*	1.2	4.0
1999.....	*	*	*	*	*	*	*	1.1	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	0.3	0.3	1.2	3.6
1999.....	*	*	*	*	*	*	0.4	1.1	3.9

White, male

2003.....	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	1.4	3.0
1999.....	*	*	*	*	*	*	*	1.3	4.3

White, female

2003.....	*	*	*	*	*	*	*	*	2.1
2002.....	*	*	*	*	*	*	*	0.7	3.6
2001.....	*	*	*	*	*	*	*	0.9	3.1
2000.....	*	*	*	*	*	*	*	1.1	3.9
1999.....	*	*	*	*	*	*	*	1.0	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Meningitis (G00,G03)

All races 3/, both sexes

2003.....	0.2	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5

All races 3/, male

2003.....	*	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.6	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7

All races 3/, female

2003.....	*	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Meningitis (G00,G03)

White, both sexes

2003.....	0.1	0.2	0.2	0.3	0.3	0.5	0.4	0.7	0.6	1.0
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3

White, male

2003.....	*	0.3	0.3	0.3	0.4	0.6	*	0.9	*	*
2002.....	*	0.3	*	0.3	*	0.6	0.8	*	*	*
2001.....	*	0.2	0.3	0.4	0.5	*	0.7	0.8	*	*
2000.....	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
1999.....	*	0.3	*	0.4	*	0.5	0.7	*	1.3	*

White, female

2003.....	*	*	*	*	*	0.5	*	0.6	*	0.9
2002.....	*	*	*	*	*	*	0.5	0.5	*	1.2
2001.....	*	*	*	*	0.4	*	*	*	0.8	0.8
2000.....	*	*	*	*	*	*	*	0.6	1.1	1.0
1999.....	*	*	*	*	0.5	*	0.6	0.8	1.0	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

All races 3/, both sexes

2003.....	*	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.8	6.6	17.0	44.8	88.3	130.1
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4

All races 3/, male

2003.....	*	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.9	2.3	9.2	24.3	66.2	133.0	210.9
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6

All races 3/, female

2003.....	*	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.5	11.1	29.5	60.9	95.3
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

White, both sexes

2003.....	*	*	0.2	0.7	2.0	6.8	21.2	52.4	99.0	146.0
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.7	1.9	7.2	18.1	47.5	93.1	138.0
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2

White, male

2003.....	*	*	*	0.8	2.7	9.4	30.9	78.4	148.9	233.3
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.8	25.8	69.5	139.4	224.5
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7

White, female

2003.....	*	*	*	0.6	1.4	4.5	13.3	33.6	68.2	107.4
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.9	11.9	31.6	64.6	101.0
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

Black, both sexes

2003.....	*	*	*	*	1.9	4.1	10.7	22.8	38.3	52.8
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.0	8.7	21.0	39.2	51.7
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0

Black, male

2003.....	*	*	*	*	*	5.4	16.8	37.0	59.1	97.0
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.1	36.4	64.8	79.8
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7

Black, female

2003.....	*	*	*	*	*	*	6.6	14.4	27.5	35.6
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.7	11.1	25.1	40.0
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

All races 3/, both sexes

2003.....	*	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.3	9.2	29.3	85.3	223.0	659.0
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3

All races 3/, male

2003.....	*	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	3.0	9.7	29.3	78.5	203.7	495.4
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7

All races 3/, female

2003.....	*	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.8	29.3	90.1	234.9	729.5
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

White, both sexes

2003.....	*	*	0.3	1.1	3.6	10.3	34.5	101.8	264.4	834.0
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.2	3.5	9.8	30.4	88.1	230.5	687.6
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9

White, male

2003.....	*	*	0.3	1.1	3.0	10.3	33.1	94.9	238.5	630.3
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.3	30.3	81.2	211.2	517.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9

White, female

2003.....	*	*	0.4	1.0	4.2	10.3	35.7	106.8	280.3	924.3
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.8	9.3	30.4	93.2	242.4	760.5
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

Black, both sexes

2003.....	*	*	*	*	1.8	10.5	30.2	91.4	207.9	583.8
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.1	25.7	71.3	179.1	446.3
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4

Black, male

2003.....	*	*	*	*	*	11.1	32.5	93.1	206.0	459.8
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.5	26.4	62.4	153.6	342.7
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5

Black, female

2003.....	*	*	*	*	*	10.1	28.6	90.3	208.9	631.9
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	6.9	25.1	77.1	193.2	489.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

2003.....	50.3	87.7	142.8	224.5	363.7	574.0	946.1	1,627.3	2,918.8	7,138.0
2002.....	49.7	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001.....	48.8	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000.....	49.2	89.4	149.5	251.3	413.0	655.3	1,080.0	1,825.5	3,282.3	7,925.3
1999.....	50.0	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1

All races 3/, male

2003.....	67.7	123.1	204.5	314.7	497.9	765.9	1,225.3	2,051.8	3,517.1	7,329.3
2002.....	67.9	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001.....	66.5	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000.....	67.5	128.3	214.3	356.8	567.6	870.8	1,415.0	2,264.4	3,896.4	8,015.6
1999.....	68.9	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2

All races 3/, female

2003.....	33.2	53.3	83.7	139.7	241.8	407.6	722.2	1,325.1	2,552.9	7,053.5
2002.....	31.6	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001.....	31.5	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000.....	31.3	51.8	88.1	153.7	274.4	472.1	813.7	1,510.0	2,905.8	7,886.5
1999.....	31.4	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2003.....	43.4	76.1	124.8	203.3	333.1	537.6	907.0	1,593.5	2,909.9	7,233.1
2002.....	43.6	76.6	127.4	211.8	344.7	567.9	957.1	1,663.0	3,023.3	7,482.7
2001.....	41.6	74.2	127.7	212.4	355.6	580.3	980.2	1,721.1	3,122.8	7,770.6
2000.....	42.1	77.0	132.3	227.4	380.3	621.8	1,041.6	1,783.6	3,264.0	8,043.5
1999.....	42.4	76.6	132.4	230.2	385.3	644.7	1,068.3	1,848.2	3,345.3	8,299.2

White, male

2003.....	60.6	110.4	183.6	289.9	463.8	727.3	1,183.5	2,022.8	3,520.9	7,466.8
2002.....	62.0	111.9	186.7	301.4	479.1	765.2	1,253.1	2,112.9	3,642.0	7,766.6
2001.....	59.1	109.0	185.6	301.7	493.8	781.8	1,286.2	2,192.9	3,777.7	8,060.7
2000.....	59.9	115.3	193.8	327.4	530.6	838.4	1,381.3	2,233.1	3,904.3	8,188.3
1999.....	60.8	113.0	193.5	329.9	533.4	874.1	1,420.2	2,361.9	4,089.2	8,717.5

White, female

2003.....	26.2	42.1	67.2	120.1	212.2	370.0	680.7	1,283.1	2,533.3	7,129.6
2002.....	25.2	41.6	69.3	126.1	220.5	394.1	715.8	1,341.0	2,644.0	7,359.3
2001.....	24.0	39.8	71.0	127.1	228.1	403.3	732.3	1,386.9	2,726.9	7,649.2
2000.....	24.2	39.1	72.6	133.0	242.4	433.9	767.5	1,457.5	2,869.7	7,981.6
1999.....	24.0	40.6	72.8	135.3	248.8	444.5	788.7	1,487.4	2,906.2	8,131.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2003.....	104.2	183.8	305.8	451.7	695.7	983.6	1,489.0	2,249.7	3,405.2	6,862.0
2002.....	99.1	184.1	302.3	468.4	697.7	1,001.3	1,510.2	2,343.9	3,555.3	7,081.7
2001.....	105.7	181.9	307.2	471.1	713.7	1,057.5	1,548.8	2,422.8	3,623.2	7,316.7
2000.....	104.8	195.1	314.0	496.2	745.3	1,021.2	1,608.8	2,505.4	3,881.4	7,538.2
1999.....	109.8	188.5	311.0	509.4	743.3	1,093.2	1,704.6	2,553.3	3,787.7	7,582.5

Black, male

2003.....	129.8	239.7	412.9	610.0	924.3	1,273.7	1,900.2	2,789.8	4,035.4	6,547.7
2002.....	122.7	237.2	406.5	633.2	934.4	1,292.7	1,908.8	2,803.5	4,093.1	6,929.0
2001.....	129.8	237.2	405.7	627.8	934.4	1,342.7	1,988.0	2,943.9	4,284.2	7,363.9
2000.....	131.5	254.2	431.4	700.4	998.4	1,272.1	1,945.0	2,889.9	4,331.1	7,093.9
1999.....	138.3	246.5	423.5	686.4	993.1	1,400.0	2,146.5	3,046.5	4,407.7	7,921.1

Black, female

2003.....	81.7	135.2	215.1	321.5	517.3	768.3	1,211.6	1,928.7	3,075.0	6,984.1
2002.....	78.2	138.0	213.9	333.0	512.5	785.0	1,242.7	2,069.3	3,275.3	7,140.2
2001.....	84.3	134.0	223.4	342.6	540.2	846.9	1,256.2	2,111.8	3,286.5	7,299.2
2000.....	81.1	145.0	218.2	337.4	556.8	834.7	1,370.4	2,257.7	3,634.0	7,723.8
1999.....	84.7	138.5	215.6	364.7	547.4	865.8	1,411.4	2,256.2	3,480.4	7,456.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes

2003.....	40.5	71.3	116.6	185.0	296.1	456.3	730.9	1,218.3	2,151.6	5,278.4
2002.....	40.2	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001.....	39.1	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000.....	39.1	71.9	122.6	207.9	335.1	521.8	837.6	1,373.1	2,413.1	5,848.7
1999.....	40.1	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0

All races 3/, male

2003.....	56.9	104.0	173.3	268.1	416.1	624.7	975.4	1,589.0	2,695.8	5,621.5
2002.....	57.7	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001.....	55.8	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000.....	56.5	108.4	182.6	304.3	472.8	713.1	1,128.6	1,755.5	2,966.4	6,100.8
1999.....	58.1	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6

All races 3/, female

2003.....	24.3	39.5	62.4	106.7	187.0	310.3	534.8	954.5	1,818.9	5,126.7
2002.....	22.9	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001.....	22.6	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000.....	22.0	36.7	65.9	118.8	211.6	359.0	606.3	1,098.1	2,073.9	5,740.1
1999.....	22.4	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

2003.....	35.9	63.6	104.7	170.9	275.7	432.1	706.5	1,199.6	2,153.0	5,363.5
2002.....	36.3	64.0	108.1	177.9	284.5	457.0	744.1	1,255.7	2,234.0	5,529.1
2001.....	34.2	61.8	108.6	178.5	294.7	468.4	765.0	1,297.7	2,308.3	5,752.1
2000.....	34.5	64.0	111.3	191.5	312.8	499.8	812.2	1,347.1	2,408.1	5,953.9
1999.....	34.9	64.1	112.7	193.7	321.3	524.0	839.7	1,403.0	2,484.1	6,152.7

White, male

2003.....	52.1	95.4	159.4	251.9	394.1	599.0	949.5	1,575.7	2,710.0	5,747.2
2002.....	53.9	96.8	164.4	261.0	406.6	631.3	1,006.4	1,654.3	2,805.0	5,939.8
2001.....	50.9	94.4	163.4	261.7	420.5	647.2	1,031.2	1,716.7	2,896.4	6,157.2
2000.....	51.6	100.1	168.9	284.0	447.8	693.1	1,108.8	1,739.7	2,984.7	6,257.6
1999.....	52.4	98.8	170.3	286.1	454.8	730.0	1,148.4	1,851.7	3,144.4	6,669.0

White, female

2003.....	19.6	32.1	51.1	93.1	166.1	284.6	507.6	927.7	1,809.7	5,193.6
2002.....	18.6	31.4	53.1	98.3	171.5	303.6	530.3	970.4	1,884.1	5,350.6
2001.....	17.5	29.4	54.9	99.1	178.6	311.5	549.5	1,001.0	1,952.9	5,582.5
2000.....	17.3	28.4	55.4	104.2	189.0	332.2	573.0	1,062.4	2,053.1	5,824.0
1999.....	17.4	29.7	56.4	105.8	198.3	344.3	594.5	1,088.0	2,094.3	5,945.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2003.....	79.0	139.6	232.4	348.4	533.8	752.7	1,115.2	1,638.3	2,445.2	4,941.2
2002.....	74.8	138.7	232.1	365.8	542.7	764.2	1,126.8	1,715.1	2,557.1	5,115.2
2001.....	79.5	137.2	232.2	369.7	546.5	809.7	1,160.0	1,780.6	2,619.7	5,275.7
2000.....	77.5	144.5	239.3	386.1	577.5	782.7	1,219.5	1,843.0	2,789.8	5,403.4
1999.....	82.9	143.0	240.2	400.6	579.9	852.1	1,289.1	1,878.6	2,749.2	5,487.4

Black, male

2003.....	102.7	188.1	322.4	481.2	718.0	997.2	1,460.0	2,083.2	2,981.9	4,850.3
2002.....	96.7	186.2	319.6	503.6	740.6	1,001.5	1,448.8	2,095.3	3,039.3	5,125.7
2001.....	101.9	184.4	314.7	503.1	731.0	1,039.8	1,518.8	2,214.9	3,181.5	5,459.9
2000.....	101.1	196.1	338.2	556.5	788.9	990.2	1,497.3	2,163.3	3,177.1	5,198.6
1999.....	109.3	193.3	335.5	556.7	789.2	1,111.8	1,652.9	2,277.2	3,259.4	5,858.8

Black, female

2003.....	58.2	97.4	156.3	239.2	390.1	571.3	882.6	1,373.9	2,164.0	4,976.5
2002.....	55.5	97.5	157.9	252.6	388.0	588.1	910.7	1,488.0	2,306.1	5,111.2
2001.....	59.7	96.3	162.0	260.3	401.4	639.8	921.0	1,521.3	2,333.5	5,207.3
2000.....	56.5	100.7	158.5	253.5	420.1	628.5	1,022.5	1,636.6	2,576.7	5,489.0
1999.....	59.6	99.6	159.4	273.1	415.8	659.7	1,047.8	1,638.6	2,496.4	5,348.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes

2003.....	0.3	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.5	5.6	8.1	11.7	16.6
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0

All races 3/, male

2003.....	0.2	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.7	1.1	1.9	2.3	3.6	5.7	7.7	11.6
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3

All races 3/, female

2003.....	0.3	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.2	9.8	14.2	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, both sexes

2003.....	0.2	0.4	0.5	0.8	1.6	3.1	4.7	8.6	12.4	21.8
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.6	5.7	8.6	12.4	17.4
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9

White, male

2003.....	*	0.3	0.3	0.7	1.3	2.3	3.7	6.9	9.8	17.3
2002.....	*	0.3	0.5	0.9	1.5	1.8	4.4	5.9	8.8	12.6
2001.....	0.3	0.4	0.4	1.0	1.2	2.3	3.3	6.3	8.0	13.3
2000.....	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.0	8.3	12.1
1999.....	0.2	0.4	0.6	1.0	1.3	2.6	4.9	5.5	9.4	11.1

White, female

2003.....	0.2	0.5	0.6	1.0	1.9	3.8	5.5	9.8	14.0	23.8
2002.....	0.2	0.5	0.9	1.4	2.5	3.7	5.9	10.6	14.2	21.6
2001.....	0.3	0.4	0.7	1.1	2.3	3.9	6.4	9.9	14.7	21.0
2000.....	0.4	0.6	0.8	1.5	2.4	4.7	7.5	10.5	15.0	19.7
1999.....	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.4	22.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2003.....	*	1.2	1.4	2.2	3.2	3.1	4.8	3.9	*	7.6
2002.....	*	0.8	1.1	2.0	2.9	3.2	*	4.8	*	*
2001.....	0.9	1.1	1.6	2.2	2.0	3.3	5.1	4.9	*	6.8
2000.....	0.9	1.0	1.1	2.0	2.2	3.3	3.6	3.7	*	8.5
1999.....	0.7	0.9	1.6	2.2	2.1	2.8	4.2	4.3	*	8.3

Black, male

2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2003.....	*	1.6	1.7	*	3.3	*	6.2	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.3	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

All races 3/, both sexes

2003.....	3.9	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.8	6.7	9.4	13.3	16.7	23.3	36.3	66.7	190.6
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9

All races 3/, male

2003.....	5.2	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.4	9.7	12.3	17.8	21.0	27.4	36.6	60.9	132.0
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6

All races 3/, female

2003.....	2.6	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.4	13.0	20.0	36.2	70.2	215.8
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

White, both sexes

2003.....	2.7	4.4	6.6	8.5	11.4	14.7	20.2	33.2	61.2	185.3
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.4	6.3	9.5	12.7	17.9	30.2	58.7	179.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3

White, male

2003.....	4.0	6.4	9.7	12.2	15.0	19.4	23.6	36.1	57.9	131.9
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.6	8.7	12.9	16.3	21.7	30.6	52.6	123.4
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7

White, female

2003.....	1.3	2.4	3.5	5.0	8.1	10.7	17.4	31.1	63.3	209.0
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.7	2.2	4.0	6.3	9.5	14.7	30.0	62.5	203.9
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

Black, both sexes

2003.....	12.8	23.9	32.8	41.9	48.3	63.0	77.1	113.3	145.7	332.3
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	18.1	26.3	38.1	48.4	52.5	80.9	111.5	172.1	340.9
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1

Black, male

2003.....	14.9	29.8	41.5	55.1	59.8	86.7	99.9	129.6	149.1	287.8
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.6	22.0	37.9	50.0	66.7	67.0	91.5	112.9	176.0	252.2
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6

Black, female

2003.....	11.0	18.7	25.5	31.0	39.4	45.5	61.8	103.6	143.9	349.6
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.7	16.8	28.9	34.8	41.7	73.5	110.6	169.9	378.0
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Hypertensive heart and renal disease (I13)

All races 3/, both sexes

2003.....	0.3	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.5	0.8	1.0	1.6	3.1	5.1	9.9	24.0
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1

All races 3/, male

2003.....	0.3	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	4.9	10.6	26.4
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3

All races 3/, female

2003.....	0.2	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.8	5.2	9.5	23.0
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart and renal disease (I13)

White, both sexes

2003.....	*	0.1	0.2	0.3	0.6	0.9	2.1	4.2	8.3	23.4
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.1	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	21.8
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1

White, male

2003.....	*	*	*	0.5	0.8	1.1	2.2	4.9	9.3	24.3
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.6	4.3	9.0	23.2
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8

White, female

2003.....	*	*	*	*	0.5	0.7	1.9	3.7	7.8	22.9
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.5	21.2
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart and renal disease (I13)

Black, both sexes

2003.....	1.5	2.3	3.8	5.3	5.5	7.4	11.0	16.5	24.1	53.1
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.4	8.0	11.8	17.7	32.4	50.5
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4

Black, male

2003.....	2.1	3.4	4.6	7.6	6.4	6.6	11.8	15.5	24.4	50.0
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.8	3.1	5.3	5.8	9.0	13.1	13.2	33.2	58.5
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8

Black, female

2003.....	*	*	3.1	3.5	4.8	8.0	10.4	17.1	24.0	54.3
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.0	7.2	10.8	20.5	32.0	47.1
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2003.....	23.5	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.4	48.4	88.4	155.7	253.4	395.3	636.3	1,033.1	1,777.2	4,127.8
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1

All races 3/, male

2003.....	35.5	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	36.7	77.8	137.5	237.0	369.9	559.2	886.6	1,369.4	2,269.9	4,461.0
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5

All races 3/, female

2003.....	11.6	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.2	20.1	41.8	80.5	148.9	255.9	437.3	791.3	1,475.1	3,984.3
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

White, both sexes

2003.....	22.4	43.5	76.1	126.4	208.2	322.6	526.3	877.2	1,542.2	3,671.7
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.0	45.5	83.5	147.4	241.1	384.0	623.6	1,019.8	1,779.6	4,210.4
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2

White, male

2003.....	34.4	68.8	121.6	194.2	308.7	464.4	731.5	1,195.8	2,028.8	4,106.1
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.5	75.0	132.2	226.3	356.7	551.5	879.5	1,365.5	2,293.4	4,591.4
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9

White, female

2003.....	10.4	18.4	31.5	61.3	115.2	197.3	358.4	646.9	1,242.2	3,479.3
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.4	16.5	36.3	72.8	135.0	238.6	417.1	769.0	1,463.3	4,047.5
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

Black, both sexes

2003.....	35.2	71.5	133.3	213.8	352.2	503.0	756.8	1,139.8	1,696.1	3,310.1
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.0	79.4	145.2	254.5	397.4	544.4	858.8	1,301.6	1,964.6	3,694.1
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8

Black, male

2003.....	50.2	103.2	195.6	302.7	490.5	677.1	1,027.8	1,501.3	2,138.8	3,285.3
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.7	114.7	209.5	382.9	554.3	696.8	1,086.3	1,571.1	2,281.4	3,585.3
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1

Black, female

2003.....	21.9	43.9	80.5	140.7	244.3	373.7	574.0	924.9	1,464.2	3,319.7
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.5	92.7	154.6	280.6	431.1	697.5	1,128.1	1,790.3	3,739.6
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2003.....	9.6	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.3	38.5	67.8	112.4	168.7	258.1	405.6	655.8	1,321.0
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3

All races 3/, male

2003.....	14.4	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.4	34.0	58.9	101.5	161.7	234.9	352.4	525.6	823.1	1,488.0
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1

All races 3/, female

2003.....	5.0	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	9.0	19.2	36.7	68.2	112.3	183.1	319.3	553.3	1,249.0
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

White, both sexes

2003.....	9.3	18.2	32.1	54.4	89.1	132.3	205.3	324.2	532.9	1,132.6
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.4	20.7	37.2	66.2	110.1	166.5	255.3	401.7	656.7	1,338.4
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2

White, male

2003.....	14.1	28.7	51.2	83.0	130.8	188.3	280.0	430.3	685.7	1,289.3
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.1	33.9	58.2	99.9	160.3	235.5	353.1	526.8	830.9	1,519.3
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0

White, female

2003.....	4.5	7.7	13.4	26.8	50.5	82.8	144.1	247.5	438.7	1,063.2
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.6	7.7	16.8	34.2	64.0	106.7	176.5	311.0	549.5	1,261.1
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2003.....	14.4	27.5	49.5	79.4	124.3	180.3	270.2	401.4	597.0	1,094.9
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.0	30.1	57.4	96.1	153.2	210.6	325.9	492.4	729.2	1,278.9
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8

Black, male

2003.....	20.0	38.6	70.9	108.1	167.9	230.2	353.4	516.2	736.1	1,113.5
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	42.1	77.1	137.2	205.6	259.4	399.6	570.9	843.5	1,309.0
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7

Black, female

2003.....	9.5	18.0	31.5	55.7	90.3	143.3	214.0	333.2	524.2	1,087.7
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.9	41.4	64.1	114.1	174.2	273.7	441.9	666.3	1,266.4
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

All races 3/, both sexes

2003.....	0.3	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.5	21.6
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9

All races 3/, male

2003.....	0.5	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.7	2.5	3.5	4.5	5.3	7.5	11.5	19.6
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3

All races 3/, female

2003.....	*	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	2.0	3.0	4.8	8.3	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

White, both sexes

2003.....	0.3	0.4	0.8	1.1	1.6	2.6	3.4	5.0	8.1	20.2
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.3	2.9	3.8	5.6	9.4	21.9
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4

White, male

2003.....	0.4	0.6	1.2	1.5	2.2	3.6	4.8	6.7	8.9	19.7
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.1	5.0	7.1	11.4	20.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9

White, female

2003.....	*	0.2	0.4	0.7	0.9	1.6	2.4	3.8	7.7	20.5
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.4	1.9	2.9	4.5	8.1	22.7
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

Black, both sexes

2003.....	0.7	1.1	1.5	2.7	3.3	4.9	6.2	8.7	11.7	19.8
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.2	3.8	4.8	5.0	6.1	10.0	12.7	22.0
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2

Black, male

2003.....	*	1.7	2.2	4.6	5.0	7.4	7.9	14.2	19.2	24.0
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.8	6.0	6.9	8.0	8.6	12.7	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*

Black, female

2003.....	*	*	*	*	*	*	5.1	5.4	7.7	18.2
2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.2	*	*	8.2	11.4	23.1
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2003.....	13.5	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.2	26.7	48.8	86.2	138.5	223.5	374.2	621.6	1,111.9	2,785.2
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0

All races 3/, male

2003.....	20.7	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	20.9	43.0	76.9	132.9	204.7	319.8	528.9	836.3	1,435.4	2,953.4
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1

All races 3/, female

2003.....	6.5	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.7	11.0	22.1	43.0	79.2	141.6	251.3	467.3	913.5	2,712.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2003.....	12.9	24.9	43.2	71.0	117.6	187.7	317.6	548.0	1,001.1	2,518.8
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.4	24.4	45.4	79.8	128.7	214.6	364.5	612.4	1,113.5	2,850.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6

White, male

2003.....	19.9	39.5	69.2	109.6	175.8	272.6	446.7	758.8	1,334.2	2,797.1
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.0	40.4	72.6	124.2	193.1	312.0	521.5	831.5	1,451.1	3,052.1
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1

White, female

2003.....	5.8	10.5	17.7	33.8	63.8	112.8	211.9	395.6	795.8	2,395.5
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.6	8.6	19.0	38.0	69.6	130.1	237.8	453.5	905.7	2,763.7
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, both sexes

2003.....	20.1	42.8	82.2	131.7	224.6	317.7	480.4	729.7	1,087.4	2,195.3
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.4	48.4	85.5	154.6	239.4	328.8	526.8	799.2	1,222.7	2,393.2
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8

Black, male

2003.....	29.4	62.9	122.5	189.9	317.6	439.5	666.5	970.9	1,383.5	2,147.8
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.2	71.2	128.6	239.7	341.8	429.4	678.1	987.4	1,422.9	2,257.2
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5

Black, female

2003.....	11.9	25.3	48.2	83.8	152.1	227.3	354.9	586.3	932.3	2,213.8
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	29.0	50.4	88.4	163.2	254.1	419.5	678.0	1,112.5	2,450.1
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2003.....	4.7	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.5	17.9	28.9	41.3	56.3	83.0	125.1	209.6	527.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4

All races 3/, male

2003.....	7.4	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.0	15.7	29.1	46.6	62.4	82.1	116.5	164.1	249.0	522.6
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6

All races 3/, female

2003.....	2.0	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.7	3.6	7.2	12.5	22.3	34.4	56.4	97.1	185.4	529.8
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, both sexes

2003.....	4.4	9.0	16.7	24.2	35.4	47.5	66.6	103.0	179.6	439.3
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.4	15.9	25.0	35.5	50.2	74.7	116.0	199.8	523.6
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5

White, male

2003.....	6.9	14.8	27.7	38.3	53.4	69.2	92.5	136.9	224.4	458.7
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.1	26.3	40.5	54.9	74.2	105.5	151.9	238.0	520.4
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2

White, female

2003.....	1.8	3.2	5.9	10.6	18.7	28.3	45.4	78.5	152.0	430.7
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.3	17.7	29.4	49.8	89.9	176.2	525.0
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Black, both sexes

2003.....	7.3	17.4	35.9	53.2	91.8	111.4	149.2	214.1	282.8	543.2
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.8	37.7	66.3	95.1	115.1	175.2	240.6	354.7	648.3
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6

Black, male

2003.....	11.4	26.4	57.8	81.3	140.0	161.3	228.9	301.3	380.2	583.9
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	31.1	60.2	111.1	143.3	161.5	245.0	326.0	419.2	647.0
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9

Black, female

2003.....	3.7	9.6	17.3	30.0	54.2	74.4	95.5	162.2	231.7	527.4
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.1	19.3	31.6	59.1	80.5	125.7	185.6	319.2	648.8
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2003.....	8.9	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	8.9	17.2	30.9	57.3	97.2	167.2	291.2	496.5	902.3	2,257.6
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6

All races 3/, male

2003.....	13.3	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	13.9	27.3	47.8	86.4	142.2	237.7	412.4	672.2	1,186.4	2,430.8
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4

All races 3/, female

2003.....	4.5	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.4	14.8	30.5	56.9	107.2	194.9	370.2	728.1	2,183.1
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2003.....	8.5	16.0	26.5	46.8	82.2	140.3	251.0	445.0	821.5	2,079.5
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.4	16.0	29.4	54.9	93.2	164.4	289.8	496.5	913.8	2,326.5
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1

White, male

2003.....	12.9	24.7	41.5	71.3	122.4	203.4	354.1	622.0	1,109.8	2,338.4
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.5	26.3	46.2	83.7	138.2	237.7	416.0	679.6	1,213.1	2,531.7
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9

White, female

2003.....	4.0	7.3	11.8	23.3	45.1	84.5	166.6	317.1	643.8	1,964.9
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.1	27.7	51.9	100.7	188.0	363.6	729.4	2,238.8
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2003.....	12.8	25.4	46.4	78.5	132.8	206.3	331.2	515.6	804.7	1,652.1
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.0	28.6	47.9	88.2	144.4	213.8	351.6	558.6	868.0	1,745.0
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2

Black, male

2003.....	18.0	36.5	64.7	108.6	177.6	278.2	437.5	669.6	1,003.3	1,563.9
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.2	40.1	68.4	128.6	198.5	267.9	433.0	661.5	1,003.7	1,610.1
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5

Black, female

2003.....	8.3	15.7	30.9	53.8	97.9	153.0	259.4	424.0	700.6	1,686.4
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.9	31.1	56.8	104.1	173.5	293.8	492.4	793.3	1,801.3
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

All races 3/, both sexes

2003.....	12.6	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.2	17.8	26.4	40.7	65.1	104.8	169.3	290.4	547.6	1,489.7
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9

All races 3/, male

2003.....	15.6	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.3	23.3	34.2	52.9	82.3	128.9	207.6	338.9	617.2	1,469.8
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9

All races 3/, female

2003.....	9.5	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.1	12.4	18.9	29.4	49.8	84.2	139.0	255.6	504.8	1,498.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

White, both sexes

2003.....	10.5	15.2	21.4	34.8	53.8	90.7	153.2	276.4	528.9	1,461.4
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.1	14.7	22.4	36.2	59.6	98.7	162.8	284.5	549.2	1,524.5
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3

White, male

2003.....	13.5	19.7	27.5	44.5	68.3	111.8	188.4	332.0	604.3	1,467.6
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	12.9	19.8	29.1	47.4	75.8	122.0	201.4	333.4	621.3	1,507.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5

White, female

2003.....	7.5	10.8	15.4	25.5	40.4	72.1	124.3	236.1	482.4	1,458.6
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.3	9.6	15.9	25.5	44.7	78.6	131.7	249.1	504.8	1,531.7
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

Black, both sexes

2003.....	28.9	40.8	61.1	85.3	124.6	176.1	265.5	364.8	574.9	1,238.1
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	44.0	64.1	87.4	124.2	174.5	264.5	408.4	616.4	1,309.4
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8

Black, male

2003.....	34.7	51.0	79.5	113.5	158.3	223.6	318.0	434.1	668.1	1,220.9
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	55.8	87.1	117.1	160.6	215.3	304.0	463.0	685.7	1,294.1
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1

Black, female

2003.....	23.8	31.9	45.4	62.1	98.2	140.9	230.2	323.7	526.1	1,244.8
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	34.1	45.4	64.3	97.0	144.2	236.4	373.3	578.2	1,315.8
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute and subacute endocarditis (I33)

All races 3/, both sexes

2003.....	0.3	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.4	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9

All races 3/, male

2003.....	0.4	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.4	1.6	3.1	4.0	3.9
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7

All races 3/, female

2003.....	0.2	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute and subacute endocarditis (I33)

White, both sexes

2003.....	0.3	0.3	0.4	0.5	0.8	1.0	1.4	2.3	2.8	3.1
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0

White, male

2003.....	0.3	0.4	0.5	0.6	1.0	1.2	1.5	2.8	3.9	4.4
2002.....	0.3	0.4	0.6	0.4	0.9	1.3	1.7	2.7	3.9	3.6
2001.....	0.3	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.3	3.9
2000.....	0.4	0.5	0.6	0.7	0.9	1.2	1.7	3.1	3.9	4.0
1999.....	0.3	0.4	0.5	0.8	0.7	1.2	1.6	3.0	4.0	5.1

White, female

2003.....	*	0.2	0.2	0.4	0.6	0.8	1.3	1.8	2.1	2.6
2002.....	*	*	*	0.3	0.7	0.7	1.2	1.8	2.3	2.3
2001.....	*	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000.....	*	*	0.3	0.4	0.5	1.0	1.1	1.6	2.0	2.1
1999.....	*	*	*	*	0.6	0.8	1.3	1.6	2.0	2.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2003.....	0.2	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.4	0.4	0.6	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0

All races 3/, male

2003.....	0.3	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.2	0.3	*	0.5	0.4	0.6	1.1	1.4	1.7	2.0
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5

All races 3/, female

2003.....	0.2	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.1	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, both sexes

2003.....	0.2	0.3	0.2	0.4	0.4	0.8	0.8	1.0	1.5	1.7
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0

White, male

2003.....	0.2	0.3	0.3	0.4	0.5	1.0	1.1	1.1	1.6	1.7
2002.....	0.3	0.3	0.3	0.4	0.6	0.6	1.1	1.1	1.2	2.5
2001.....	*	0.2	0.3	0.4	0.6	0.5	0.9	1.3	1.1	2.0
2000.....	0.2	*	*	0.5	*	*	1.1	1.2	1.8	2.0
1999.....	0.3	0.3	*	0.6	0.6	0.7	0.9	1.3	1.5	2.5

White, female

2003.....	*	*	*	0.4	*	0.7	0.6	0.8	1.4	1.7
2002.....	*	*	*	0.4	*	0.5	0.9	1.0	1.4	2.0
2001.....	*	0.3	*	0.3	*	*	0.9	1.2	1.4	1.9
2000.....	*	*	*	*	*	0.6	0.7	0.9	1.1	1.7
1999.....	*	0.3	*	*	0.5	0.7	0.8	1.1	1.4	1.8

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Heart failure (I50)

All races 3/, both sexes

2003.....	1.0	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.1	6.2	13.0	24.2	47.1	93.6	201.6	675.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9

All races 3/, male

2003.....	1.3	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	2.0	3.6	7.5	15.6	28.7	56.1	106.0	226.1	637.8
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0

All races 3/, female

2003.....	0.7	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.6	20.4	40.0	84.7	186.6	692.3
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Heart failure (I50)

White, both sexes

2003.....	0.7	1.3	2.6	5.6	10.7	22.4	43.2	89.5	195.3	660.4
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.6	5.5	11.9	22.9	45.5	92.8	203.0	695.9
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2

White, male

2003.....	0.9	1.7	3.0	7.0	13.1	27.0	52.1	105.9	217.1	650.8
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.1	6.9	14.2	27.7	54.9	105.8	228.3	658.2
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0

White, female

2003.....	0.4	0.9	2.1	4.2	8.5	18.3	35.9	77.6	181.9	664.7
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.1	4.1	9.7	18.8	38.0	83.5	187.5	712.1
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Heart failure (I50)

Black, both sexes

2003.....	3.3	4.7	9.4	15.3	27.7	41.1	70.9	114.7	203.6	531.8
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.9	14.2	24.6	39.6	71.9	120.2	218.6	559.8
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9

Black, male

2003.....	3.9	5.5	11.6	19.4	35.9	51.0	79.5	124.6	227.4	513.0
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.6	9.6	15.5	31.0	43.6	79.0	129.7	239.1	536.4
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5

Black, female

2003.....	2.8	4.0	7.6	11.9	21.4	33.8	65.1	108.8	191.1	539.2
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.5	13.2	19.8	36.7	66.9	114.1	207.3	569.6
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes

2003.....	11.0	15.5	21.3	31.7	46.3	71.8	113.2	186.0	328.5	781.5
2002.....	11.2	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001.....	10.5	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000.....	10.8	15.7	22.5	33.5	51.1	78.9	120.0	193.5	341.9	809.2
1999.....	11.1	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1

All races 3/, male

2003.....	13.8	19.7	27.5	40.8	58.6	88.8	139.6	224.8	380.7	795.9
2002.....	14.2	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001.....	13.4	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000.....	13.5	20.5	29.6	44.1	65.4	98.3	148.8	228.5	385.6	826.1
1999.....	13.9	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7

All races 3/, female

2003.....	8.4	11.5	15.4	23.1	35.1	57.0	92.0	158.4	296.6	775.1
2002.....	8.2	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001.....	7.6	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000.....	8.2	11.1	15.8	23.6	38.2	62.4	97.1	168.3	315.1	802.0
1999.....	8.3	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2003.....	9.4	13.4	18.2	28.4	42.0	66.5	107.8	183.7	329.3	796.1
2002.....	9.7	13.3	18.8	28.9	42.7	69.1	111.9	191.2	339.3	806.1
2001.....	8.7	12.9	18.9	28.3	44.0	70.3	112.3	186.3	333.4	809.9
2000.....	9.0	13.1	19.2	29.8	46.7	74.2	115.1	188.4	342.0	824.1
1999.....	9.2	12.9	19.4	29.8	48.2	74.9	118.6	192.7	346.9	839.0

White, male

2003.....	12.0	17.3	23.6	36.5	53.7	82.6	133.8	222.2	381.7	810.7
2002.....	12.6	17.7	24.8	36.9	54.7	85.0	138.8	229.4	391.0	834.2
2001.....	11.5	17.0	24.7	35.4	56.5	86.9	138.2	226.4	388.3	831.0
2000.....	11.4	17.6	25.2	39.3	60.3	92.6	143.8	223.3	387.3	843.3
1999.....	11.9	17.3	25.1	39.0	62.4	97.5	148.4	232.5	409.5	883.9

White, female

2003.....	6.8	9.5	12.9	20.5	31.1	52.3	86.5	155.9	297.0	789.7
2002.....	6.7	9.0	13.0	21.4	31.5	55.1	90.0	163.8	307.6	793.9
2001.....	5.8	8.8	13.4	21.5	32.5	55.8	91.3	158.0	300.1	801.1
2000.....	6.6	8.7	13.3	20.8	34.1	58.2	91.9	163.2	314.1	815.9
1999.....	6.5	8.6	13.7	21.0	35.2	55.1	94.9	164.7	310.0	821.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2003.....	24.5	34.4	49.8	67.9	94.2	132.5	190.7	246.1	366.8	703.7
2002.....	23.6	36.5	49.5	70.6	94.5	127.7	180.3	263.4	376.5	738.5
2001.....	25.1	36.9	50.0	73.6	91.5	135.2	179.9	276.1	383.5	730.2
2000.....	25.2	37.9	54.3	70.7	97.6	132.3	189.4	283.9	394.2	743.9
1999.....	26.3	38.5	51.8	77.4	109.1	140.1	206.0	283.3	400.8	746.1

Black, male

2003.....	29.7	43.0	65.2	90.9	120.3	170.1	234.2	304.9	436.3	707.9
2002.....	28.2	47.7	62.8	90.7	117.2	157.1	226.3	314.2	434.1	753.7
2001.....	30.0	45.9	60.5	95.7	121.2	158.8	227.2	347.1	479.1	805.6
2000.....	30.8	47.6	75.2	99.2	127.5	167.0	222.4	328.2	442.5	753.4
1999.....	31.8	47.7	67.7	103.6	142.8	183.3	265.3	341.0	503.1	849.5

Black, female

2003.....	19.9	26.8	36.8	48.9	73.9	104.5	161.3	211.1	330.4	702.0
2002.....	19.5	26.8	38.3	54.1	76.8	105.8	149.4	233.0	346.5	732.8
2001.....	20.8	29.1	41.0	55.4	68.2	117.7	148.4	233.8	334.8	702.2
2000.....	20.3	29.7	37.3	48.5	75.4	106.5	165.9	255.4	367.7	739.9
1999.....	21.5	30.6	38.3	56.1	82.7	108.1	166.8	248.6	350.0	707.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes

2003.....	1.1	1.8	3.3	5.0	8.0	13.1	21.3	38.4	69.9	188.9
2002.....	1.0	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001.....	1.0	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000.....	1.1	1.8	3.0	4.5	7.9	11.6	19.3	33.8	63.6	160.8
1999.....	0.9	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1

All races 3/, male

2003.....	1.3	2.5	4.2	6.0	9.2	14.9	22.5	37.5	66.1	153.5
2002.....	1.3	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001.....	1.3	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000.....	1.4	2.3	3.9	5.5	9.2	11.6	20.8	33.8	62.0	135.1
1999.....	1.1	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4

All races 3/, female

2003.....	0.9	1.2	2.4	4.0	6.9	11.4	20.4	39.1	72.2	204.6
2002.....	0.7	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001.....	0.7	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000.....	0.8	1.3	2.2	3.5	6.7	11.6	18.2	33.8	64.5	171.9
1999.....	0.7	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

White, both sexes

2003.....	0.7	1.2	2.3	3.3	5.8	9.8	16.6	33.0	64.1	181.0
2002.....	0.6	1.0	1.9	3.0	5.4	9.0	16.8	29.8	60.4	170.4
2001.....	0.5	1.1	1.8	2.9	5.3	9.2	15.3	30.9	58.6	163.3
2000.....	0.6	1.0	1.8	3.0	5.5	8.7	15.5	28.9	57.8	151.5
1999.....	0.5	1.0	1.6	2.7	5.0	9.1	15.2	27.6	53.3	142.0

White, male

2003.....	0.8	1.6	2.9	4.1	6.6	11.6	18.0	32.7	60.3	145.3
2002.....	0.8	1.4	2.4	3.8	6.3	9.8	18.3	30.8	57.7	145.2
2001.....	0.7	1.4	2.2	3.5	5.9	10.3	16.4	31.4	54.5	142.2
2000.....	0.8	1.4	2.5	3.8	6.6	9.0	17.0	29.4	56.6	126.4
1999.....	0.7	1.3	2.1	3.5	5.8	10.4	16.6	28.1	52.5	121.0

White, female

2003.....	0.5	0.7	1.6	2.4	5.0	8.2	15.5	33.2	66.5	196.8
2002.....	0.4	0.7	1.4	2.3	4.5	8.3	15.6	29.0	62.0	181.3
2001.....	0.3	0.7	1.4	2.4	4.8	8.2	14.5	30.5	61.1	172.2
2000.....	0.4	0.6	1.1	2.3	4.5	8.4	14.4	28.6	58.5	162.3
1999.....	0.3	0.8	1.2	1.9	4.3	7.9	14.0	27.2	53.8	150.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

2003.....	3.9	7.1	12.0	20.5	29.0	42.9	68.9	102.1	145.4	301.1
2002.....	3.8	7.4	10.9	18.4	24.3	36.5	61.5	96.4	151.0	308.8
2001.....	4.4	7.8	12.5	18.0	27.0	40.4	56.3	89.8	148.2	288.0
2000.....	4.2	7.4	13.2	18.2	29.3	38.8	58.3	92.7	136.1	284.1
1999.....	4.0	6.3	12.8	17.0	27.0	38.0	57.2	95.0	138.7	269.4

Black, male

2003.....	4.5	9.8	16.4	24.3	36.2	50.0	74.6	102.2	147.7	274.2
2002.....	4.7	9.2	13.3	21.7	25.7	40.6	67.1	105.2	164.3	274.5
2001.....	5.5	10.1	16.3	21.7	30.3	48.8	61.9	95.2	158.1	275.7
2000.....	5.3	9.3	16.5	23.6	35.9	38.4	60.9	89.7	135.3	249.0
1999.....	4.6	7.8	16.6	19.5	31.0	43.1	62.1	98.8	155.2	266.0

Black, female

2003.....	3.3	4.7	8.4	17.4	23.3	37.6	65.1	102.0	144.3	311.5
2002.....	3.0	5.9	8.8	15.6	23.2	33.5	57.8	91.1	144.0	322.0
2001.....	3.4	5.8	9.3	15.0	24.5	34.2	52.5	86.5	143.1	292.6
2000.....	3.2	5.7	10.4	14.0	24.3	39.1	56.5	94.7	136.6	298.8
1999.....	3.5	5.0	9.5	15.0	24.0	34.2	54.0	92.7	130.5	270.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2003.....	7.4	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.6	13.2	19.6	31.1	54.5	92.0	171.3	333.0	659.7	1,568.4
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8

All races 3/, male

2003.....	7.6	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.7	14.3	21.8	35.9	63.5	104.2	194.4	361.0	684.5	1,408.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8

All races 3/, female

2003.....	7.2	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.1	17.5	26.8	46.4	81.6	153.0	312.8	644.6	1,637.4
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

White, both sexes

2003.....	5.6	9.0	14.2	22.6	38.3	70.8	140.3	283.2	565.9	1,382.2
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.8	9.8	15.6	25.6	47.0	83.1	161.3	321.4	651.1	1,583.8
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6

White, male

2003.....	5.9	10.1	16.0	24.7	44.5	81.8	158.5	310.0	590.4	1,247.0
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.0	10.8	17.2	28.9	54.5	93.5	182.4	348.4	675.9	1,421.8
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7

White, female

2003.....	5.3	7.9	12.3	20.5	32.5	61.1	125.4	263.8	550.7	1,442.1
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.8	14.0	22.5	40.2	74.1	144.2	301.8	635.9	1,653.1
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

Black, both sexes

2003.....	18.6	31.4	52.7	72.4	113.1	157.4	256.3	433.2	677.0	1,348.7
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.8	53.0	78.5	115.3	168.1	278.6	476.9	802.9	1,533.9
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5

Black, male

2003.....	18.8	34.2	62.6	90.6	141.7	184.9	307.4	503.0	727.9	1,180.3
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.4	42.1	64.0	102.1	143.7	203.9	324.1	524.9	850.1	1,346.2
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3

Black, female

2003.....	18.5	29.0	44.4	57.3	90.7	137.1	221.8	391.8	650.3	1,414.2
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	34.1	44.0	60.0	94.1	141.5	246.4	446.0	776.9	1,612.4
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

All races 3/, both sexes

2003.....	0.2	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.3	10.6	22.7	52.9	186.3
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0

All races 3/, male

2003.....	0.2	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.7	6.7	13.2	27.0	58.9	167.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0

All races 3/, female

2003.....	*	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.6	19.6	49.2	194.5
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

White, both sexes

2003.....	0.2	0.3	0.6	1.0	2.4	4.4	8.9	20.2	44.1	159.5
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.5	5.3	10.6	23.1	53.4	193.0
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7

White, male

2003.....	0.2	0.5	0.8	1.4	3.1	5.2	10.1	23.8	48.4	143.3
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.7	13.4	27.7	59.4	173.0
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7

White, female

2003.....	*	*	0.4	0.7	1.7	3.6	7.8	17.6	41.4	166.7
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.3	19.7	49.6	201.6
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

Black, both sexes

2003.....	*	*	1.3	1.9	3.5	6.5	12.7	20.8	45.4	121.1
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.4	12.8	23.5	56.3	138.6
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3

Black, male

2003.....	*	*	*	*	4.8	8.1	18.1	28.8	59.1	104.3
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.3	8.2	13.4	24.6	65.6	136.2
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0

Black, female

2003.....	*	*	*	*	*	5.3	9.1	16.0	38.3	127.6
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.5	*	12.4	22.9	51.2	139.6
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2003.....	1.2	2.4	3.5	5.7	11.3	20.5	34.2	56.6	82.7	146.0
2002.....	1.3	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001.....	1.3	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000.....	1.3	2.1	3.6	6.5	12.9	24.7	41.1	62.9	93.0	161.1
1999.....	1.4	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2

All races 3/, male

2003.....	1.7	3.2	5.1	7.9	15.9	29.4	46.1	78.9	111.3	179.7
2002.....	1.8	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001.....	1.8	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000.....	1.6	2.8	5.0	9.5	18.3	35.2	58.0	87.0	124.7	203.9
1999.....	1.6	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5

All races 3/, female

2003.....	0.8	1.6	2.1	3.7	7.2	12.8	24.6	40.7	65.2	131.1
2002.....	0.7	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001.....	0.9	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000.....	0.9	1.5	2.1	3.7	8.0	15.8	27.7	45.6	73.6	142.6
1999.....	1.1	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of circulatory system (I71-I78)

White, both sexes

2003.....	1.1	2.0	3.1	5.5	11.0	20.6	34.7	57.5	82.9	146.9
2002.....	1.1	1.9	2.7	5.8	11.4	22.4	37.0	58.7	84.8	148.9
2001.....	1.1	1.8	2.9	5.6	11.7	22.6	36.8	60.6	88.4	158.2
2000.....	1.0	1.8	3.1	6.1	12.5	24.9	41.9	63.0	93.6	161.2
1999.....	1.2	1.9	3.0	6.3	12.6	23.7	40.8	64.1	94.8	165.1

White, male

2003.....	1.5	2.8	4.6	7.7	15.4	29.7	47.4	80.5	111.9	184.1
2002.....	1.6	2.7	3.8	7.9	16.0	32.5	49.8	80.4	116.1	190.9
2001.....	1.4	2.5	3.8	7.9	16.4	31.5	51.7	82.0	124.0	201.3
2000.....	1.3	2.5	4.4	9.0	18.1	36.2	59.7	87.9	127.8	209.5
1999.....	1.5	2.7	4.0	9.2	18.6	34.4	58.4	91.1	131.1	211.1

White, female

2003.....	0.7	1.3	1.7	3.4	6.9	12.6	24.4	40.9	65.0	130.5
2002.....	0.6	1.0	1.7	3.7	7.2	13.5	26.5	43.1	65.6	130.7
2001.....	0.7	1.1	2.0	3.4	7.4	14.7	24.7	45.5	67.0	140.1
2000.....	0.7	1.1	1.8	3.3	7.3	15.1	27.6	45.0	72.6	140.6
1999.....	0.9	1.0	1.9	3.6	7.1	14.5	26.9	45.1	73.4	146.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2003.....	2.6	5.3	7.3	8.5	16.3	24.0	35.9	55.3	92.1	150.0
2002.....	2.8	5.3	6.7	11.2	16.5	25.7	37.8	58.4	83.5	156.8
2001.....	3.2	4.0	6.5	10.7	18.0	22.9	42.8	69.3	96.2	174.7
2000.....	3.2	4.6	7.0	11.4	18.5	26.2	39.5	69.2	96.3	178.1
1999.....	2.8	4.5	6.1	11.4	13.5	26.1	38.6	64.3	96.5	178.9

Black, male

2003.....	3.5	6.8	9.9	11.2	23.5	33.5	40.1	72.6	118.9	138.7
2002.....	4.1	6.7	9.8	16.5	23.0	30.7	48.7	65.7	100.1	166.2
2001.....	4.5	5.5	9.4	13.1	23.8	28.0	55.9	79.0	127.6	165.7
2000.....	4.2	5.6	10.6	15.8	23.6	31.4	49.4	87.4	102.9	163.9
1999.....	3.1	6.1	8.6	14.1	17.8	31.9	49.2	74.6	117.7	184.0

Black, female

2003.....	1.8	4.0	5.1	6.2	10.7	17.0	33.0	45.0	78.1	154.3
2002.....	1.7	4.0	4.0	6.8	11.5	22.0	30.5	54.1	74.8	153.2
2001.....	2.0	2.7	4.1	8.7	13.4	19.1	34.1	63.5	80.1	178.1
2000.....	2.3	3.7	4.1	7.9	14.7	22.4	32.5	57.5	92.7	184.1
1999.....	2.5	3.2	4.0	9.1	10.1	21.7	31.5	58.1	86.0	177.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

All races 3/, both sexes

2003.....	0.8	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.3	4.4	8.5	16.5	28.1	41.1	57.0	78.1
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3

All races 3/, male

2003.....	1.3	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	2.0	3.6	6.8	13.4	25.3	42.2	60.8	83.6	120.8
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3

All races 3/, female

2003.....	0.4	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.2	8.9	17.0	26.9	40.6	59.8
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

White, both sexes

2003.....	0.7	1.4	2.1	3.6	7.3	14.0	23.6	38.1	52.2	72.4
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.1	4.1	8.5	16.9	29.4	42.0	58.5	79.9
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2

White, male

2003.....	1.2	2.1	3.4	5.7	11.4	21.1	34.6	56.2	76.5	110.0
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.3	6.5	13.5	26.6	44.3	62.4	87.3	126.6
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8

White, female

2003.....	0.3	0.7	0.8	1.5	3.5	7.9	14.6	25.0	37.2	55.7
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.6	0.9	1.9	4.0	8.6	17.4	27.2	40.8	59.9
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

Black, both sexes

2003.....	1.8	3.8	4.7	5.7	9.0	11.7	17.9	25.9	36.0	46.4
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.8	7.4	9.4	13.7	20.1	33.9	42.1	62.4
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3

Black, male

2003.....	2.8	5.8	7.1	8.0	14.7	16.3	20.4	32.9	49.5	51.1
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.7	6.4	10.8	13.4	16.9	26.4	46.0	44.8	63.9
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2

Black, female

2003.....	*	2.1	2.7	3.8	4.7	8.4	16.2	21.7	29.0	44.6
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.4	11.3	15.6	26.1	40.6	61.8
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes

2003.....	0.4	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.3	13.0	21.8	36.1	82.9
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9

All races 3/, male

2003.....	0.3	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.9	1.5	2.6	5.0	9.9	15.9	26.2	41.1	83.1
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2

All races 3/, female

2003.....	0.5	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.8	6.9	10.7	18.7	33.0	82.9
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2003.....	0.3	0.6	1.0	1.9	3.7	6.6	11.1	19.5	30.7	74.6
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.9	12.5	21.0	35.1	81.4
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9

White, male

2003.....	0.3	0.7	1.2	2.0	4.0	8.7	12.8	24.4	35.4	74.1
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.6	9.6	15.4	25.4	40.5	82.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3

White, female

2003.....	0.4	0.6	0.9	1.9	3.4	4.8	9.8	15.9	27.8	74.8
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.2	17.9	31.8	80.7
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2003.....	0.8	1.5	2.6	2.8	7.3	12.3	18.0	29.4	56.1	103.6
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.2	4.0	9.1	12.5	19.4	35.3	54.2	115.7
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6

Black, male

2003.....	*	*	2.8	3.2	8.9	17.2	19.7	39.7	69.4	87.6
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	2.0	4.2	5.0	10.2	14.5	22.9	41.4	58.1	100.0
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Black, female

2003.....	*	1.8	2.4	2.4	6.0	8.6	16.8	23.3	49.1	109.8
2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.2	11.1	17.0	31.4	52.1	122.3
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2003.....	0.9	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.3	1.5	1.7	2.7	3.5	5.4	7.7	12.0	25.2
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4

All races 3/, male

2003.....	1.0	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.8	2.0	2.7	3.9	5.7	7.9	12.2	21.1
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8

All races 3/, female

2003.....	0.9	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.6	1.0	1.3	1.5	2.6	3.1	5.1	7.6	11.9	27.0
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2003.....	0.8	1.1	1.4	1.6	2.2	3.0	4.7	6.5	11.6	23.6
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	3.0	5.1	7.3	11.9	25.2
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2

White, male

2003.....	0.9	1.2	1.6	2.1	2.3	3.2	4.6	6.4	10.2	18.5
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.5	5.6	7.6	12.1	21.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7

White, female

2003.....	0.7	1.1	1.2	1.1	2.2	2.9	4.7	6.5	12.4	25.8
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.2	1.4	2.4	2.5	4.8	7.1	11.8	27.0
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2003.....	2.3	2.6	2.5	3.6	5.2	6.2	9.3	13.8	17.8	33.6
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.3	4.0	5.2	8.4	9.0	14.6	14.7	30.1
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4

Black, male

2003.....	1.8	3.4	3.4	3.6	7.3	7.6	10.2	15.5	20.7	32.3
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.2	3.9	5.0	5.8	9.0	8.3	12.7	*	27.7
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*

Black, female

2003.....	2.7	1.9	1.8	3.6	3.6	5.1	8.6	12.8	16.2	34.0
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.0	9.5	15.8	15.5	31.1
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

2003.....	2.7	4.2	6.4	8.2	15.2	26.6	49.4	103.3	216.8	666.1
2002.....	2.6	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001.....	2.6	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000.....	2.9	3.9	5.7	9.0	15.8	26.9	53.3	106.8	242.9	734.4
1999.....	2.7	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8

All races 3/, male

2003.....	3.1	4.9	8.1	9.6	17.9	33.3	62.1	130.1	272.3	757.8
2002.....	3.1	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001.....	3.0	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000.....	3.7	4.7	7.2	11.1	19.2	33.4	66.3	131.7	304.1	846.3
1999.....	3.1	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7

All races 3/, female

2003.....	2.3	3.5	4.8	6.8	12.7	20.7	39.2	84.2	182.9	625.6
2002.....	2.2	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001.....	2.2	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000.....	2.1	3.2	4.4	7.1	12.6	21.3	42.9	88.8	205.4	686.2
1999.....	2.3	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

White, both sexes

2003.....	2.3	3.7	5.7	7.4	14.0	25.2	47.9	102.4	218.4	679.3
2002.....	2.3	3.8	5.0	7.7	13.8	24.5	49.9	105.4	228.0	709.1
2001.....	2.2	3.3	4.8	7.5	13.0	23.3	48.1	98.6	216.6	695.0
2000.....	2.5	3.4	5.0	8.1	14.5	24.9	51.0	105.6	242.2	749.9
1999.....	2.1	3.4	4.6	7.7	13.1	24.0	48.8	105.5	235.5	764.3

White, male

2003.....	2.6	4.3	7.2	8.5	15.7	30.8	59.2	127.2	271.4	767.1
2002.....	2.7	4.4	6.0	9.1	16.5	29.8	62.4	131.4	286.5	815.9
2001.....	2.5	4.2	5.7	9.0	15.4	28.3	59.3	125.5	272.3	814.4
2000.....	3.1	3.9	6.0	9.6	17.5	30.2	62.9	129.5	301.8	864.2
1999.....	2.3	4.2	5.7	8.6	15.1	30.1	60.9	132.2	301.9	905.7

White, female

2003.....	2.1	3.1	4.3	6.3	12.4	20.2	38.6	84.5	185.7	640.4
2002.....	1.9	3.2	3.9	6.3	11.3	19.8	39.7	86.8	192.1	662.6
2001.....	1.8	2.5	4.0	6.0	10.7	19.0	39.0	79.6	182.9	645.0
2000.....	1.8	2.8	4.0	6.7	11.7	20.3	41.4	88.3	205.6	701.0
1999.....	1.9	2.6	3.6	6.7	11.2	18.7	39.1	86.8	196.4	707.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

Black, both sexes

2003.....	5.5	8.5	12.6	15.7	27.1	41.6	70.1	122.5	205.6	525.4
2002.....	5.2	7.5	10.1	17.2	26.3	35.5	68.9	123.3	231.6	575.2
2001.....	5.8	9.0	10.7	15.5	23.3	34.3	65.7	133.7	224.0	581.3
2000.....	6.1	8.5	12.6	18.1	27.5	44.2	76.8	124.8	256.0	602.5
1999.....	6.9	9.7	11.8	16.2	26.6	36.8	75.2	118.7	219.9	634.8

Black, male

2003.....	7.0	10.4	16.7	21.3	39.7	61.6	100.5	177.6	285.0	641.2
2002.....	6.2	9.9	12.9	23.7	36.0	48.1	89.8	173.2	321.7	689.4
2001.....	7.0	11.4	15.7	20.8	33.6	50.1	94.9	203.5	309.0	714.8
2000.....	8.3	11.6	18.2	27.0	36.8	62.1	102.6	166.2	338.7	697.0
1999.....	9.0	13.4	17.2	21.7	34.2	51.9	109.6	160.0	312.2	803.8

Black, female

2003.....	4.1	7.0	9.1	11.1	17.3	26.9	49.6	89.8	164.0	480.4
2002.....	4.3	5.5	7.7	12.0	18.7	26.1	54.9	93.6	184.6	531.5
2001.....	4.8	6.9	6.5	11.2	15.2	22.7	46.2	92.1	180.7	531.7
2000.....	4.2	5.8	8.0	11.1	20.6	30.9	58.5	98.2	210.5	562.9
1999.....	5.1	6.6	7.1	11.8	20.6	25.6	52.4	93.8	174.1	571.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza (J10-J11)

All races 3/, both sexes

2003.....	*	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.6	6.2	19.9
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2

All races 3/, male

2003.....	*	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.6	6.2	19.6
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9

All races 3/, female

2003.....	*	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.2	20.0
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza (J10-J11)

White, both sexes

2003.....	*	0.1	0.2	0.3	0.3	0.8	1.4	2.9	5.1	18.2
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.9	1.4	2.7	6.5	21.1
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8

White, male

2003.....	*	*	*	0.3	*	0.9	1.4	3.1	5.2	18.3
2002.....	*	*	*	*	*	*	*	1.0	2.8	9.7
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	0.5	1.1	1.7	2.7	6.6	20.6
1999.....	*	*	*	*	0.5	0.6	1.2	2.6	7.5	21.7

White, female

2003.....	*	*	*	*	0.4	0.7	1.4	2.8	5.0	18.1
2002.....	*	*	*	*	*	*	*	0.8	2.2	9.3
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	*	*	0.6	0.7	1.2	2.7	6.4	21.3
1999.....	*	*	*	*	*	0.7	1.1	2.0	5.7	23.2

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

All races 3/, both sexes

2003.....	2.6	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.6	8.7	15.3	26.1	51.9	104.2	236.8	714.5
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6

All races 3/, male

2003.....	3.0	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	7.0	10.8	18.8	32.4	64.6	129.1	298.0	826.7
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7

All races 3/, female

2003.....	2.2	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.3	6.8	12.1	20.7	41.7	86.3	199.2	666.1
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

White, both sexes

2003.....	2.3	3.6	5.6	7.1	13.7	24.4	46.5	99.5	213.3	661.1
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.3	4.9	7.8	13.9	24.0	49.6	102.9	235.8	728.8
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6

White, male

2003.....	2.5	4.2	7.0	8.2	15.4	29.9	57.8	124.1	266.2	748.8
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.9	5.8	9.3	17.0	29.2	61.2	126.8	295.2	843.6
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0

White, female

2003.....	2.0	3.0	4.2	6.0	12.1	19.5	37.2	81.7	180.7	622.3
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.7	2.7	3.9	6.4	11.2	19.6	40.2	85.6	199.2	679.7
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

Black, both sexes

2003.....	5.4	8.3	12.4	15.4	27.0	41.0	69.6	121.7	203.6	519.6
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.3	12.5	17.9	27.2	43.7	75.4	124.0	253.7	593.1
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1

Black, male

2003.....	7.0	10.0	16.5	20.8	39.5	60.8	100.2	176.6	283.5	638.1
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.3	18.0	26.6	36.6	61.6	101.6	164.3	337.9	685.3
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3

Black, female

2003.....	4.1	6.7	8.9	11.0	17.1	26.3	49.0	89.0	161.7	473.5
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.7	8.0	11.1	20.1	30.4	56.7	97.9	207.3	554.5
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2003.....	*	*	*	*	*	*	*	0.5	0.8	3.5
2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.6
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2003.....	*	*	*	*	*	*	*	*	0.8	3.9
2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	0.9	3.7
2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

White, male

2003.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	0.8	*	3.6
1999.....	*	*	*	*	*	*	*	0.8	1.6	4.0

White, female

2003.....	*	*	*	*	*	*	*	*	0.8	4.2
2002.....	*	*	*	*	*	*	*	*	0.9	3.7
2001.....	*	*	*	*	*	*	*	*	0.8	3.9
2000.....	*	*	*	*	*	*	*	*	1.0	5.1
1999.....	*	*	*	*	*	*	*	0.5	1.1	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	0.5	2.0
2001.....	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	0.4	0.8	3.6

All races 3/, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	2.2
1999.....	*	*	*	*	*	*	*	*	2.6

All races 3/, female

2003.....	*	*	*	*	*	*	*	*	2.5
2002.....	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	0.7	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	0.4	0.9	3.8

White, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	2.3
1999.....	*	*	*	*	*	*	*	*	2.6

White, female

2003.....	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	2.3
2001.....	*	*	*	*	*	*	*	*	2.5
2000.....	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	0.7	4.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2003.....	2.8	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.7	5.9	12.0	29.9	63.4	125.5	221.0	328.0	477.6	640.1
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0

All races 3/, male

2003.....	2.5	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.3	12.4	32.4	69.3	141.9	259.8	396.9	619.0	918.3
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7

All races 3/, female

2003.....	3.1	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.7	27.6	58.0	111.6	190.2	278.4	390.9	520.2
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2003.....	2.5	5.6	11.7	28.0	67.3	129.5	223.2	347.2	478.9	664.5
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.6	30.3	66.2	132.4	232.8	343.1	498.6	668.1
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9

White, male

2003.....	2.2	6.1	12.0	29.4	71.5	139.3	252.4	406.9	585.6	894.4
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.7	32.2	70.8	145.8	268.7	408.1	638.8	951.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4

White, female

2003.....	2.8	5.1	11.4	26.7	63.5	120.8	199.3	304.0	413.2	562.8
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.4	28.5	62.0	120.7	203.9	295.9	412.2	547.0
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2003.....	5.7	9.1	16.7	31.3	52.7	86.5	146.7	211.9	271.9	353.0
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.2	18.8	33.5	58.0	93.6	149.8	212.4	283.1	374.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2

Black, male

2003.....	5.1	9.8	17.7	38.0	67.7	112.0	212.5	326.8	471.8	601.6
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.7	20.8	41.3	76.8	132.6	215.7	314.6	441.7	639.6
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9

Black, female

2003.....	6.2	8.6	15.8	25.7	41.1	67.6	102.4	143.5	167.1	256.4
2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.7	17.2	27.4	44.0	64.6	103.1	146.6	195.9	264.1
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes

2003.....	*	*	*	0.1	0.2	0.4	0.7	1.4	2.6	7.3
2002.....	*	*	0.1	0.1	0.3	0.5	1.0	1.8	2.9	8.1
2001.....	*	0.1	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000.....	*	*	*	0.2	0.4	0.7	1.0	2.1	3.6	11.7
1999.....	*	*	0.1	0.3	0.6	0.7	1.2	1.9	4.1	10.7

All races 3/, male

2003.....	*	*	*	*	*	*	0.8	1.9	2.6	7.0
2002.....	*	*	*	*	*	0.7	1.1	2.2	2.9	8.8
2001.....	*	*	*	*	0.4	0.8	1.1	2.2	2.5	9.2
2000.....	*	*	*	*	0.5	0.9	1.0	2.7	4.8	12.2
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.4	4.7	11.9

All races 3/, female

2003.....	*	*	*	*	*	0.4	0.7	1.1	2.5	7.5
2002.....	*	*	*	*	*	*	0.9	1.4	2.9	7.8
2001.....	*	*	*	*	*	0.4	0.7	1.4	2.6	9.1
2000.....	*	*	*	*	0.4	0.5	1.1	1.7	2.9	11.4
1999.....	*	*	*	*	0.6	0.5	1.0	1.5	3.8	10.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, chronic and unspecified (J40-J42)

White, both sexes

2003.....	*	*	*	*	0.2	0.5	0.7	1.5	2.6	7.5
2002.....	*	*	*	*	0.3	0.6	1.1	1.9	3.0	8.6
2001.....	*	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	9.6
2000.....	*	*	*	*	0.5	0.7	1.1	2.2	3.7	12.5
1999.....	*	*	*	0.3	0.6	0.7	1.2	1.9	4.1	11.2

White, male

2003.....	*	*	*	*	*	*	0.7	1.9	2.6	7.3
2002.....	*	*	*	*	*	0.7	1.2	2.4	2.9	9.3
2001.....	*	*	*	*	*	0.8	1.0	2.2	2.5	9.4
2000.....	*	*	*	*	0.5	1.0	1.0	2.8	5.1	13.2
1999.....	*	*	*	0.4	0.5	0.9	1.5	2.4	4.4	12.4

White, female

2003.....	*	*	*	*	*	0.5	0.8	1.2	2.6	7.6
2002.....	*	*	*	*	*	*	1.0	1.6	3.0	8.3
2001.....	*	*	*	*	*	*	0.7	1.5	2.8	9.7
2000.....	*	*	*	*	0.5	0.5	1.1	1.7	2.9	12.3
1999.....	*	*	*	*	0.7	0.5	1.0	1.5	3.9	10.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Emphysema (J43)

All races 3/, both sexes

2003.....	0.3	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.1	10.6	21.0	36.0	47.5	61.0	62.5
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1

All races 3/, male

2003.....	0.3	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.7	11.4	24.9	42.9	59.3	82.3	98.8
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4

All races 3/, female

2003.....	0.3	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.8	17.6	30.5	39.0	47.9	46.9
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Emphysema (J43)

White, both sexes

2003.....	0.3	0.7	1.6	4.2	9.9	18.0	30.5	44.3	54.2	57.0
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.3	22.3	38.4	50.2	64.1	65.1
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7

White, male

2003.....	0.3	1.0	1.9	4.5	10.6	19.5	35.9	53.2	69.6	80.4
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.9	25.5	45.1	62.0	85.7	100.7
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2

White, female

2003.....	0.3	0.5	1.4	3.9	9.2	16.7	26.0	37.8	44.7	46.6
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.6	4.8	10.8	19.5	33.0	41.7	50.7	49.8
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Emphysema (J43)

Black, both sexes

2003.....	*	1.0	1.3	3.6	7.2	10.2	16.4	25.2	30.5	30.9
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.6	4.7	7.8	14.3	19.3	24.6	32.1	37.3
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6

Black, male

2003.....	*	*	2.0	5.2	10.2	13.3	24.6	43.4	50.2	70.9
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.7	6.2	11.3	24.2	28.7	33.7	52.3	80.9
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6

Black, female

2003.....	*	*	*	2.3	4.8	7.9	10.9	14.4	20.1	15.4
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.0	12.7	18.8	21.0	19.1
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

All races 3/, both sexes

2003.....	1.1	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.5	4.6	5.7	8.8	16.5
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3

All races 3/, male

2003.....	0.9	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.7	1.9	2.8	3.4	3.9	6.6	12.2
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7

All races 3/, female

2003.....	1.4	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.1	5.6	7.0	10.1	18.3
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

White, both sexes

2003.....	0.8	1.1	1.1	1.5	1.8	2.3	2.6	4.4	6.7	14.3
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.1	5.2	8.5	15.7
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6

White, male

2003.....	0.6	0.9	0.8	1.2	1.2	1.5	2.0	3.0	5.4	10.6
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.1	3.1	3.6	6.2	10.8
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8

White, female

2003.....	1.0	1.3	1.5	1.7	2.3	3.0	3.1	5.4	7.5	15.9
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.5	4.9	6.4	9.9	17.8
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

Black, both sexes

2003.....	3.4	4.1	5.7	5.2	5.0	6.6	7.5	8.2	10.4	12.0
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.5	5.5	6.7	7.3	8.5	9.7	10.7	10.3	22.3
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4

Black, male

2003.....	2.7	3.5	4.4	3.8	4.8	4.9	*	9.1	*	*
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.1	3.7	6.5	6.3	8.2	6.7	*	*	23.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*

Black, female

2003.....	4.1	4.7	6.7	6.3	5.1	7.9	8.6	7.6	11.6	13.8
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.7	7.0	6.8	8.1	8.7	11.8	14.7	11.0	21.8
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic lower respiratory diseases (J44,J47)

All races 3/, both sexes

2003.....	1.4	3.6	8.6	21.5	52.2	101.4	178.5	281.8	394.6	558.9
2002.....	1.4	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001.....	1.3	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000.....	1.2	3.4	8.4	22.6	49.8	100.4	179.4	272.7	404.3	549.5
1999.....	1.2	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8

All races 3/, male

2003.....	1.3	4.1	9.1	23.3	57.0	112.2	206.5	338.3	493.3	769.0
2002.....	1.6	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001.....	1.3	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000.....	1.2	3.9	8.9	24.9	55.5	113.3	212.6	331.0	525.2	795.1
1999.....	1.2	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7

All races 3/, female

2003.....	1.4	3.2	8.0	19.8	47.8	92.1	156.0	241.5	334.2	466.0
2002.....	1.3	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001.....	1.4	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000.....	1.2	2.9	7.9	20.6	44.7	89.4	153.0	230.8	330.1	443.7
1999.....	1.1	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic lower respiratory diseases (J44,J47)

White, both sexes

2003.....	1.3	3.8	8.9	22.3	55.5	108.7	189.4	297.0	415.4	585.8
2002.....	1.4	3.6	8.8	22.9	51.9	105.4	185.7	294.0	418.6	580.7
2001.....	1.4	3.3	8.6	23.4	53.1	106.6	187.6	289.8	404.4	581.3
2000.....	1.2	3.4	8.5	23.4	52.4	106.5	189.3	285.5	422.3	574.8
1999.....	1.1	3.0	8.4	24.4	54.2	108.1	191.8	292.0	422.6	580.3

White, male

2003.....	1.2	4.2	9.3	23.6	59.4	117.8	213.8	348.8	507.9	796.0
2002.....	1.6	3.8	9.5	23.7	54.7	114.7	211.4	353.0	526.8	810.4
2001.....	1.3	3.9	9.0	24.8	56.2	115.4	215.7	351.7	514.4	817.8
2000.....	1.2	3.8	8.6	25.2	57.0	117.3	219.6	339.7	541.9	826.7
1999.....	1.1	3.5	8.6	26.2	60.3	123.4	230.5	365.1	558.5	897.0

White, female

2003.....	1.5	3.3	8.5	21.0	51.8	100.6	169.5	259.5	358.4	492.7
2002.....	1.3	3.5	8.2	22.1	49.3	97.1	164.8	251.8	352.3	480.9
2001.....	1.4	2.8	8.2	22.1	50.2	98.9	164.8	246.0	337.9	482.3
2000.....	1.1	3.0	8.3	21.7	48.2	97.2	164.8	246.2	348.7	467.1
1999.....	1.1	2.6	8.3	22.7	48.7	94.7	161.1	240.6	342.4	453.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic lower respiratory diseases (J44,J47)

Black, both sexes

2003.....	1.8	3.8	9.4	22.1	40.4	69.5	122.2	178.0	228.7	304.6
2002.....	1.7	3.9	10.1	20.3	41.8	67.8	116.9	184.3	252.5	320.1
2001.....	1.6	4.1	9.4	20.7	37.6	75.7	119.1	173.3	238.4	320.9
2000.....	1.8	3.9	10.5	22.0	42.7	70.0	119.7	175.3	237.8	311.4
1999.....	2.0	4.8	8.5	23.9	43.1	77.1	129.9	199.1	248.8	313.7

Black, male

2003.....	1.9	4.5	11.0	28.8	52.4	93.8	181.3	273.0	410.5	520.3
2002.....	2.0	4.8	12.7	27.5	56.2	88.8	174.3	291.6	407.3	565.0
2001.....	1.6	4.4	11.3	26.6	50.4	105.7	179.4	285.3	408.5	583.2
2000.....	1.7	5.2	13.3	28.5	58.7	99.7	179.1	274.5	377.7	532.1
1999.....	2.5	6.0	11.0	31.2	61.3	109.6	192.3	325.8	441.2	607.0

Black, female

2003.....	1.7	3.1	8.0	16.6	31.0	51.5	82.4	121.5	133.5	220.8
2002.....	1.4	3.1	8.0	14.4	30.5	52.2	78.4	120.2	171.9	226.3
2001.....	1.5	3.8	7.7	15.8	27.5	53.5	78.9	106.5	151.7	223.5
2000.....	1.9	2.9	8.1	16.8	30.8	48.0	77.5	111.4	160.8	219.2
1999.....	1.5	3.7	6.5	17.9	28.8	53.1	88.4	122.7	153.4	204.1

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.8	14.4	30.5	66.2	185.2
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6

All races 3/, male

2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	4.0	9.3	20.4	41.6	93.8	261.9
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4

All races 3/, female

2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.7	9.6	22.4	49.4	152.1
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

White, both sexes

2003.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	3.0	6.5	13.6	30.1	66.1	188.9
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0

White, male

2003.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.7	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.4	2.4	3.7	8.7	18.9	40.7	94.7	270.9
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6

White, female

2003.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.6	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.3	22.5	48.5	153.8
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2003.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.4
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.3	2.9	5.8	10.8	24.9	39.1	74.5	160.3
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1

Black, male

2003.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.3	3.4	7.6	16.7	39.2	60.1	94.6	193.7
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5

Black, female

2003.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	144.2
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.5	6.5	14.7	25.5	63.5	146.3
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2003.....	1.3	2.6	4.0	6.6	11.6	21.0	36.2	57.3	83.8	138.1
2002.....	1.5	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001.....	1.5	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000.....	1.7	2.6	4.7	7.8	12.8	23.2	37.3	60.5	90.0	139.9
1999.....	1.6	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6

All races 3/, male

2003.....	1.4	2.8	4.3	7.5	13.8	25.9	43.8	71.8	105.5	165.0
2002.....	1.6	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001.....	1.8	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000.....	1.8	2.9	5.1	8.9	14.9	27.2	46.0	75.9	110.7	170.5
1999.....	1.7	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3

All races 3/, female

2003.....	1.3	2.5	3.7	5.7	9.6	16.8	30.0	47.0	70.5	126.2
2002.....	1.4	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001.....	1.3	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000.....	1.6	2.4	4.2	6.8	10.9	19.7	30.4	49.5	77.4	126.7
1999.....	1.4	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2003.....	1.1	2.2	3.7	6.3	11.5	21.5	37.4	58.5	87.4	143.2
2002.....	1.3	2.1	3.8	6.7	11.6	21.0	38.2	58.7	85.5	137.9
2001.....	1.3	2.3	3.9	7.0	12.4	23.0	38.7	63.5	92.3	146.0
2000.....	1.6	2.3	4.3	7.6	12.6	23.2	38.2	62.0	91.8	144.3
1999.....	1.3	2.3	3.8	6.4	12.6	22.6	37.0	60.6	89.5	150.7

White, male

2003.....	1.2	2.5	4.1	7.1	13.7	26.4	45.2	72.7	110.0	170.5
2002.....	1.4	2.1	4.0	7.6	13.4	25.3	48.3	72.5	106.8	162.0
2001.....	1.7	2.5	4.4	7.6	14.5	27.5	48.3	78.4	113.3	173.5
2000.....	1.6	2.4	4.7	8.6	14.6	27.3	46.9	77.5	113.3	175.9
1999.....	1.5	2.4	4.2	6.7	14.7	27.7	47.0	76.5	117.6	185.7

White, female

2003.....	1.0	2.0	3.3	5.4	9.6	17.2	30.9	48.2	73.6	131.0
2002.....	1.2	2.0	3.6	5.9	10.0	17.2	30.0	48.7	72.5	127.4
2001.....	1.0	2.0	3.4	6.4	10.4	19.1	31.0	52.9	79.6	134.4
2000.....	1.5	2.1	3.9	6.6	10.7	19.5	31.2	50.7	78.5	130.8
1999.....	1.1	2.2	3.5	6.2	10.7	18.2	29.1	49.5	72.9	136.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2003.....	3.1	5.7	7.3	10.5	14.4	20.2	29.1	51.4	50.0	89.6
2002.....	2.9	5.5	6.9	11.1	16.8	20.7	33.9	40.5	68.3	90.3
2001.....	3.3	5.7	9.3	10.9	17.0	24.2	36.1	50.2	66.9	109.6
2000.....	3.0	6.1	8.1	11.8	15.7	27.0	33.1	49.9	74.2	98.5
1999.....	3.4	5.8	7.1	11.7	15.4	25.0	38.0	52.8	66.2	111.6

Black, male

2003.....	2.5	5.5	7.3	12.6	18.5	25.6	37.8	70.8	65.0	113.6
2002.....	3.4	5.7	7.6	12.4	21.2	22.2	44.4	53.3	86.4	100.8
2001.....	3.2	6.2	9.4	10.5	21.6	28.0	39.4	60.4	85.1	129.3
2000.....	3.3	7.4	9.1	13.9	18.4	31.2	38.6	65.1	81.4	122.4
1999.....	3.3	6.6	7.8	13.7	17.8	28.3	44.1	66.5	83.7	145.3

Black, female

2003.....	3.6	5.8	7.3	8.8	11.1	16.3	23.3	39.9	42.2	80.2
2002.....	2.3	5.3	6.2	10.0	13.3	19.6	26.9	32.9	58.9	86.3
2001.....	3.3	5.3	9.1	11.3	13.4	21.4	33.9	44.1	57.6	102.3
2000.....	2.6	5.0	7.2	10.2	13.7	23.9	29.2	40.2	70.3	88.5
1999.....	3.5	5.1	6.6	10.1	13.5	22.5	34.0	44.6	57.5	99.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

All races 3/, both sexes

2003.....	0.3	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.7	1.0	1.5	2.3	3.2	5.1	8.7	15.8	35.1
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6

All races 3/, male

2003.....	0.4	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.6	2.1	3.0	4.0	7.0	11.0	17.5	38.6
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5

All races 3/, female

2003.....	0.2	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	1.0	1.7	2.5	3.7	7.0	14.7	33.6
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

White, both sexes

2003.....	0.3	0.5	0.7	1.1	1.6	2.4	4.2	7.2	13.4	26.9
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.3	3.1	5.1	8.6	16.1	35.9
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3

White, male

2003.....	0.3	0.7	1.0	1.3	2.2	3.1	5.3	8.6	15.6	29.0
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	3.0	3.9	6.8	10.7	17.6	38.8
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4

White, female

2003.....	0.2	0.3	0.5	0.9	1.0	1.7	3.3	6.2	12.1	26.0
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	7.0	15.2	34.6
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

Black, both sexes

2003.....	*	1.2	1.8	2.1	2.4	2.1	4.5	8.7	14.5	16.9
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.1	2.6	2.8	3.9	5.7	9.5	11.5	26.7
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5

Black, male

2003.....	*	1.6	3.1	3.1	3.9	*	7.2	11.4	23.6	*
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.4	3.6	*	*	9.6	12.3	*	28.7
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1

Black, female

2003.....	*	*	*	*	*	*	*	7.1	9.7	15.8
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.6	9.1	25.8
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of appendix (K35-K38)

All races 3/, both sexes

2003.....	*	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

All races 3/, male

2003.....	*	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.5	*	0.6	1.0	1.6	1.9
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9

All races 3/, female

2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.3	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of appendix (K35-K38)

White, both sexes

2003.....	*	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5

White, male

2003.....	*	*	*	*	*	*	0.7	0.8	2.4	2.5
2002.....	*	*	*	*	0.5	*	0.7	1.2	1.6	2.0
2001.....	*	*	*	*	*	*	1.0	0.7	1.5	2.5
2000.....	*	*	*	*	0.5	*	*	0.9	1.5	1.9
1999.....	*	*	*	*	*	*	*	0.9	*	3.1

White, female

2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.9	2.1
2001.....	*	*	*	*	*	*	*	*	*	1.9
2000.....	*	*	*	*	*	*	*	*	1.2	1.6
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hernia (K40-K46)

All races 3/, both sexes

2003.....	0.1	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	12.9
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7

All races 3/, male

2003.....	*	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	2.9	5.6	11.9
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2

All races 3/, female

2003.....	*	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.1	4.5	13.4
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hernia (K40-K46)

White, both sexes

2003.....	*	0.2	0.3	0.4	0.7	1.1	1.6	2.8	4.7	12.3
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.2
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0

White, male

2003.....	*	*	0.4	0.4	0.7	1.1	1.6	3.1	5.1	12.5
2002.....	*	*	*	0.4	0.5	0.8	1.6	2.9	5.4	11.9
2001.....	*	*	0.3	*	0.7	0.9	1.4	2.3	5.2	11.8
2000.....	*	*	0.3	*	0.6	0.8	1.5	3.0	5.6	12.1
1999.....	*	*	*	0.4	0.8	0.8	1.6	2.5	6.6	12.9

White, female

2003.....	*	*	*	0.5	0.7	1.0	1.6	2.6	4.5	12.2
2002.....	*	*	*	0.4	0.6	0.9	1.6	3.0	5.4	14.0
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	5.0	13.2
2000.....	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
1999.....	*	*	*	*	0.5	0.7	1.5	2.5	4.9	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

2003.....	9.3	15.9	21.1	22.0	24.4	28.0	31.2	31.0	28.5	20.1
2002.....	9.7	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001.....	9.8	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000.....	10.1	17.0	19.0	22.1	26.5	29.2	31.2	31.9	29.8	22.8
1999.....	9.9	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2

All races 3/, male

2003.....	12.8	23.4	32.3	32.4	35.6	38.4	40.0	38.1	35.0	27.5
2002.....	12.9	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001.....	13.2	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000.....	14.2	25.7	29.3	32.3	38.5	39.9	41.0	40.0	34.0	32.6
1999.....	14.5	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1

All races 3/, female

2003.....	5.9	8.6	10.3	12.2	14.2	19.0	24.1	26.0	24.5	16.8
2002.....	6.5	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001.....	6.4	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000.....	6.1	8.5	9.3	12.7	15.8	20.1	23.5	26.1	27.2	18.6
1999.....	5.4	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2003.....	9.7	16.2	21.0	22.0	24.6	28.9	32.8	32.9	29.8	20.9
2002.....	10.0	16.1	20.2	21.8	24.8	28.9	32.7	34.2	30.9	22.3
2001.....	9.8	17.2	19.6	20.7	25.3	29.7	32.3	32.6	30.0	23.2
2000.....	10.0	16.9	18.4	21.8	26.8	29.9	32.3	33.4	30.7	23.7
1999.....	9.7	15.8	18.3	20.8	26.3	30.2	32.6	33.7	32.0	23.8

White, male

2003.....	13.4	24.0	32.0	32.2	35.9	39.5	41.8	40.1	36.2	28.4
2002.....	13.5	23.8	30.9	32.1	34.7	38.2	41.6	41.6	38.1	29.6
2001.....	13.2	25.5	29.3	30.6	35.6	40.2	42.1	39.1	37.2	33.5
2000.....	14.2	26.2	28.3	31.9	38.8	41.0	42.2	41.5	35.4	33.9
1999.....	14.2	24.6	28.4	31.5	37.4	42.2	42.5	41.1	40.3	34.2

White, female

2003.....	5.9	8.4	10.2	12.2	14.2	19.6	25.4	27.8	25.9	17.5
2002.....	6.5	8.5	9.8	11.9	15.6	20.7	25.4	28.9	26.5	19.1
2001.....	6.3	8.9	10.0	11.3	15.8	20.4	24.3	28.0	25.7	18.9
2000.....	5.8	7.9	8.8	12.3	15.7	20.2	24.3	27.5	27.9	19.3
1999.....	5.1	7.2	8.5	10.7	16.2	19.8	24.7	28.4	27.2	19.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2003.....	7.7	15.0	25.6	26.4	25.0	25.4	21.7	14.1	14.7	12.3
2002.....	7.9	17.4	25.0	24.6	25.5	22.9	20.0	20.1	16.0	12.2
2001.....	10.6	19.2	26.8	27.7	23.6	26.9	22.6	21.0	14.6	10.4
2000.....	10.9	19.5	26.8	26.3	26.2	26.5	24.5	18.4	18.3	13.8
1999.....	11.9	22.5	22.9	28.2	30.7	27.7	25.3	23.2	18.8	17.2

Black, male

2003.....	9.7	21.4	41.5	42.2	38.9	36.7	28.9	19.6	22.1	21.9
2002.....	9.8	23.4	38.2	36.9	39.0	34.9	29.0	23.0	21.4	*
2001.....	14.5	28.2	42.2	43.8	38.4	43.2	33.7	26.9	*	*
2000.....	14.5	27.1	43.1	40.2	39.0	36.4	35.7	26.4	19.1	*
1999.....	17.1	34.5	36.6	44.1	42.4	43.4	37.3	31.8	25.3	*

Black, female

2003.....	5.9	9.3	12.2	13.4	14.1	17.0	16.8	10.9	10.8	8.5
2002.....	6.3	12.2	13.7	14.5	14.9	14.1	13.9	18.4	13.1	10.7
2001.....	7.1	11.3	13.7	14.5	12.0	14.8	15.2	17.5	14.7	9.7
2000.....	7.7	13.1	13.5	15.4	16.8	19.1	16.5	13.2	17.8	12.0
1999.....	7.3	12.2	11.4	15.2	21.5	16.2	17.3	18.0	15.6	15.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcoholic liver disease (K70)

All races 3/, both sexes

2003.....	5.9	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.1	10.9	11.8	13.1	11.5	9.2	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8

All races 3/, male

2003.....	8.0	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.3	15.4	17.4	18.3	20.7	19.0	15.8	13.5	7.6	5.0
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1

All races 3/, female

2003.....	3.8	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.8	4.8	4.9	5.8	6.2	5.2	4.0	2.6	2.5	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcoholic liver disease (K70)

White, both sexes

2003.....	6.2	9.7	11.8	11.8	11.6	11.2	8.6	7.0	4.5	1.9
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.5	10.0	10.7	11.6	13.1	11.8	9.4	7.3	4.4	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7

White, male

2003.....	8.4	14.6	18.4	18.3	18.7	17.9	14.6	11.8	8.2	3.7
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.3	15.8	17.1	18.2	20.9	19.4	16.0	13.8	7.7	5.2
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8

White, female

2003.....	3.9	4.9	5.4	5.5	5.1	5.3	3.7	3.5	2.2	1.1
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.7	4.4	4.5	5.4	6.0	5.2	4.1	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcoholic liver disease (K70)

Black, both sexes

2003.....	4.2	7.6	12.7	12.2	12.9	11.2	6.9	*	*	*
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.2	13.8	14.1	12.7	11.4	8.3	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*

Black, male

2003.....	5.4	12.0	20.6	19.5	20.8	18.7	12.5	*	*	*
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.5	15.4	21.8	21.3	18.8	18.4	15.9	10.9	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*

Black, female

2003.....	3.0	3.7	6.0	6.1	6.8	5.7	*	*	*	*
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.6	7.2	8.4	8.2	6.1	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes

2003.....	3.4	6.5	9.3	10.4	12.7	17.0	22.9	24.5	24.1	18.2
2002.....	3.7	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001.....	3.6	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000.....	3.6	6.9	8.1	10.3	13.4	17.7	22.0	24.7	25.4	20.6
1999.....	3.5	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4

All races 3/, male

2003.....	4.8	9.2	14.0	14.3	17.0	20.7	25.8	26.8	26.8	23.9
2002.....	4.9	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001.....	4.8	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000.....	4.9	10.3	11.9	14.0	17.8	21.0	25.2	26.5	26.4	27.5
1999.....	5.2	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0

All races 3/, female

2003.....	2.0	3.8	4.8	6.6	8.9	13.8	20.5	22.8	22.5	15.8
2002.....	2.6	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001.....	2.5	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000.....	2.2	3.6	4.4	6.9	9.6	14.9	19.5	23.5	24.7	17.6
1999.....	1.8	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2003.....	3.5	6.5	9.2	10.2	13.0	17.7	24.2	26.0	25.3	19.0
2002.....	3.9	6.5	8.7	10.4	12.8	18.0	22.8	27.4	26.4	20.2
2001.....	3.6	6.6	8.5	9.7	13.1	18.5	23.1	25.5	26.0	20.9
2000.....	3.5	6.9	7.6	10.2	13.6	18.1	22.8	26.1	26.4	21.4
1999.....	3.4	6.0	7.6	9.6	13.9	18.7	23.2	26.5	26.8	21.1

White, male

2003.....	5.0	9.4	13.6	13.8	17.2	21.5	27.2	28.2	27.9	24.7
2002.....	5.2	9.5	13.0	14.3	15.3	20.5	25.8	30.5	30.0	25.1
2001.....	4.7	9.6	12.5	13.0	16.2	22.4	26.7	26.6	30.0	28.1
2000.....	5.0	10.4	11.2	13.7	17.9	21.6	26.1	27.7	27.7	28.7
1999.....	5.1	9.2	11.3	13.4	17.3	22.8	26.3	29.1	30.5	28.4

White, female

2003.....	1.9	3.6	4.8	6.7	9.2	14.3	21.7	24.3	23.7	16.5
2002.....	2.5	3.5	4.6	6.6	10.5	15.7	20.4	25.1	24.3	18.1
2001.....	2.4	3.8	4.6	6.6	10.2	15.1	20.2	24.7	23.5	17.9
2000.....	2.1	3.4	4.2	6.9	9.7	15.0	20.2	24.9	25.6	18.2
1999.....	1.8	2.9	3.9	6.0	10.7	15.1	20.7	24.6	24.6	18.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic liver disease and cirrhosis (K73-K74)

Black, both sexes

2003.....	3.6	7.4	13.0	14.2	12.1	14.2	14.8	11.1	10.9	10.8
2002.....	3.7	9.2	12.8	12.1	13.5	13.4	12.6	13.9	12.0	9.8
2001.....	4.7	10.2	12.1	13.6	12.2	13.9	14.8	15.3	11.6	7.7
2000.....	4.1	8.4	13.0	12.2	13.5	15.1	16.1	12.1	13.8	12.2
1999.....	4.7	9.8	10.1	13.4	15.1	14.8	14.9	15.2	13.3	12.8

Black, male

2003.....	4.3	9.5	20.9	22.7	18.1	18.0	16.4	14.6	*	*
2002.....	4.2	11.9	19.5	19.5	18.6	18.4	16.0	12.4	*	*
2001.....	6.4	15.3	19.9	20.3	19.7	20.3	18.5	18.1	*	*
2000.....	5.0	11.7	21.3	18.9	20.1	17.9	19.8	15.5	*	*
1999.....	7.0	15.3	16.4	19.2	19.1	19.2	18.7	17.6	*	*

Black, female

2003.....	2.9	5.6	6.2	7.3	7.4	11.3	13.7	9.0	10.1	8.1
2002.....	3.3	6.9	7.1	6.1	9.4	9.6	10.3	14.8	10.7	9.9
2001.....	3.2	5.8	5.5	8.1	6.2	9.2	12.3	13.6	13.1	*
2000.....	3.2	5.5	6.2	7.0	8.5	13.0	13.6	10.0	15.1	11.6
1999.....	2.6	5.0	4.8	8.7	12.0	11.5	12.4	13.7	12.1	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2003.....	0.1	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	2.0	3.0	5.7	10.3	24.5
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9

All races 3/, male

2003.....	*	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.4	0.8	1.2	2.3	3.7	7.2	12.7	26.2
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7

All races 3/, female

2003.....	*	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.9	23.8
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2003.....	*	0.2	0.3	0.4	0.8	1.8	3.6	5.9	11.3	21.9
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.9	2.8	5.6	10.2	25.3
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4

White, male

2003.....	*	*	0.4	0.5	0.9	2.1	4.4	7.5	13.9	23.0
2002.....	*	0.3	0.5	0.8	1.1	2.3	4.0	7.4	13.3	24.7
2001.....	*	0.3	0.4	0.8	1.4	1.9	4.4	7.6	12.0	26.7
2000.....	*	*	0.3	0.7	1.2	2.3	3.5	7.3	13.0	27.5
1999.....	*	0.3	0.3	0.7	1.2	1.9	4.2	7.0	13.8	28.1

White, female

2003.....	*	*	0.3	0.3	0.7	1.6	2.8	4.7	9.8	21.4
2002.....	*	*	0.3	0.4	0.9	1.7	3.1	4.1	9.0	23.0
2001.....	*	0.3	*	0.4	1.1	1.8	3.1	4.4	8.4	22.8
2000.....	*	*	0.3	0.4	0.8	1.5	2.2	4.4	8.5	24.4
1999.....	*	0.3	0.4	0.6	0.6	1.4	2.7	4.8	9.3	22.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2003.....	*	*	1.0	*	*	3.0	2.8	6.0	10.9	20.7
2002.....	*	*	*	*	*	2.3	4.6	7.7	12.8	14.8
2001.....	*	*	*	1.4	2.0	3.5	6.2	6.1	11.6	19.7
2000.....	*	*	*	*	*	3.0	4.9	6.6	11.5	17.9
1999.....	*	*	*	*	2.0	3.2	4.1	4.6	11.3	15.3

Black, male

2003.....	*	*	*	*	*	*	*	*	*	20.9
2002.....	*	*	*	*	*	*	*	*	16.0	*
2001.....	*	*	*	*	*	*	*	*	*	27.2
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2003.....	*	*	*	*	*	*	*	*	9.3	20.7
2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	5.0	6.5	12.3	20.0
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes

2003.....	2.3	3.8	6.1	10.4	17.9	30.7	50.6	84.5	143.9	293.1
2002.....	2.1	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001.....	2.1	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000.....	2.0	3.5	5.6	9.5	17.2	29.8	47.8	79.5	133.9	274.2
1999.....	2.0	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9

All races 3/, male

2003.....	2.7	4.5	7.6	12.2	19.6	34.6	58.1	102.9	187.9	387.6
2002.....	2.3	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001.....	2.3	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000.....	2.5	4.4	6.5	10.9	18.9	33.3	54.8	98.0	170.3	361.2
1999.....	2.3	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1

All races 3/, female

2003.....	1.9	3.2	4.7	8.6	16.4	27.4	44.7	71.4	117.1	251.4
2002.....	1.9	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001.....	1.9	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000.....	1.6	2.5	4.7	8.3	15.6	26.8	42.3	66.2	111.6	236.7
1999.....	1.7	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2003.....	1.5	2.6	4.3	7.8	14.1	25.2	43.5	77.6	136.0	284.8
2002.....	1.4	2.6	3.9	7.4	13.1	24.4	43.2	78.1	135.5	278.8
2001.....	1.4	2.4	4.1	7.3	12.8	25.8	43.1	75.1	127.9	278.2
2000.....	1.4	2.3	3.8	6.9	13.3	24.1	41.2	73.0	126.1	265.7
1999.....	1.3	2.0	3.5	6.6	12.2	23.0	40.4	70.9	118.5	255.7

White, male

2003.....	1.9	3.1	5.1	9.2	15.5	29.1	50.5	96.9	179.7	383.6
2002.....	1.6	3.0	4.6	8.0	14.3	28.4	53.0	99.0	177.9	377.4
2001.....	1.6	2.8	4.9	8.2	14.4	30.5	51.3	94.6	169.0	388.9
2000.....	1.7	2.9	4.4	8.0	15.1	28.3	49.3	91.8	164.2	358.8
1999.....	1.5	2.4	4.0	7.8	14.0	26.0	48.9	91.6	159.6	360.8

White, female

2003.....	1.1	2.1	3.4	6.5	12.8	21.8	37.7	63.6	109.1	241.0
2002.....	1.3	2.3	3.3	6.7	12.0	20.8	35.1	63.2	109.4	235.9
2001.....	1.2	1.9	3.3	6.3	11.4	21.7	36.4	61.3	103.1	231.9
2000.....	1.1	1.7	3.2	5.9	11.7	20.5	34.6	59.4	102.6	225.9
1999.....	1.1	1.6	3.1	5.5	10.6	20.4	33.6	56.4	94.3	213.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2003.....	7.9	12.6	21.0	34.3	54.2	83.8	127.8	171.0	260.7	440.5
2002.....	6.7	11.7	21.6	33.7	50.4	79.2	126.6	178.0	241.5	441.6
2001.....	6.7	12.5	19.8	32.5	53.1	88.6	119.8	172.3	248.6	429.6
2000.....	6.3	12.9	20.4	32.8	51.3	80.3	118.6	166.0	252.8	407.4
1999.....	7.0	11.8	19.6	29.1	47.1	77.8	117.0	163.7	246.2	433.3

Black, male

2003.....	8.8	14.8	29.3	43.3	62.7	94.1	155.7	191.7	333.7	518.2
2002.....	7.6	14.7	26.8	40.0	61.2	89.3	149.6	207.6	281.2	561.8
2001.....	7.0	16.5	22.8	37.7	55.5	101.1	130.9	189.1	309.8	511.7
2000.....	7.9	17.9	25.5	39.0	56.7	83.2	121.7	191.7	285.6	452.3
1999.....	8.2	15.5	23.4	29.2	54.5	81.8	130.6	208.0	327.0	526.1

Black, female

2003.....	7.2	10.7	13.9	26.9	47.5	76.2	109.0	158.7	222.4	410.4
2002.....	5.9	9.1	17.2	28.5	42.0	71.8	111.1	160.3	220.8	395.5
2001.....	6.4	9.1	17.1	28.2	51.2	79.4	112.3	162.3	217.5	399.1
2000.....	4.8	8.7	16.3	27.9	47.2	78.1	116.4	149.6	234.7	388.6
1999.....	5.9	8.6	16.4	29.1	41.3	74.8	108.0	137.0	206.1	398.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes

2003.....	*	*	*	*	*	*	*	0.6	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	0.4	0.4	0.9
2000.....	*	*	*	*	*	*	*	0.7	1.1
1999.....	*	*	*	*	*	0.3	0.5	0.5	0.8

All races 3/, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	*

All races 3/, female

2003.....	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	0.5	*	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races 3/, both sexes

2003.....	*	*	0.2	0.1	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.2	1.7	2.8
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	0.1	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	2.7
1999.....	*	*	0.2	0.2	0.3	0.4	0.7	1.4	2.0	3.2

All races 3/, male

2003.....	*	*	0.2	*	*	0.5	0.6	1.3	2.0	4.1
2002.....	*	*	*	0.3	*	*	0.9	1.6	2.1	3.8
2001.....	*	*	*	*	*	0.5	1.0	1.3	2.3	3.7
2000.....	*	*	*	*	*	0.6	0.8	1.5	2.0	4.3
1999.....	*	*	*	*	*	0.5	0.6	1.9	2.7	5.0

All races 3/, female

2003.....	*	*	*	*	*	*	0.8	0.9	1.4	2.2
2002.....	*	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	*	0.5	0.5	0.7	1.0	2.4
2000.....	*	*	*	*	*	*	0.6	0.8	1.8	2.1
1999.....	*	*	*	*	*	0.4	0.7	1.1	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2003.....	*	*	*	0.2	0.2	0.4	0.6	0.9	1.5	2.6
2002.....	*	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.5	2.9
2000.....	*	*	*	0.2	*	0.4	0.7	1.0	1.9	2.4
1999.....	*	*	*	*	0.3	0.3	0.6	1.4	1.8	2.9

White, male

2003.....	*	*	*	*	*	0.5	*	1.1	1.9	3.8
2002.....	*	*	*	*	*	*	0.9	1.5	2.1	3.8
2001.....	*	*	*	*	*	*	1.1	1.2	2.3	4.0
2000.....	*	*	*	*	*	*	0.8	1.4	2.1	3.8
1999.....	*	*	*	*	*	*	0.6	1.9	2.6	4.6

White, female

2003.....	*	*	*	*	*	*	0.6	0.8	1.3	2.1
2002.....	*	*	*	*	*	*	0.5	0.7	1.4	2.0
2001.....	*	*	*	*	*	*	0.5	0.7	0.9	2.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.8
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

All races 3/, both sexes

2003.....	2.2	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	1.9	3.3	5.4	9.3	16.8	29.2	46.8	78.2	131.1	270.1
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8

All races 3/, male

2003.....	2.6	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.3	4.2	6.3	10.6	18.4	32.6	53.7	96.2	167.5	355.0
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6

All races 3/, female

2003.....	1.8	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.5	2.4	4.6	8.0	15.3	26.3	41.4	65.2	108.9	233.5
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

White, both sexes

2003.....	1.4	2.5	4.1	7.6	13.8	24.7	42.7	76.4	133.9	281.5
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	13.0	23.6	40.2	71.7	123.4	262.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9

White, male

2003.....	1.8	3.0	4.9	9.0	15.3	28.4	49.7	95.6	177.3	379.3
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.6	2.7	4.3	7.7	14.7	27.7	48.2	90.0	161.3	353.3
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9

White, female

2003.....	1.0	2.1	3.3	6.4	12.4	21.4	36.9	62.4	107.1	238.2
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.7	11.4	20.1	33.8	58.5	100.0	223.0
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

Black, both sexes

2003.....	7.8	12.4	20.5	34.2	53.6	82.9	126.1	168.8	257.4	436.4
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.5	19.5	32.4	50.6	78.9	117.6	163.7	249.3	401.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3

Black, male

2003.....	8.6	14.4	28.7	43.3	61.9	92.6	153.7	188.1	328.5	513.0
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.5	24.0	38.7	55.6	81.2	121.1	189.4	283.1	443.8
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2

Black, female

2003.....	7.0	10.7	13.7	26.7	47.1	75.7	107.4	157.4	220.1	406.7
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.3	15.9	27.5	46.9	77.2	115.0	147.2	230.6	383.7
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes

2003.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.8	5.9
2002.....	*	*	0.1	0.2	0.3	0.5	0.7	1.6	2.3	6.3
2001.....	*	0.1	0.1	0.2	0.2	0.4	0.6	1.7	2.4	6.0
2000.....	*	0.1	0.1	0.2	0.3	0.4	0.8	1.5	2.7	7.4
1999.....	*	0.1	0.2	0.2	0.3	0.5	0.9	1.5	2.7	7.5

All races 3/, male

2003.....	*	*	*	*	*	*	*	0.8	2.4	5.4
2002.....	*	*	*	*	*	*	0.7	1.4	1.4	5.3
2001.....	*	*	*	*	*	*	0.5	1.2	2.1	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.1	5.3
1999.....	*	*	*	*	*	*	0.6	1.2	2.6	6.8

All races 3/, female

2003.....	*	*	*	0.3	*	0.8	0.7	1.6	3.1	6.1
2002.....	*	*	*	*	0.4	0.6	0.7	1.7	2.8	6.7
2001.....	*	*	0.2	*	*	*	0.6	2.1	2.6	6.7
2000.....	*	*	*	0.3	0.4	0.5	1.1	1.9	3.1	8.4
1999.....	*	*	0.3	*	*	0.6	1.1	1.7	2.8	7.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2003.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.7	6.0
2002.....	*	*	*	0.2	0.2	0.5	0.8	1.6	2.3	6.4
2001.....	*	*	*	*	*	0.3	0.6	1.7	2.2	6.1
2000.....	*	*	*	0.2	0.3	0.4	0.7	1.4	2.6	7.5
1999.....	*	*	0.2	0.2	0.2	0.5	0.9	1.5	2.8	7.7

White, male

2003.....	*	*	*	*	*	*	*	0.8	2.2	5.8
2002.....	*	*	*	*	*	*	0.8	1.4	1.4	5.4
2001.....	*	*	*	*	*	*	*	1.1	2.0	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.1	5.6
1999.....	*	*	*	*	*	*	*	1.3	2.6	7.2

White, female

2003.....	*	*	*	*	*	0.8	0.7	1.6	3.1	6.0
2002.....	*	*	*	*	*	0.6	0.8	1.7	2.9	6.8
2001.....	*	*	*	*	*	*	0.7	2.2	2.4	6.9
2000.....	*	*	*	*	*	0.5	1.0	1.7	2.9	8.3
1999.....	*	*	*	*	*	0.6	1.1	1.7	2.9	7.9

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Pregnancy, childbirth and the puerperium (000-099)

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*

[illegible]

2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2003.....	1.4	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.9	2.2	2.6	2.6	2.5	3.3	5.0	7.4
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8

All races 3/, male

2003.....	1.5	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.4	1.6	2.3	2.0	2.7	2.8	2.6	3.2	5.3	10.1
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5

All races 3/, female

2003.....	1.3	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.9	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2003.....	1.5	1.5	1.9	2.5	2.5	2.3	2.3	3.1	4.5	6.9
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.7	2.5	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

White, male

2003.....	1.6	1.5	2.0	2.6	2.6	2.3	2.3	2.8	4.3	6.9
2002.....	1.4	1.8	2.0	2.3	2.8	2.6	2.7	3.4	4.6	7.3
2001.....	1.6	1.5	1.9	2.2	2.6	2.7	3.2	3.9	4.6	7.5
2000.....	1.4	1.6	2.3	2.1	2.8	2.6	2.6	3.4	5.5	10.6
1999.....	1.3	1.9	2.0	2.6	2.6	2.5	2.6	3.0	5.8	8.6

White, female

2003.....	1.3	1.4	1.9	2.4	2.4	2.2	2.2	3.3	4.7	6.9
2002.....	1.2	1.5	2.0	2.4	2.2	2.4	2.4	3.0	5.0	6.7
2001.....	1.3	1.4	1.9	2.2	2.8	2.1	2.3	3.1	4.2	7.3
2000.....	1.1	1.4	1.7	2.4	2.5	2.7	2.4	3.4	5.0	6.5
1999.....	1.2	1.7	1.7	2.3	2.3	2.9	2.3	3.6	4.6	7.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2003.....	5.9	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.5	8.7	8.2	8.7	10.3	14.4	19.3	30.2	57.0	205.4
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All races 3/, male

2003.....	7.4	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.1	11.8	11.5	11.6	13.6	19.2	22.6	35.5	61.4	173.4
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All races 3/, female

2003.....	4.4	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.1	5.9	7.4	10.2	16.7	26.3	54.3	219.2
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2003.....	5.6	6.6	6.2	6.3	7.6	10.1	16.6	26.5	56.1	228.3
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.3	8.2	7.4	7.9	9.6	13.1	18.7	29.5	56.4	209.9
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2

White, male

2003.....	7.0	8.6	8.2	8.1	9.7	12.8	20.4	31.0	57.8	180.0
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.0	11.2	10.3	10.3	12.6	17.4	21.8	34.8	61.1	178.2
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3

White, female

2003.....	4.2	4.6	4.2	4.7	5.6	7.7	13.5	23.3	55.0	249.7
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.7	6.8	9.5	16.3	25.7	53.5	223.4
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2003.....	9.7	13.4	13.4	13.0	15.7	17.9	30.3	34.9	60.9	210.9
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	14.0	16.5	16.2	19.2	27.2	29.5	44.4	76.6	186.9
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4

Black, male

2003.....	12.5	17.5	17.8	18.5	22.8	25.9	38.8	41.5	70.1	193.9
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.8	25.0	26.1	26.6	39.6	36.3	52.8	82.2	149.0
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0

Black, female

2003.....	7.3	9.9	9.7	8.4	10.2	12.1	24.6	30.9	56.1	217.5
2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	10.0	9.5	8.6	13.7	18.0	24.6	39.0	73.5	202.8
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,
N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2003.....	23.1	33.3	45.3	56.6	80.9	114.5	178.3	303.7	559.8	1,508.4
2002.....	23.5	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001.....	21.7	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000.....	21.9	31.3	41.1	53.7	75.1	107.2	170.4	289.1	542.5	1,426.8
1999.....	21.6	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7

All races 3/, male

2003.....	27.8	40.3	55.2	65.6	91.0	125.7	191.0	316.5	563.7	1,258.1
2002.....	28.2	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001.....	25.7	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000.....	26.1	39.0	48.8	62.0	84.7	119.4	184.6	298.0	529.4	1,201.4
1999.....	26.6	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4

All races 3/, female

2003.....	18.5	26.5	36.0	48.1	71.7	104.8	168.0	294.7	557.4	1,619.1
2002.....	18.9	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001.....	17.7	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000.....	17.8	23.9	33.8	46.0	66.4	96.9	159.2	282.7	550.5	1,523.9
1999.....	16.6	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,
N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2003.....	21.4	30.8	41.6	53.5	77.1	111.9	175.4	301.8	562.0	1,534.2
2002.....	21.6	30.9	39.2	53.1	73.4	107.7	170.2	295.6	556.2	1,517.6
2001.....	19.3	28.7	37.3	49.5	72.6	108.9	169.2	292.6	541.2	1,498.0
2000.....	19.5	27.5	37.6	49.7	70.8	104.0	166.8	285.8	540.6	1,448.3
1999.....	19.1	26.3	34.8	47.4	68.6	103.2	163.6	277.1	512.1	1,434.0

White, male

2003.....	26.4	38.0	50.8	62.0	86.2	121.6	186.0	311.4	561.0	1,279.1
2002.....	26.8	38.2	48.7	60.6	83.4	118.6	181.2	304.6	553.0	1,274.7
2001.....	23.4	35.3	45.1	55.3	81.8	119.9	182.1	302.0	540.6	1,282.9
2000.....	23.7	35.3	44.9	57.5	78.6	114.8	179.4	292.9	525.4	1,215.9
1999.....	24.3	32.5	41.6	55.4	76.8	111.8	179.1	290.9	525.5	1,231.6

White, female

2003.....	16.3	23.6	32.6	45.3	68.6	103.4	166.8	294.9	562.7	1,647.2
2002.....	16.3	23.6	29.9	46.0	64.2	98.1	161.2	289.2	558.2	1,623.1
2001.....	15.3	22.3	29.7	44.0	64.2	99.2	158.8	285.9	541.6	1,588.0
2000.....	15.3	19.8	30.6	42.4	63.8	94.6	156.7	280.6	549.9	1,547.7
1999.....	13.8	20.2	28.3	39.9	61.0	95.7	151.2	267.4	504.2	1,515.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,
N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2003.....	38.7	58.2	83.5	96.1	131.7	161.7	241.2	382.2	626.5	1,435.7
2002.....	41.6	60.3	78.4	96.1	123.7	155.7	242.8	369.7	642.1	1,425.4
2001.....	41.2	63.0	79.4	95.6	122.1	163.6	237.0	380.4	612.1	1,418.0
2000.....	41.8	65.3	78.2	96.8	123.6	151.2	231.8	373.8	645.8	1,392.5
1999.....	42.5	61.8	75.4	92.1	117.2	167.2	239.7	369.5	556.5	1,286.2

Black, male

2003.....	42.0	65.9	102.4	113.0	158.7	191.6	289.7	441.8	710.2	1,224.1
2002.....	44.2	70.9	94.6	116.3	146.1	185.2	288.4	419.3	693.1	1,319.8
2001.....	47.0	74.8	95.9	114.8	141.6	196.9	273.3	445.6	659.7	1,313.9
2000.....	47.9	76.8	94.5	119.5	156.9	180.4	266.7	412.9	682.4	1,222.8
1999.....	47.6	73.3	89.5	111.0	134.9	189.5	274.1	434.1	635.7	1,291.3

Black, female

2003.....	35.7	51.5	67.6	82.2	110.7	139.6	208.5	346.7	582.6	1,517.9
2002.....	39.3	51.0	64.6	79.6	106.1	133.8	212.2	340.0	615.5	1,465.9
2001.....	36.2	52.9	65.3	79.8	106.7	138.9	212.8	341.4	587.9	1,456.6
2000.....	36.5	55.6	64.9	79.2	98.8	129.4	207.0	348.7	625.7	1,463.4
1999.....	37.9	52.0	63.5	76.6	103.3	150.6	216.9	330.7	517.3	1,284.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2003.....	40.4	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.1	34.5	31.3	30.2	32.7	35.8	49.5	75.5	125.7	269.9
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4

All races 3/, male

2003.....	58.0	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	52.6	52.2	47.2	43.9	46.2	48.9	65.1	98.0	162.3	341.2
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4

All races 3/, female

2003.....	23.0	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	17.9	17.5	16.3	17.6	20.5	24.7	37.0	59.3	103.3	239.2
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2003.....	41.5	41.0	35.7	32.1	32.5	37.3	50.9	80.2	136.5	291.1
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	34.7	33.4	29.6	29.1	32.1	35.0	48.7	75.7	128.1	279.6
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4

White, male

2003.....	59.6	58.6	50.6	45.6	46.1	50.8	66.9	105.8	177.3	365.7
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	51.8	50.2	44.0	41.3	44.4	46.7	64.0	96.9	164.0	352.7
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6

White, female

2003.....	23.4	23.4	21.1	19.2	19.9	25.4	37.9	61.7	111.3	258.0
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.6	16.9	15.7	17.6	20.9	24.9	36.4	60.3	106.0	248.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2003.....	41.7	47.9	49.1	43.7	40.4	47.7	53.8	67.7	105.1	160.2
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.7	47.8	48.6	42.6	39.8	44.3	59.6	75.6	104.3	181.6
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6

Black, male

2003.....	60.8	73.9	75.2	70.1	64.8	74.4	74.2	105.0	152.8	225.2
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.6	76.6	81.0	74.6	68.4	71.0	82.2	115.2	153.6	254.3
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4

Black, female

2003.....	24.8	25.3	27.0	22.0	21.4	28.0	40.1	45.6	80.1	134.9
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.2	22.2	17.7	18.5	24.4	43.6	50.1	77.2	151.2
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2003.....	17.5	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.7	16.3	15.7	15.9	16.6	17.0	19.6	25.1	30.2	31.1
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0

All races 3/, male

2003.....	25.5	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.6	23.9	22.9	22.8	23.0	23.0	26.5	33.9	43.4	56.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5

All races 3/, female

2003.....	9.5	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	8.9	9.0	9.0	9.5	10.8	11.8	14.2	18.8	22.2	20.0
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transport accidents (V01-V99,Y85)

White, both sexes

2003.....	17.7	16.9	16.1	15.7	15.8	16.0	19.2	23.9	30.0	30.5
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.6	16.1	15.3	15.7	16.6	16.9	19.5	25.3	30.9	31.9
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3

White, male

2003.....	25.8	24.9	23.3	22.8	22.6	22.2	25.4	32.7	42.6	52.8
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.3	23.5	22.1	22.0	22.5	22.3	26.1	33.7	43.8	58.8
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7

White, female

2003.....	9.6	9.1	9.1	8.8	9.6	10.5	14.0	17.6	22.2	20.6
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	8.9	8.8	8.7	9.7	11.2	12.1	14.1	19.2	23.0	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transport accidents (V01-V99,Y85)

Black, both sexes

2003.....	18.7	18.1	18.2	18.9	17.8	18.5	17.9	22.8	21.8	21.0
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.7	20.0	18.4	16.6	17.5	21.2	19.4	21.8	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7

Black, male

2003.....	28.3	29.1	28.9	30.3	27.6	28.6	24.0	37.4	37.7	42.7
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	30.9	30.1	31.8	32.3	28.8	29.9	31.9	33.2	39.0	41.5
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0

Black, female

2003.....	10.3	8.4	9.2	9.6	10.2	11.0	13.7	14.1	13.5	12.6
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.3	7.5	7.6	8.3	13.6	10.6	12.3	13.3
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2003.....	16.0	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.0	14.8	14.1	14.4	14.9	15.2	18.3	23.8	28.7	30.0
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All races 3/, male

2003.....	23.1	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.7	21.4	20.1	20.2	20.1	20.1	24.1	31.6	40.7	54.6
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All races 3/, female

2003.....	9.0	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.3	8.3	8.3	9.0	10.2	11.1	13.8	18.2	21.4	19.4
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2003.....	16.2	15.2	14.4	13.8	14.0	14.6	17.8	22.6	28.8	29.5
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	14.8	14.6	13.5	14.2	14.9	15.1	18.2	24.0	29.4	30.7
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7

White, male

2003.....	23.2	22.1	20.5	19.7	19.4	19.9	23.0	30.3	40.6	50.7
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.3	21.1	19.3	19.5	19.4	19.4	23.7	31.4	41.0	56.4
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3

White, female

2003.....	9.0	8.4	8.6	8.1	9.1	10.0	13.5	17.1	21.5	20.1
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.3	8.2	8.0	9.1	10.7	11.4	13.7	18.6	22.2	19.8
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2003.....	17.5	16.2	16.8	17.5	16.4	16.9	17.2	21.3	20.6	20.7
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.9	18.9	16.9	15.6	15.9	20.0	18.2	20.9	20.4
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5

Black, male

2003.....	26.4	25.8	26.0	27.6	24.5	25.1	23.0	35.1	34.7	41.7
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.3	27.0	29.8	29.4	26.8	26.7	30.0	31.0	38.2	40.4
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Black, female

2003.....	9.7	7.9	9.1	9.2	10.1	10.8	13.3	13.0	13.2	12.6
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.1	10.0	7.2	7.3	8.0	12.9	10.0	11.4	12.0
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2003.....	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.6	0.7	0.5
2002.....	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	*
2001.....	0.6	0.7	0.6	0.5	0.6	0.7	0.6	0.8	0.8	0.7
2000.....	0.7	0.6	0.6	0.5	0.5	0.6	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.3	1.4	1.3

All races 3/, male

2003.....	0.9	1.1	0.8	0.9	0.8	0.8	0.7	0.8	1.3	*
2002.....	0.9	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.2	*
2001.....	1.0	1.0	0.9	0.8	0.9	1.1	0.9	1.2	*	*
2000.....	1.2	0.9	0.9	0.8	1.0	0.9	1.0	1.1	1.3	*
1999.....	1.5	1.4	1.2	1.2	1.1	1.2	1.2	1.7	2.1	2.9

All races 3/, female

2003.....	0.3	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.4	0.2	*	*	*	*	0.5	0.7	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2003.....	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.6	*
2001.....	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.8	0.7	0.6
2000.....	0.7	0.6	0.6	0.4	0.6	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.2	1.4	1.3

White, male

2003.....	0.9	0.9	0.8	0.9	0.8	0.7	0.7	0.8	1.2	*
2002.....	0.8	0.9	0.8	0.6	0.7	0.5	0.7	*	1.3	*
2001.....	0.9	0.9	0.9	0.8	0.9	1.0	0.9	1.3	*	*
2000.....	1.2	0.8	0.9	0.6	1.0	0.8	0.9	1.1	1.3	*
1999.....	1.5	1.3	1.2	1.3	1.1	1.2	1.1	1.7	2.0	2.8

White, female

2003.....	0.2	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.5	0.3	0.4	0.4	0.5	*	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2003.....	0.9	1.1	1.1	1.2	1.2	0.9	0.8	0.8	0.5	*
2002.....	0.9	1.1	1.0	1.1	1.1	1.0	0.8	0.8	0.7	0.4
2001.....	1.2	1.1	1.2	1.0	1.1	1.0	0.9	0.8	0.7	0.5
2000.....	1.0	0.9	1.1	1.0	1.1	1.2	0.8	0.6	0.7	*
1999.....	0.9	1.0	1.0	1.1	1.1	0.9	0.8	0.5	0.6	*

All races 3/, male

2003.....	1.6	1.7	2.0	2.0	2.3	1.7	1.5	1.6	*	*
2002.....	1.5	1.9	1.6	1.9	1.9	2.0	1.7	1.4	1.5	*
2001.....	1.8	1.8	1.9	1.7	1.9	1.8	1.7	1.3	1.4	*
2000.....	1.7	1.6	1.9	1.8	1.9	2.0	1.4	1.2	1.3	*
1999.....	1.5	1.7	1.8	2.0	1.9	1.6	1.6	0.9	*	*

All races 3/, female

2003.....	0.3	0.4	0.3	0.4	*	*	*	*	*	*
2002.....	0.3	0.3	0.3	0.3	0.4	*	*	*	*	*
2001.....	0.5	0.4	0.4	0.4	*	*	*	0.5	*	*
2000.....	0.3	0.3	0.3	*	0.4	0.5	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2003.....	1.0	1.1	1.2	1.3	1.3	0.9	0.9	0.8	0.5	*
2002.....	1.0	1.2	1.0	1.1	1.2	1.0	0.9	0.8	0.7	*
2001.....	1.3	1.1	1.2	1.0	1.2	1.0	1.0	0.8	0.7	*
2000.....	1.1	1.0	1.1	1.1	1.2	1.2	0.8	0.6	0.7	*
1999.....	1.0	1.1	1.1	1.1	1.1	0.9	0.9	0.5	0.5	*

White, male

2003.....	1.7	1.9	2.1	2.2	2.5	1.6	1.7	1.6	*	*
2002.....	1.6	2.1	1.7	1.9	2.1	2.0	1.8	1.3	1.5	*
2001.....	2.0	1.8	2.0	1.6	2.1	1.8	1.8	1.3	1.5	*
2000.....	1.8	1.6	2.0	1.9	2.0	2.1	1.5	1.2	1.5	*
1999.....	1.5	1.8	1.8	2.0	1.9	1.6	1.6	0.9	*	*

White, female

2003.....	0.4	0.4	0.4	0.4	*	*	*	*	*	*
2002.....	0.3	0.2	0.4	0.4	0.4	*	*	*	*	*
2001.....	0.6	0.4	0.5	0.5	*	*	*	*	*	*
2000.....	0.4	0.3	0.3	*	0.4	*	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

2003.....	22.9	23.9	20.3	17.0	16.9	21.9	31.9	54.9	104.1	249.2
2002.....	22.3	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001.....	19.4	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000.....	18.4	18.2	15.6	14.3	16.1	18.8	29.8	50.4	95.5	238.8
1999.....	18.8	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3

All races 3/, male

2003.....	32.4	34.1	28.7	24.0	24.3	29.7	41.8	72.0	133.0	300.5
2002.....	32.3	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001.....	28.7	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000.....	28.1	28.3	24.3	21.1	23.2	25.9	38.7	64.2	118.9	284.5
1999.....	29.1	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9

All races 3/, female

2003.....	13.5	14.1	12.3	10.4	10.2	15.1	23.9	42.7	86.4	226.5
2002.....	12.5	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001.....	10.2	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000.....	9.0	8.5	7.4	8.1	9.7	12.9	22.8	40.5	81.1	219.2
1999.....	8.7	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nontransport accidents (W00-X59,Y86)

White, both sexes

2003.....	23.8	24.0	19.5	16.5	16.7	21.4	31.8	56.2	106.5	260.6
2002.....	23.1	21.6	17.7	14.7	15.5	20.8	30.5	55.6	104.3	256.5
2001.....	19.3	19.0	15.4	13.4	15.3	20.0	29.4	54.6	102.1	255.7
2000.....	18.1	17.3	14.3	13.4	15.5	18.2	29.3	50.4	97.1	247.8
1999.....	18.5	17.1	13.2	12.9	14.2	19.7	28.7	51.9	102.6	258.1

White, male

2003.....	33.8	33.8	27.2	22.8	23.4	28.6	41.5	73.1	134.7	312.9
2002.....	33.3	31.1	25.2	20.9	21.7	28.2	41.1	72.1	130.1	302.8
2001.....	28.7	28.6	22.4	19.0	21.1	27.1	38.1	72.0	128.6	314.0
2000.....	27.5	26.6	21.9	19.3	21.9	24.4	37.9	63.3	120.2	294.0
1999.....	28.5	26.6	20.0	18.6	20.9	26.5	38.6	64.8	135.0	319.8

White, female

2003.....	13.8	14.3	12.0	10.4	10.4	14.9	23.9	44.0	89.1	237.4
2002.....	12.8	12.1	10.4	8.8	9.7	14.3	21.9	43.8	88.5	236.3
2001.....	9.7	9.4	8.5	8.1	10.0	13.8	22.3	42.2	86.1	231.2
2000.....	8.6	8.1	7.0	7.9	9.6	12.7	22.3	41.1	83.0	228.0
1999.....	8.5	7.8	6.5	7.4	8.0	13.7	20.8	42.9	83.5	233.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2003.....	22.9	29.9	30.9	24.8	22.6	29.3	36.0	44.9	83.3	139.2
2002.....	22.9	29.4	28.6	25.2	23.2	29.9	35.0	54.8	72.7	155.0
2001.....	24.9	29.6	28.0	22.8	25.1	26.9	39.0	56.1	84.8	166.1
2000.....	24.4	28.1	28.7	24.2	23.2	26.8	38.5	56.2	82.5	159.9
1999.....	25.6	30.7	26.2	24.6	28.6	32.7	45.6	65.7	95.9	179.8

Black, male

2003.....	32.4	44.8	46.3	39.8	37.2	45.8	50.3	67.6	115.2	182.5
2002.....	32.7	42.9	46.3	42.1	38.0	47.1	50.1	79.5	105.5	214.4
2001.....	36.0	47.8	46.1	37.2	38.8	40.9	56.2	80.4	133.2	220.1
2000.....	37.7	46.5	49.3	42.3	39.6	41.1	50.3	81.9	114.6	212.8
1999.....	40.7	51.5	44.1	42.2	42.2	50.6	70.2	95.0	128.2	248.4

Black, female

2003.....	14.5	16.8	17.8	12.5	11.1	17.0	26.4	31.5	66.5	122.3
2002.....	14.2	17.7	13.6	11.2	11.6	17.2	24.9	40.1	55.7	132.3
2001.....	15.2	13.8	12.5	11.0	14.2	16.5	27.6	41.6	60.1	146.1
2000.....	12.7	12.4	11.9	10.2	11.0	16.1	30.1	39.6	64.9	137.8
1999.....	12.3	12.9	11.0	10.2	18.0	19.4	29.3	48.1	79.9	154.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

All races 3/, both sexes

2003.....	1.7	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.1	2.7	3.4	4.7	6.3	12.2	22.1	44.9	111.1
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2

All races 3/, male

2003.....	2.7	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.6	3.4	4.3	5.1	7.0	8.7	15.8	28.9	56.5	138.0
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5

All races 3/, female

2003.....	0.6	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.3	9.3	17.3	37.8	99.5
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

White, both sexes

2003.....	1.7	2.3	2.7	3.7	5.4	8.2	15.2	30.1	60.0	144.4
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.6	3.3	4.7	6.3	12.5	23.1	46.9	117.8
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9

White, male

2003.....	2.8	3.6	4.1	5.3	7.3	10.6	19.5	38.6	76.4	175.1
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	4.0	4.8	7.0	8.6	16.3	29.8	58.8	145.7
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1

White, female

2003.....	0.6	1.0	1.3	2.1	3.6	6.1	11.6	24.0	49.9	130.8
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.4	18.3	39.6	105.8
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

Black, both sexes

2003.....	1.5	3.0	3.6	4.3	5.5	7.6	10.6	13.4	28.2	44.1
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.7	4.5	5.2	6.8	9.7	11.6	21.2	38.9
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3

Black, male

2003.....	2.6	4.8	5.8	6.5	8.1	12.8	12.2	18.3	35.4	56.3
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.5	6.7	8.1	9.1	10.2	11.8	17.8	25.7	56.4
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6

Black, female

2003.....	*	1.4	1.7	2.5	3.5	3.7	9.5	10.6	24.4	39.3
2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.3	8.1	7.6	18.7	31.6
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2003.....	1.1	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6

All races 3/, male

2003.....	1.8	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.5	1.8	1.2	2.7	2.6
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3

All races 3/, female

2003.....	0.3	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.2	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental drowning and submersion (W65-W74)

White, both sexes

2003.....	1.0	1.0	1.0	1.0	0.9	1.1	1.1	1.2	1.3	1.6
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.9	1.1	0.9	1.8	1.4
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6

White, male

2003.....	1.8	1.5	1.5	1.6	1.4	1.7	1.9	1.8	2.4	3.3
2002.....	1.9	1.4	1.4	1.3	1.2	1.4	1.6	1.8	2.4	3.7
2001.....	1.6	1.7	1.3	1.1	1.3	1.4	1.6	1.8	2.3	3.5
2000.....	1.6	1.5	1.5	1.6	1.6	1.5	1.7	1.1	2.9	2.6
1999.....	1.7	1.8	1.7	1.4	1.4	1.4	1.5	1.9	2.7	2.4

White, female

2003.....	0.3	0.4	0.4	0.4	0.4	0.6	0.6	0.8	0.7	0.9
2002.....	0.4	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001.....	0.4	0.3	0.3	0.4	0.5	*	*	0.7	0.7	1.0
2000.....	0.3	0.3	0.3	*	*	*	0.6	0.7	1.1	1.0
1999.....	0.4	0.4	0.3	0.4	*	*	0.7	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

[illegible]

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2003.....	1.0	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	0.9	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.4	6.1
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9

All races 3/, male

2003.....	1.2	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.9	2.0	2.3	2.5	3.2	4.4	5.8	8.9
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3

All races 3/, female

2003.....	0.7	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.8	0.9	1.2	1.6	1.8	2.8	3.5	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, both sexes

2003.....	1.0	1.2	1.2	1.2	1.4	1.8	2.2	2.6	3.7	4.7
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.8	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7

White, male

2003.....	1.3	1.5	1.5	1.6	1.9	2.2	3.0	3.2	4.2	6.7
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.0	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	7.8
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0

White, female

2003.....	0.7	0.8	0.9	0.8	1.0	1.5	1.6	2.2	3.4	3.8
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2003.....	1.4	2.3	2.9	2.8	2.9	4.4	6.4	8.7	12.4	14.9
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.7	4.0	4.5	4.8	7.0	11.4	16.8	20.4
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7

Black, male

2003.....	1.5	3.3	3.2	4.6	3.9	6.4	8.9	15.5	17.0	*
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.3	4.2	6.0	7.4	7.2	9.9	15.9	21.6	25.5
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3

Black, female

2003.....	*	1.4	2.6	*	*	*	4.7	*	10.1	13.4
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	5.0	8.5	14.2	18.2
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2003.....	15.5	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.1	10.0	6.2	3.4	2.3	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4

All races 3/, male

2003.....	20.7	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.3	14.9	9.1	4.1	2.8	1.8	1.8	1.6	2.8	5.9
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4

All races 3/, female

2003.....	10.3	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	5.9	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2003.....	16.3	15.7	10.6	5.8	3.4	2.0	2.0	2.0	2.7	3.5
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	10.9	9.4	5.4	3.0	2.1	1.5	1.4	1.7	2.7	4.0
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3

White, male

2003.....	21.8	20.9	13.8	7.0	4.2	2.3	2.2	2.3	3.6	3.7
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.0	13.8	7.7	3.4	2.4	1.6	1.6	1.6	2.9	5.6
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5

White, female

2003.....	10.8	10.5	7.3	4.7	2.6	1.7	1.8	1.8	2.1	3.4
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.7	5.1	3.1	2.8	1.9	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2003.....	15.2	18.8	16.9	9.6	4.8	5.5	*	*	*	*
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.7	14.7	6.9	4.0	2.5	*	*	*	8.2
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*

Black, male

2003.....	20.5	26.7	25.7	15.8	8.7	10.3	*	*	*	*
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	27.0	24.5	12.0	7.4	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*

Black, female

2003.....	10.5	11.9	9.5	4.5	*	*	*	*	*	*
2002.....	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001.....	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000.....	8.6	8.0	6.7	2.9	*	*	*	*	*	*
1999.....	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2003.....	3.4	3.8	4.0	4.7	5.5	8.1	11.2	19.8	38.1	102.5
2002.....	3.3	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001.....	3.2	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000.....	3.4	4.0	4.2	5.0	6.4	7.9	12.4	22.0	41.7	115.6
1999.....	3.7	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9

All races 3/, male

2003.....	5.5	6.0	6.1	7.1	8.7	11.4	15.1	26.5	47.6	120.2
2002.....	5.4	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001.....	5.3	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000.....	5.6	6.6	6.8	7.9	9.3	10.9	15.7	27.6	50.8	128.3
1999.....	6.1	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7

All races 3/, female

2003.....	1.4	1.6	2.1	2.5	2.7	5.3	8.2	15.0	32.3	94.7
2002.....	1.3	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001.....	1.2	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000.....	1.3	1.4	1.7	2.4	3.8	5.3	9.7	18.0	36.2	110.1
1999.....	1.4	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2003.....	3.5	3.7	3.9	4.6	5.4	8.0	11.1	20.0	38.4	106.3
2002.....	3.4	3.5	4.0	4.6	5.6	7.9	11.0	21.4	39.1	108.8
2001.....	3.2	3.8	4.4	4.5	5.9	7.7	11.9	22.3	42.5	117.7
2000.....	3.4	3.7	4.0	4.8	6.2	7.5	12.0	21.7	42.2	119.2
1999.....	3.7	3.6	4.2	4.7	5.8	8.6	12.7	22.8	46.8	132.4

White, male

2003.....	5.7	5.8	5.9	6.9	8.2	11.4	14.7	26.6	47.3	123.9
2002.....	5.6	5.9	6.1	6.9	8.5	10.9	14.9	28.1	47.2	121.2
2001.....	5.2	6.2	6.6	6.5	8.5	10.7	15.3	29.1	51.9	137.0
2000.....	5.5	6.2	6.5	7.4	8.8	10.3	15.3	26.6	50.4	131.3
1999.....	6.0	5.5	6.8	7.1	8.5	11.3	17.0	28.7	59.0	155.2

White, female

2003.....	1.3	1.6	1.9	2.4	2.7	5.0	8.1	15.3	32.9	98.5
2002.....	1.2	1.1	1.9	2.4	2.9	5.3	7.8	16.5	34.2	103.4
2001.....	1.1	1.5	2.3	2.6	3.5	5.1	9.1	17.4	36.8	109.7
2000.....	1.3	1.4	1.6	2.3	3.9	5.1	9.3	18.1	37.1	114.0
1999.....	1.5	1.6	1.7	2.4	3.3	6.2	9.3	18.7	39.6	123.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2003.....	3.7	4.7	6.2	6.7	8.0	11.2	14.7	19.6	39.6	73.2
2002.....	3.3	4.3	5.1	7.3	7.3	15.9	14.3	27.8	39.0	83.1
2001.....	3.9	5.4	5.8	5.9	10.4	10.4	18.7	27.3	43.9	96.1
2000.....	3.9	5.8	5.8	8.0	8.7	11.8	18.0	30.3	41.0	90.3
1999.....	4.7	6.6	6.6	9.3	10.9	13.9	22.9	36.2	53.7	105.5

Black, male

2003.....	5.6	7.9	9.2	10.4	13.9	15.3	22.3	29.2	56.8	91.8
2002.....	4.3	6.4	8.6	12.7	11.3	23.7	18.7	40.9	53.5	106.1
2001.....	6.6	8.6	9.5	9.9	14.6	15.8	25.2	35.3	72.2	114.6
2000.....	6.5	10.4	10.6	14.4	14.5	17.4	22.0	45.1	63.1	114.9
1999.....	8.1	11.1	11.0	15.4	16.1	20.8	34.9	50.3	69.8	153.5

Black, female

2003.....	2.0	2.0	3.7	3.7	3.5	8.2	9.5	13.8	30.6	66.0
2002.....	2.4	2.5	2.1	2.9	4.2	10.2	11.4	20.0	31.4	74.3
2001.....	1.6	2.7	2.7	2.7	7.0	6.4	14.3	22.5	29.4	89.3
2000.....	1.5	1.8	*	2.9	4.4	7.6	15.1	20.8	28.8	80.0
1999.....	1.6	2.7	2.8	4.3	6.8	8.9	14.9	27.7	45.8	87.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2003.....	15.9	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 4/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.3	15.0	14.2	12.8	11.6	11.4	13.9	16.6	19.5	19.4
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All races 3/, male

2003.....	24.5	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 4/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.7	23.6	21.9	20.4	19.0	20.1	26.3	34.3	44.1	54.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All races 3/, female

2003.....	7.4	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 4/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.9	5.8	4.9	4.0	4.1	3.8	4.3	4.3
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2003.....	17.9	18.1	17.7	16.2	13.8	13.0	14.7	16.7	18.9	18.1
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 4/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	16.8	16.6	15.5	14.0	12.8	12.5	14.9	17.6	20.6	20.8
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7

White, male

2003.....	27.3	27.3	27.2	25.4	22.9	22.9	27.9	34.0	42.9	51.4
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 4/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	25.9	25.8	23.6	22.0	20.7	21.7	27.9	36.5	47.0	58.8
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5

White, female

2003.....	8.4	9.0	8.3	7.4	5.5	4.3	3.9	4.2	4.1	3.3
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 4/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.8	7.6	7.6	6.4	5.5	4.4	4.4	4.0	4.3	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2003.....	7.2	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.5	7.9	7.5	7.8	8.4	10.4	12.4	14.2	13.0
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6

All races 3/, male

2003.....	12.1	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.5	12.9	13.6	13.4	14.2	16.1	21.4	27.6	34.8	40.7
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6

All races 3/, female

2003.....	2.3	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.3	2.5	2.1	2.1	1.8	1.6	1.4	1.5	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2003.....	8.3	9.1	9.4	9.5	9.1	9.5	11.1	12.9	14.1	12.9
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.3	8.4	8.8	8.2	8.6	9.3	11.2	13.4	15.3	14.1
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6

White, male

2003.....	13.8	15.1	16.2	16.4	16.8	18.3	22.5	28.3	34.8	39.9
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	13.9	14.3	15.0	14.4	15.5	17.6	22.9	29.9	37.6	44.3
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9

White, female

2003.....	2.7	3.1	2.6	2.9	2.0	1.8	1.8	1.8	1.4	0.9
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2003.....	8.7	8.2	7.5	6.2	4.5	3.4	3.5	3.7	4.6	5.1
2002.....	8.7	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 4/.....	8.2	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000.....	7.9	7.5	6.3	5.3	3.8	3.0	3.5	4.2	5.3	6.4
1999.....	7.7	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7

All races 3/, male

2003.....	12.4	11.2	9.9	8.5	5.8	4.7	5.4	5.5	7.7	11.1
2002.....	12.1	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 4/.....	11.6	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000.....	11.2	10.7	8.3	7.1	4.9	4.0	4.9	6.6	9.3	13.8
1999.....	10.8	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5

All races 3/, female

2003.....	5.1	5.3	5.1	4.1	3.2	2.4	2.0	2.4	2.6	2.4
2002.....	5.4	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 4/.....	4.8	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000.....	4.6	4.5	4.4	3.7	2.8	2.2	2.5	2.4	2.9	3.2
1999.....	4.5	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2003.....	9.6	9.1	8.3	6.7	4.7	3.5	3.6	3.8	4.8	5.2
2002.....	9.7	9.6	7.9	6.4	4.6	3.8	4.0	4.3	5.3	6.2
2001 4/.....	9.1	9.1	7.2	5.9	4.2	3.8	3.7	4.2	5.0	5.5
2000.....	8.5	8.3	6.7	5.8	4.1	3.2	3.7	4.2	5.3	6.7
1999.....	8.4	7.6	6.7	5.1	4.1	3.8	3.8	4.7	5.9	7.1

White, male

2003.....	13.5	12.2	11.0	9.0	6.1	4.7	5.4	5.6	8.1	11.5
2002.....	13.4	12.6	10.9	8.7	6.2	5.0	5.8	6.7	8.4	13.4
2001 4/.....	12.7	12.6	9.3	7.9	5.3	5.5	5.3	6.3	9.0	12.9
2000.....	12.0	11.5	8.6	7.6	5.2	4.1	5.0	6.6	9.4	14.5
1999.....	11.9	10.4	8.6	6.7	5.2	5.3	5.5	6.7	10.0	16.6

White, female

2003.....	5.7	5.9	5.7	4.5	3.5	2.5	2.2	2.4	2.8	2.4
2002.....	6.0	6.5	5.0	4.2	3.0	2.7	2.5	2.6	3.4	3.1
2001 4/.....	5.4	5.6	5.1	4.1	3.1	2.3	2.5	2.8	2.5	2.3
2000.....	5.1	5.0	4.8	4.0	3.2	2.3	2.6	2.5	2.8	3.3
1999.....	4.9	4.7	4.9	3.5	3.0	2.6	2.4	3.3	3.5	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races 3/, both sexes

2003.....	6.4	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 4/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.7	2.5	2.4	2.3	2.6	2.3
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4

All races 3/, male

2003.....	9.5	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 4/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.9	6.1	5.3	3.9	3.4	3.4	2.7	3.7	3.1
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4

All races 3/, female

2003.....	3.4	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 4/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.5	2.8	2.1	1.7	1.5	1.7	1.6	2.0	2.0	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2003.....	4.2	3.9	3.1	2.4	2.0	1.9	1.9	2.2	2.1	2.0
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 4/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.1	1.8	2.0	2.1	2.0	2.1
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1

White, male

2003.....	5.8	5.4	4.5	3.3	2.9	2.7	2.6	2.9	2.5	3.0
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 4/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.3	5.1	4.1	4.0	2.9	2.3	2.6	2.4	2.5	2.7
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7

White, female

2003.....	2.7	2.4	1.7	1.6	1.0	1.2	1.3	1.6	1.8	1.6
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 4/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.4	2.3	1.8	1.4	1.3	1.5	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2003.....	21.4	17.7	12.7	9.1	6.8	5.7	6.9	6.6	6.6	*
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 4/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	17.1	12.8	9.3	7.5	7.9	6.7	5.4	10.0	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*

Black, male

2003.....	36.8	30.0	21.8	17.3	11.4	10.8	11.5	12.3	*	*
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 4/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	29.5	23.8	16.7	13.6	14.5	11.8	*	19.1	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*

Black, female

2003.....	7.9	7.0	5.0	2.3	3.2	*	*	*	*	*
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 4/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2003.....	3.8	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.7	2.9	2.2	1.8	1.2	1.1	1.0	0.7	0.6	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All races 3/, male

2003.....	5.9	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.7	4.6	3.4	2.7	1.9	1.3	1.3	0.7	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All races 3/, female

2003.....	1.7	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.4	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2003.....	2.4	2.2	1.7	1.2	1.0	0.7	0.8	0.6	0.6	0.5
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.6	1.4	1.0	0.9	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

White, male

2003.....	3.4	3.1	2.5	1.8	1.4	1.1	1.1	0.7	*	*
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.6	3.0	2.2	2.1	1.4	1.0	1.0	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*

White, female

2003.....	1.3	1.2	0.9	0.8	0.5	*	0.5	0.5	0.7	*
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.8	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2003.....	13.7	10.2	6.8	4.6	3.1	2.4	*	*	*	*
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.6	7.2	4.2	2.7	2.7	2.6	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*

Black, male

2003.....	24.3	19.1	11.9	9.1	5.2	*	*	*	*	*
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.5	17.2	13.9	7.4	5.6	5.0	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Black, female

2003.....	4.4	2.5	2.5	*	*	*	*	*	*	*
2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2003.....	2.6	2.5	1.9	1.5	1.3	1.5	1.5	1.8	1.8	1.7
2002.....	2.6	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 4/.....	4.9	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000.....	2.9	2.4	1.8	1.7	1.4	1.4	1.4	1.6	2.1	1.7
1999.....	2.9	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8

All races 3/, male

2003.....	3.6	3.3	2.9	2.2	1.9	2.1	2.0	2.7	2.6	2.5
2002.....	3.4	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 4/.....	7.0	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000.....	3.9	3.4	2.7	2.6	2.1	2.1	2.1	2.0	3.0	2.4
1999.....	4.0	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6

All races 3/, female

2003.....	1.6	1.6	1.0	0.9	0.7	1.0	1.0	1.1	1.3	1.4
2002.....	1.8	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 4/.....	2.9	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000.....	1.8	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999.....	1.8	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2003.....	1.9	1.7	1.4	1.2	1.0	1.2	1.1	1.5	1.5	1.6
2002.....	1.9	1.8	1.3	1.4	1.3	1.2	1.2	1.0	1.3	1.4
2001 4/.....	4.2	3.7	2.9	2.6	1.8	1.7	1.4	1.4	1.5	1.8
2000.....	1.9	1.7	1.4	1.3	1.0	1.0	1.2	1.4	1.6	1.5
1999.....	2.0	1.7	1.5	1.3	1.0	1.4	1.2	1.4	1.4	1.5

White, male

2003.....	2.4	2.3	2.0	1.5	1.5	1.6	1.5	2.2	2.2	2.3
2002.....	2.6	2.5	1.6	1.9	1.9	1.7	1.8	1.2	1.5	2.0
2001 4/.....	6.0	5.3	4.2	3.9	2.5	2.4	1.8	1.7	2.0	2.3
2000.....	2.6	2.1	1.9	1.9	1.5	1.3	1.7	1.7	2.2	2.0
1999.....	2.8	2.4	2.1	1.9	1.5	2.1	1.3	1.8	2.2	1.9

White, female

2003.....	1.3	1.2	0.9	0.8	0.5	0.9	0.8	1.0	1.1	1.2
2002.....	1.2	1.2	0.9	0.9	0.7	0.7	0.7	0.8	1.2	1.2
2001 4/.....	2.3	2.0	1.6	1.4	1.1	1.1	1.1	1.2	1.2	1.6
2000.....	1.2	1.2	0.9	0.6	0.6	0.7	0.7	1.1	1.3	1.3
1999.....	1.2	0.9	1.0	0.8	0.6	0.8	1.0	1.0	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2003.....	7.7	7.5	5.9	4.5	3.6	3.3	4.4	4.8	5.1	*
2002.....	7.5	8.0	5.9	4.6	3.5	3.1	3.8	5.7	*	*
2001 4/.....	10.1	9.7	7.2	5.6	4.6	4.8	4.0	4.7	*	*
2000.....	9.2	7.5	5.5	5.0	4.8	5.2	4.1	4.1	7.1	*
1999.....	9.0	6.5	5.3	4.3	3.2	4.4	3.4	5.2	*	*

Black, male

2003.....	12.5	10.9	9.9	8.1	6.2	6.2	6.9	9.1	*	*
2002.....	9.7	13.4	10.3	8.5	6.9	*	7.3	10.1	*	*
2001 4/.....	13.9	14.7	11.3	8.8	6.9	6.9	7.1	*	*	*
2000.....	12.6	12.4	9.9	9.3	8.0	9.5	6.4	*	*	*
1999.....	13.6	10.0	9.0	7.6	4.5	8.6	*	*	*	*

Black, female

2003.....	3.5	4.5	2.5	*	*	*	*	*	*	*
2002.....	5.5	3.4	2.1	*	*	*	*	*	*	*
2001 4/.....	6.8	5.3	3.7	3.1	*	*	*	*	*	*
2000.....	6.2	3.3	2.0	*	*	*	*	*	*	*
1999.....	5.0	3.5	2.2	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2003.....	3.9	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.0	2.7	1.7	1.2	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

All races 3/, male

2003.....	4.9	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	0.9	1.1	1.6
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1

All races 3/, female

2003.....	2.9	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.2	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	4.0	3.6	2.5	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.6	0.5	0.6	0.8	1.2
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4

White, male

2003.....	5.0	4.5	3.2	1.8	1.0	0.7	0.7	0.9	1.1	1.5
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	3.9	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1

White, female

2003.....	3.1	2.8	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.8	1.7	1.2	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2003.....	3.8	3.5	2.5	1.5	1.0	0.7	0.6	0.6	0.8	1.1
2002.....	3.3	3.2	2.2	1.3	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	3.1	2.9	1.9	1.0	0.6	0.5	0.4	0.6	1.0	1.0
2000.....	2.9	2.7	1.6	1.1	0.7	0.5	0.5	0.5	0.7	1.3
1999.....	2.8	2.6	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3

All races 3/, male

2003.....	4.8	4.5	3.3	1.8	1.1	0.8	0.6	0.7	*	1.4
2002.....	4.1	4.0	2.8	1.7	1.3	0.9	0.8	0.7	1.0	2.2
2001.....	4.2	4.1	2.4	1.4	0.7	0.8	0.6	0.7	1.3	*
2000.....	4.0	3.8	2.2	1.5	0.9	0.8	0.7	0.8	*	*
1999.....	3.8	3.5	2.0	1.4	0.9	0.9	*	0.9	*	1.8

All races 3/, female

2003.....	2.8	2.5	1.8	1.1	0.8	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.5	1.7	0.9	0.7	0.6	0.4	0.5	0.8	1.1
2001.....	2.0	1.7	1.5	0.6	0.5	*	*	0.6	0.7	0.8
2000.....	1.9	1.6	1.1	0.7	0.4	*	*	*	*	1.2
1999.....	1.8	1.7	1.0	0.6	0.5	0.5	*	*	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	3.9	3.5	2.5	1.4	0.9	0.7	0.5	0.5	0.8	1.2
2002.....	3.4	3.2	2.1	1.3	0.9	0.7	0.6	0.5	0.8	1.4
2001.....	3.1	2.9	1.8	1.0	0.6	0.5	0.4	0.6	0.9	1.0
2000.....	2.8	2.6	1.6	0.9	0.6	0.5	0.5	0.5	0.7	1.2
1999.....	2.6	2.4	1.3	0.9	0.7	0.7	0.4	0.6	0.8	1.3

White, male

2003.....	4.8	4.4	3.1	1.7	0.9	0.6	*	0.7	*	*
2002.....	4.2	3.9	2.4	1.7	1.2	0.8	0.7	*	*	2.3
2001.....	4.1	3.9	2.2	1.3	0.6	0.7	*	0.7	1.4	*
2000.....	3.7	3.5	2.0	1.2	0.8	0.7	0.6	0.8	*	*
1999.....	3.6	3.1	1.7	1.2	0.9	0.9	*	0.8	*	*

White, female

2003.....	3.0	2.7	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.5	1.9	1.0	0.7	0.6	0.5	*	0.7	1.1
2001.....	2.1	1.8	1.5	0.7	0.6	*	*	0.6	*	0.8
2000.....	1.8	1.6	1.1	0.7	0.4	*	*	*	*	1.1
1999.....	1.7	1.7	1.0	0.7	0.6	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2003.....	0.5	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	1.0	1.2	2.1	2.8	4.4	5.8	8.1	13.5
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7

All races 3/, male

2003.....	0.5	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.1	2.8	5.4	6.8	10.2	15.2
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5

All races 3/, female

2003.....	0.4	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.2	2.0	2.7	3.6	5.0	6.8	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

White, both sexes

2003.....	0.4	0.6	0.6	1.0	1.6	2.1	3.6	5.1	7.4	11.0
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.0
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1

White, male

2003.....	0.6	0.6	0.5	1.2	1.6	2.3	4.1	5.5	9.1	12.8
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.2	1.8	2.6	5.1	6.6	10.4	15.1
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6

White, female

2003.....	0.3	0.5	0.6	0.8	1.6	2.0	3.1	4.7	6.3	10.2
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	1.0	1.8	2.7	3.4	4.5	6.5	12.1
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2003.....	*	1.3	1.6	2.5	3.0	3.6	5.8	9.5	7.6	15.2
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.2	1.7	2.6	4.7	4.5	7.8	10.5	10.9	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1

Black, male

2003.....	*	*	2.1	2.8	*	*	*	*	*	*
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.8	6.0	8.9	10.0	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Black, female

2003.....	*	*	*	*	3.0	3.8	6.4	10.6	7.7	15.4
2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	7.0	10.8	11.4	22.7
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

All races 3/, both sexes

2003.....	10.6	16.0	18.9	18.3	18.8	16.8	13.2	10.3	7.6	4.1
2002 6/.....	10.6	15.6	18.2	18.2	18.6	16.5	14.4	10.9	7.2	4.4
2001 6/.....	10.6	16.5	17.9	17.6	18.9	17.3	13.7	11.0	7.5	5.0
2000 6/.....	11.0	16.1	17.0	18.4	19.6	17.4	14.5	11.2	8.0	5.3
1999 6/.....	10.9	16.0	16.9	18.1	19.3	17.6	14.1	11.9	8.5	5.5

All races 3/, male

2003.....	15.2	24.5	29.9	29.1	30.6	27.6	22.8	18.0	15.0	7.8
2002 6/.....	15.3	23.7	29.2	29.2	31.0	27.9	23.8	18.7	13.4	9.7
2001 6/.....	15.3	25.4	28.0	28.7	31.0	28.5	23.8	19.4	13.8	11.6
2000 6/.....	16.2	25.3	27.6	29.3	32.0	29.4	25.2	20.7	14.7	11.3
1999 6/.....	16.6	25.2	27.4	29.7	31.5	29.9	24.5	21.2	16.6	11.8

All races 3/, female

2003.....	6.0	7.7	8.3	8.1	8.1	7.5	5.4	4.9	3.1	2.5
2002 6/.....	6.1	7.7	7.7	7.9	7.2	6.7	6.9	5.3	3.4	2.1
2001 6/.....	6.0	7.9	8.3	7.2	8.0	7.6	5.7	5.1	3.8	2.3
2000 6/.....	5.8	7.2	7.1	8.4	8.4	7.3	6.0	4.4	3.9	2.7
1999 6/.....	5.3	7.1	6.8	7.3	8.3	7.0	6.0	5.5	3.8	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

White, both sexes

2003.....	10.8	16.2	18.7	18.2	18.2	17.0	13.2	10.6	7.6	4.2
2002 6/.....	10.9	15.8	18.2	17.7	17.9	16.5	14.2	10.9	7.1	4.3
2001 6/.....	10.4	16.5	17.1	16.8	18.6	16.7	13.8	10.9	7.7	4.8
2000 6/.....	10.7	15.6	16.4	17.5	18.9	17.1	14.4	11.2	7.9	5.3
1999 6/.....	10.5	15.3	16.2	17.1	18.5	17.0	13.6	11.6	8.3	5.3

White, male

2003.....	15.6	24.7	29.3	28.9	29.2	27.6	22.7	18.2	14.6	7.8
2002 6/.....	15.8	24.0	29.1	28.4	29.8	27.5	23.0	18.7	13.2	9.3
2001 6/.....	14.9	25.4	26.8	27.0	30.2	27.1	23.5	19.2	14.0	11.1
2000 6/.....	15.8	24.8	26.4	27.9	30.7	28.6	24.7	20.4	14.7	11.1
1999 6/.....	16.1	24.2	26.3	27.7	30.4	28.4	23.4	20.4	16.4	11.4

White, female

2003.....	6.0	7.8	8.3	8.0	8.0	7.5	5.4	5.1	3.3	2.5
2002 6/.....	6.1	7.7	7.4	7.5	6.9	6.8	7.1	5.3	3.4	2.2
2001 6/.....	5.9	7.6	7.6	7.0	8.0	7.6	5.8	5.1	3.9	2.2
2000 6/.....	5.5	6.5	6.6	7.7	8.1	7.1	6.0	4.4	3.7	2.8
1999 6/.....	4.8	6.5	6.3	7.0	7.6	7.0	5.9	5.4	3.6	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

Black, both sexes

2003.....	8.7	15.1	22.6	22.4	26.7	19.0	14.3	9.0	8.6	*
2002 6/.....	9.0	15.8	21.8	25.3	25.3	18.7	17.7	12.5	9.2	*
2001 6/.....	12.2	18.4	27.4	26.9	24.6	25.3	14.7	13.5	7.3	*
2000 6/.....	12.9	21.9	25.0	27.0	26.4	22.9	17.5	13.6	8.5	6.3
1999 6/.....	12.9	22.8	24.1	29.3	28.9	24.9	21.2	16.4	11.6	7.7

Black, male

2003.....	12.6	24.7	39.4	37.2	49.2	33.0	27.6	17.3	21.4	*
2002 6/.....	12.5	24.4	35.3	42.0	45.5	36.4	35.1	23.4	19.1	*
2001 6/.....	18.2	28.9	43.9	47.9	44.3	47.8	29.6	26.0	*	*
2000 6/.....	19.2	33.9	42.2	45.6	46.1	41.1	34.4	27.3	16.6	*
1999 6/.....	18.9	35.9	39.9	52.9	48.6	48.3	41.4	31.3	22.7	*

Black, female

2003.....	5.3	6.7	8.4	10.3	9.2	8.6	5.3	*	*	*
2002 6/.....	5.8	8.4	10.3	11.6	9.4	5.6	6.0	6.0	*	*
2001 6/.....	6.9	9.2	13.4	9.6	9.1	8.6	4.7	6.1	*	*
2000 6/.....	7.4	11.8	10.9	12.6	11.8	9.4	5.4	*	*	*
1999 6/.....	7.6	11.6	10.8	10.1	13.5	7.5	7.9	7.4	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 years
years	years	years	years	years	years	years	years	years	and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

All races 3/, both sexes

2003.....	22.8	22.7	17.0	9.8	5.7	4.5	3.7	3.9	4.6	6.0
2002 6/.....	21.8	20.9	14.5	8.2	5.1	3.8	3.3	3.6	4.1	5.7
2001 6/.....	18.5	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.6	5.1
2000 6/.....	17.4	15.9	10.5	6.0	3.7	2.6	2.8	3.2	4.0	5.6
1999 6/.....	17.4	15.5	9.3	5.8	3.8	3.2	2.9	3.6	4.2	4.8

All races 3/, male

2003.....	29.1	29.2	21.7	11.5	6.6	5.2	3.9	4.7	5.3	7.8
2002 6/.....	28.0	26.4	19.1	10.0	5.7	4.1	3.4	3.9	4.7	7.3
2001 6/.....	24.6	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.9	5.7
2000 6/.....	23.6	22.2	14.1	6.8	3.9	2.9	2.8	3.5	4.2	7.1
1999 6/.....	24.3	21.9	12.2	7.1	4.7	3.7	3.1	3.5	4.4	6.2

All races 3/, female

2003.....	16.6	16.5	12.4	8.2	4.9	4.0	3.4	3.3	4.2	5.2
2002 6/.....	15.8	15.6	10.2	6.6	4.6	3.6	3.3	3.3	3.8	4.9
2001 6/.....	12.6	11.6	8.3	5.1	3.9	2.6	2.6	3.2	3.4	4.9
2000 6/.....	11.3	9.9	7.1	5.2	3.5	2.3	2.8	2.9	3.9	4.9
1999 6/.....	10.6	9.2	6.6	4.6	2.9	2.7	2.7	3.7	4.0	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 years
years	years	years	years	years	years	years	years	years	and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

White, both sexes

2003.....	23.9	23.1	16.5	9.5	5.6	4.4	3.8	4.2	4.9	6.3
2002 6/.....	22.6	20.8	13.8	7.9	4.9	3.9	3.5	3.7	4.3	5.8
2001 6/.....	18.7	16.9	10.8	6.1	3.9	3.1	3.0	3.5	3.7	5.1
2000 6/.....	17.1	15.1	9.6	5.7	3.7	2.6	2.9	3.4	4.1	5.5
1999 6/.....	16.9	14.5	8.6	5.5	3.5	3.1	2.9	3.8	4.4	4.8

White, male

2003.....	30.3	29.2	20.4	10.7	6.2	4.7	3.9	4.9	5.5	7.8
2002 6/.....	29.0	25.9	17.3	9.0	4.9	3.8	3.4	4.0	4.6	7.6
2001 6/.....	24.8	22.3	13.3	6.9	3.6	3.6	3.2	3.6	4.1	5.6
2000 6/.....	23.1	20.5	12.3	6.0	3.6	2.7	2.7	3.8	4.3	6.9
1999 6/.....	23.5	20.0	10.6	6.2	4.0	3.5	3.1	3.5	4.6	6.4

White, female

2003.....	17.3	17.0	12.7	8.5	5.2	4.2	3.6	3.6	4.4	5.6
2002 6/.....	16.3	15.8	10.4	7.0	4.9	3.9	3.5	3.5	4.2	4.9
2001 6/.....	12.7	11.6	8.4	5.3	4.2	2.6	2.8	3.5	3.5	4.9
2000 6/.....	11.1	9.7	6.9	5.5	3.7	2.4	3.0	3.1	4.0	4.9
1999 6/.....	10.1	9.1	6.6	4.8	3.2	2.8	2.7	4.0	4.2	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Black, both sexes

2003.....	23.6	28.2	25.4	14.6	7.8	6.4	3.2	*	*	*
2002 6/.....	24.1	28.2	25.0	13.6	8.9	3.7	*	*	*	*
2001 6/.....	23.3	28.0	21.7	10.2	7.1	3.5	3.0	*	*	6.1
2000 6/.....	23.9	26.4	21.7	9.8	4.9	3.5	*	*	*	7.5
1999 6/.....	25.7	27.8	18.4	9.8	6.5	3.9	3.0	*	*	*

Black, male

2003.....	31.1	39.3	38.5	23.3	13.1	11.6	*	*	*	*
2002 6/.....	31.1	38.7	39.9	22.8	16.0	7.0	*	*	*	*
2001 6/.....	31.7	43.3	34.4	17.1	12.6	5.3	*	*	*	*
2000 6/.....	33.8	41.8	35.6	16.7	8.2	5.7	*	*	*	*
1999 6/.....	36.8	44.3	30.2	17.6	12.1	6.5	*	*	*	*

Black, female

2003.....	17.0	18.5	14.3	7.5	3.6	*	*	*	*	*
2002 6/.....	17.9	19.1	12.4	6.1	3.3	*	*	*	*	*
2001 6/.....	15.9	14.8	10.8	4.6	*	*	*	*	*	*
2000 6/.....	15.1	13.4	10.3	4.4	*	*	*	*	*	*
1999 6/.....	15.9	13.5	8.5	3.3	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

All races 3/, both sexes

2003.....	11.5	11.4	10.9	10.4	9.6	9.7	11.3	12.9	14.2	12.5
2002.....	11.9	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001.....	11.3	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000.....	11.7	10.8	10.5	9.7	9.2	9.8	11.6	13.4	14.9	14.0
1999.....	10.9	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5

All races 3/, male

2003.....	18.9	18.9	18.6	17.8	17.6	18.5	22.5	27.8	33.8	37.8
2002.....	19.7	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001.....	18.6	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000.....	19.2	18.1	17.7	16.7	16.4	18.1	23.3	29.0	36.0	42.5
1999.....	17.7	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3

All races 3/, female

2003.....	4.2	4.1	3.5	3.4	2.4	2.1	2.3	2.3	2.2	1.3
2002.....	4.2	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001.....	4.1	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000.....	4.2	3.7	3.6	3.2	2.8	2.7	2.4	2.2	1.9	1.7
1999.....	4.2	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

White, both sexes

2003.....	11.2	11.6	11.5	11.1	10.4	10.5	12.1	13.9	15.1	13.5
2002.....	11.6	11.3	10.8	11.5	10.2	11.0	12.8	14.4	16.7	13.4
2001.....	11.0	10.9	10.9	10.9	10.5	11.3	12.1	14.6	16.1	13.8
2000.....	11.3	10.8	10.7	10.1	9.9	10.5	12.3	14.5	15.8	15.1
1999.....	10.6	10.6	10.0	10.4	9.8	10.7	12.9	14.9	15.4	14.5

White, male

2003.....	18.1	18.9	19.4	18.9	18.9	20.0	24.1	29.7	36.0	41.0
2002.....	19.1	18.4	18.5	19.7	18.1	20.3	25.8	31.2	40.3	41.7
2001.....	18.0	18.2	18.3	18.6	18.0	20.9	24.1	31.8	39.2	43.3
2000.....	18.5	17.9	17.9	17.2	17.2	19.2	24.4	31.3	38.3	46.2
1999.....	17.1	17.6	16.7	17.5	17.7	20.1	25.8	32.6	38.2	46.8

White, female

2003.....	4.2	4.4	3.6	3.7	2.5	2.2	2.3	2.4	2.2	1.3
2002.....	4.1	4.2	3.3	3.6	2.9	2.8	2.1	2.3	2.2	1.2
2001.....	4.0	3.7	3.7	3.6	3.6	2.9	2.3	2.4	2.2	1.4
2000.....	4.1	3.7	3.8	3.3	3.1	3.0	2.6	2.3	2.0	1.8
1999.....	4.1	3.8	3.4	3.6	2.5	2.5	2.7	2.4	1.9	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Black, both sexes

2003.....	16.6	12.4	10.2	7.3	6.3	5.9	6.2	6.0	6.3	*
2002.....	16.6	13.0	10.3	7.3	7.0	5.1	6.7	8.8	*	*
2001.....	15.4	11.3	9.6	8.1	5.5	5.9	7.4	6.9	8.4	*
2000.....	16.5	13.4	11.0	7.5	5.8	5.9	7.7	4.6	7.1	*
1999.....	15.6	13.6	8.6	8.5	7.9	6.4	6.9	8.4	6.1	*

Black, male

2003.....	29.4	23.4	18.2	13.7	11.8	11.3	12.2	13.2	14.8	*
2002.....	29.1	24.1	18.1	14.4	12.3	10.2	14.7	20.7	*	*
2001.....	26.8	20.4	17.4	16.3	10.4	12.2	15.8	16.3	20.9	*
2000.....	28.5	24.1	20.6	14.3	11.7	12.2	17.2	9.1	16.6	*
1999.....	26.9	24.4	15.2	17.3	15.5	12.7	15.6	19.5	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
Black, female										
2003.....	5.4	2.8	3.4	*	*	*	*	*	*	*
2002.....	5.6	3.3	3.7	*	*	*	*	*	*	*
2001.....	5.4	3.5	2.9	*	*	*	*	*	*	*
2000.....	5.9	4.4	3.1	*	*	*	*	*	*	*
1999.....	5.6	4.3	3.1	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
1/ Figures for age not stated included in "All ages" but not distributed among age groups.
2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
3/ Includes races other than white and black.
4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
5/ Included in selected categories.
6/ Figures may differ from those previously shown as they are based on a modified list of causes; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."